



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C12N 15/12, 15/11, C07K 14/52, C12N 15/62, C07K 16/22, G01N 33/68, A61K 38/18	A2	(11) International Publication Number: WO 00/34474 (43) International Publication Date: 15 June 2000 (15.06.00)
(21) International Application Number: PCT/US99/28968 (22) International Filing Date: 7 December 1999 (07.12.99) (30) Priority Data: 09/207,120 7 December 1998 (07.12.98) US 60/142,576 6 July 1999 (06.07.99) US 60/161,653 21 October 1999 (21.10.99) US 60/165,255 12 November 1999 (12.11.99) US (71) Applicant: ZYMOGENETICS, INC. [US/US]; 1201 Eastlake Avenue East, Seattle, WA 98102 (US). (72) Inventors: GAO, Zeren; 9502 179th Place Northeast #3, Redmond, WA 98052 (US). HART, Charles, E.; 20114 183rd Place Northeast, Woodinville, WA 98072 (US). PIDDINGTON, Christopher, S.; 2838 Irongate Place, Thousand Oaks, CA 91362 (US). SHEPPARD, Paul, O.; 13532 278th Drive Northeast, Granite Falls, WA 98252 (US). SHOEMAKER, Kimberly, E.; 4634 Somerset Avenue Southeast, Bellevue, WA 98006 (US). GILBERTSON, Debra, G.; 4025 Northeast 195th Place, Seattle, WA 98155 (US). WEST, James, W.; 7539 19th Northeast, Seattle, WA 98115 (US).	(74) Agent: PARKER, Gary, E.; ZymoGenetics, Inc., 1201 Eastlake Avenue East, Seattle, WA 98102 (US). (81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>Without international search report and to be republished upon receipt of that report.</i>	
(54) Title: GROWTH FACTOR HOMOLOG ZVEGF3 (57) Abstract <p>Polypeptide growth factors, methods of making them, polynucleotides encoding them, antibodies to them, and methods of using them are disclosed. The polypeptides comprises an amino acid segment that is at least 90% identical to residues 46-163 of SEQ ID NO:2 or residues 235-345 of SEQ ID NO:2. Multimers of the polypeptides are also disclosed. The polypeptides, multimeric proteins, and polynucleotides can be used in the study and regulation of cell and tissue development, as components of cell culture media, and as diagnostic agents.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

Description

GROWTH FACTOR HOMOLOG ZVEGF3

5

BACKGROUND OF THE INVENTION

In multicellular animals, cell growth, differentiation, and migration are controlled by polypeptide growth factors. These growth factors play a role in both normal development and pathogenesis, including the development of solid tumors.

Polypeptide growth factors influence cellular events by binding to cell-surface receptors, many of which are tyrosine kinases. Binding initiates a chain of signalling events within the cell, which ultimately results in phenotypic changes, such as cell division, protease production, and cell migration.

Growth factors can be classified into families on the basis of structural similarities. One such family, the PDGF (platelet derived growth factor) family, is characterized by a dimeric structure stabilized by disulfide bonds. This family includes PDGF, the placental growth factors (PIGFs), and the vascular endothelial growth factors (VEGFs). The individual polypeptide chains of these proteins form characteristic higher-order structures having a bow tie-like configuration about a cystine knot, formed by disulfide bonding between pairs of cysteine residues. Hydrophobic interactions between loops contribute to the dimerization of the two monomers. See, Daopin et al., *Science* 257:369, 1992; Laphorn et al., *Nature* 369:455, 1994. Members of this family are active as both homodimers and heterodimers. See, for example, Heldin et al., *EMBO J.* 7:1387-1393, 1988; Cao et al., *J. Biol. Chem.* 271:3154-3162, 1996. The cystine knot motif and bow tie fold are also characteristic of the growth factors transforming growth factor-beta (TGF- β) and nerve growth factor (NGF), and the glycoprotein hormones. Although their amino acid sequences are quite divergent, these proteins all contain the six conserved cysteine residues of the cystine knot.

Four vascular endothelial growth factors have been identified: VEGF, also known as vascular permeability factor (Dvorak et al., *Am. J. Pathol.* 146:1029-1039, 1995); VEGF-B (Olofsson et al., *Proc. Natl. Acad. Sci. USA* 93:2567-2581, 1996; Hayward et al., WIPO Publication WO 96/27007); VEGF-
5 C (Joukov et al., *EMBO J.* 15:290-298, 1996); and VEGF-D (Oliviero, WO 97/12972; Achen et al., WO 98/07832). Five VEGF polypeptides (121, 145, 165, 189, and 206 amino acids) arise from alternative splicing of the VEGF mRNA.

VEGFs stimulate the development of vasculature through a
10 process known as angiogenesis, wherein vascular endothelial cells re-enter the cell cycle, degrade underlying basement membrane, and migrate to form new capillary sprouts. These cells then differentiate, and mature vessels are formed. This process of growth and differentiation is regulated by a balance of pro-angiogenic and anti-angiogenic factors. Angiogenesis is central to normal
15 formation and repair of tissue, occurring in embryo development and wound healing. Angiogenesis is also a factor in the development of certain diseases, including solid tumors, rheumatoid arthritis, diabetic retinopathy, macular degeneration, and atherosclerosis.

A number of proteins from vertebrates and invertebrates have
20 been identified as influencing neural development. Among those molecules are members of the neuropilin family and the semaphorin/collapsin family. Neuronal cell outgrowths, known as processes, grow away from the cell body to form synaptic connections. Long, thin processes that carry information away from the cell body are called axons, and short, thicker processes which carry
25 information to and from the cell body are called dendrites. Axons and dendrites are collectively referred to as neurites. Neurites are extended by means of growth cones, the growing tips of neurites, which are highly motile and are ultimately responsible for increasing and extending the neuronal network in the body.

30 Three receptors for VEGF have been identified: KDR/Flk-1 (Matthews et al., *Proc. Natl. Acad. Sci. USA* 88:9026-9030, 1991), Flt-1 (de

Vries et al., *Science* 255:989-991, 1992), and neuropilin-1 (Soker et al., *Cell* 92:735-745, 1998). Neuropilin-1 is a cell-surface glycoprotein that was initially identified in *Xenopus* tadpole nervous tissues, then in chicken, mouse, and human. The primary structure of neuropilin-1 is highly conserved among these vertebrate species. Neuropilin-1 has been demonstrated to be a receptor for various members of the semaphorin family including semaphorin III (Kolodkin et al., *Cell* 90:753-762, 1997), Sema E and Sema IV (Chen et al., *Neuron* 19:547-559, 1997). A variety of activities have been associated with the binding of neuropilin-1 to its ligands. For example, binding of semaphorin III to neuropilin-1 can induce neuronal growth cone collapse and repulsion of neurites *in vitro* (Kitsukawa et al., *Neuron* 19: 995-1005, 1997).

In mice, neuropilin-1 is expressed in the cardiovascular system, nervous system, and limbs at particular developmental stages. Chimeric mice over-expressing neuropilin-1 were found to be embryonic lethal (Kitsukawa et al., *Development* 121:4309-4318, 1995). The chimeric embryos exhibited several morphological abnormalities, including excess capillaries and blood vessels, dilation of blood vessels, malformed hearts, ectopic sprouting and defasciculation of nerve fibers, and extra digits. All of these abnormalities occurred in the organs in which neuropilin-1 is expressed in normal development. Mice lacking the neuropilin-1 gene have severe cardiovascular abnormalities, including impairment of vascular network formation in the central and peripheral nervous systems (Takashima et al., American Heart Association 1998 Meeting, Abstract # 3178).

Neuropilin-1 has been identified as a cell-surface receptor for VEGF (Soker et al., *ibid.*), and displays selective binding activity for VEGF₁₆₅ over VEGF₁₂₁. It has been shown to be expressed on vascular endothelial cells and tumor cells *in vitro*. When neuropilin-1 is co-expressed in cells with KDR, neuropilin-1 enhances the binding of VEGF₁₆₅ to KDR and VEGF₁₆₅-mediated chemotaxis. Conversely, inhibition of VEGF₁₆₅ binding to neuropilin-1 inhibits its binding to KDR and its mitogenic activity for endothelial cells (Soker, et al., *ibid.*). Neuropilin-1 is also a receptor for PIGF-2 (Migdal et al., *J. Biol. Chem.*

273: 22272-22278, 1998). A second semaphorin receptor, neuropilin-2, exhibits homology with neuropilin-1 but has differs in binding specificity (Chen et al. *Neuron* 19: 547-559, 1997).

Semaphorins are a large family of molecules which share the
5 defining semaphorin domain of approximately 500 amino acids. This family can be subdivided into multiple subfamilies that contain both secreted and membrane-bound proteins. Select members of these subfamilies, class III (SemD) and class IV (SemD), form homodimers linked by disulfide bridges. In the case of SemD, there is additional proteolytic processing that creates a 65-
10 kDa isoform that lacks the 33-kDa carboxyl-terminal sequence. Dimerization is believed to be important for functional activity (Klostermann et al., *J. Biol. Chem.* 273:7326-7331, 1998). Collapsin-1, the first identified vertebrate member of the semaphorin family of axon guidance proteins, has also been shown to form covalent dimers, with dimerization necessary for collapse activity
15 (Koppel et al., *J. Biol. Chem.* 273:15708-15713, 1998).

Semaphorin III has been associated *in vitro* with regulating growth cone collapse and chemorepulsion of neurites. *In vivo* it has also been shown to be required for correct sensory afferent innervation and other aspects of development, including skeletal and cardiac defects (Fehar et al., *Nature*
20 383:525-528, 1996). Other members of the semaphorin family have been shown to be associated with other forms of biology. The human semaphorin E gene is expressed in rheumatoid synovial cells and is thought to play an immunosuppressive role via inhibition of cytokines (Mangasser-Stephan et al., *Biochem. Biophys. Res. Comm.* 234:153-156, 1997). CD100, a leukocyte
25 semaphorin, promotes B-cell aggregation and differentiation (Hall et al., *Proc. Natl. Acad. Sci. USA* 93:11780-11785, 1996). CD100 has also been shown to be expressed in many T-cell lymphomas and may be a marker of malignant T-cell neoplasms (Dorfman et al., *Am. J. Pathol.* 153:255-262, 1998). Semaphorin homologues have also been identified in DNA viruses (Lang,
30 *Genomics* 51:340-350, 1998) and in poxvirus (Comeau, et al. *Immunity* 8:473-482, 1998). Transcription of the mouse semaphorin gene, M-semaH,

correlates with metastatic ability of mouse tumor cell lines (Christensen et al., *Cancer Res.* 58:1238-1244, 1998).

The role of growth factors, other regulatory molecules, and their receptors in controlling cellular processes makes them likely candidates and targets for therapeutic intervention. Platelet-derived growth factor, for example, has been disclosed for the treatment of periodontal disease (U.S. Patent No. 5,124,316), gastrointestinal ulcers (U.S. Patent No. 5,234,908), and dermal ulcers (Robson et al., *Lancet* 339:23-25, 1992; Steed et al., *J. Vasc. Surg.* 21:71-81, 1995). Inhibition of PDGF receptor activity has been shown to reduce intimal hyperplasia in injured baboon arteries (Giese et al., Restenosis Summit VIII, Poster Session #23, 1996; U.S. Patent No. 5,620,687). Vascular endothelial growth factors (VEGFs) have been shown to promote the growth of blood vessels in ischemic limbs (Isner et al., *The Lancet* 348:370-374, 1996), and have been proposed for use as wound-healing agents, for treatment of periodontal disease, for promoting endothelialization in vascular graft surgery, and for promoting collateral circulation following myocardial infarction (WIPO Publication No. WO 95/24473; U.S. Patent No. 5,219,739). VEGFs are also useful for promoting the growth of vascular endothelial cells in culture. A soluble VEGF receptor (soluble flt-1) has been found to block binding of VEGF to cell-surface receptors and to inhibit the growth of vascular tissue in vitro (*Biotechnology News* 16(17):5-6, 1996).

In view of the proven clinical utility of hormones, there are needs in the art for additional such molecules for use as therapeutic agents, diagnostic agents, and research tools and reagents. These and other needs are addressed by the present invention.

DESCRIPTION OF THE INVENTION

The present invention provides an isolated polypeptide of at least 15 amino acid residues comprising an epitope-bearing portion of a protein of SEQ ID NO:2. Within certain embodiments, the polypeptide comprises a segment that is at least 90% identical to residues 46-163 of SEQ ID NO:2 or

residues 235-345 of SEQ ID NO:2. Within additional embodiments, the polypeptide is selected from the group consisting of residues 15-163 of SEQ ID NO:2, residues 46-163 of SEQ ID NO:2, residues 15-170 of SEQ ID NO:2, residues 46-170 of SEQ ID NO:2, residues 15-234 of SEQ ID NO:2, residues 46-234 of SEQ ID NO:2, residues 15-229 amide of SEQ ID NO:2, residues 15-230 of SEQ ID NO:2, residues 15-345 of SEQ ID NO:2, residues 46-345 of SEQ ID NO:2, residues 235-345 of SEQ ID NO:2, and residues 226-345 of SEQ ID NO:2.

The invention also provides an isolated polypeptide comprising a sequence of amino acids of the formula $R1_x-R2_y-R3_z$, wherein R1 comprises a polypeptide of from 100 to 120 residues in length that is at least 90% identical to residues 46-163 of SEQ ID NO:2, and comprises a sequence motif C[KR]Y[DNE][WYF]X{11,15}G[KR][WYF]C (SEQ ID NO:4) corresponding to residues 104-124 of SEQ ID NO:2; R2 is a polypeptide at least 90% identical to residues 164-234 of SEQ ID NO:2; R3 is a polypeptide at least 90% identical in amino acid sequence to residues 235-345 of SEQ ID NO:2 and comprises cysteine residues at positions corresponding to residues 250, 280, 284, 296, 335, and 337 of SEQ ID NO:2, a glycine residue at a position corresponding to residue 282 of SEQ ID NO:2, and a sequence motif CX{18,33}CXGXCX{6,33}CX{20,40}CXC (SEQ ID NO:3) corresponding to residues 250-337 of SEQ ID NO:2; and each of x, y, and z is individually 0 or 1, subject to the limitations that at least one of x and z is 1, and, if x and z are each 1, then y is 1. There are thus provided isolated polypeptides of the above formula wherein (a) $x=1$, (b) $z=1$, and (c) $x=1$ and $z=1$. Within certain embodiments, $x=1$ and R1 is at least 90% identical to residues 18-163 of SEQ ID NO:2. Within related embodiments, $x=1$ and R1 comprises residues 46-163 of SEQ ID NO:2. Within other embodiments, $z=1$ and R3 comprises residues 235-345 of SEQ ID NO:2. Within additional embodiments, $x=1$, $z=1$, and the polypeptide comprises residues 46-229 of SEQ ID NO:2, residues 164-345 of SEQ ID NO:2, or residues 46-345 of SEQ ID NO:2. The isolated polypeptide may further comprise cysteine residues at positions corresponding to residues

286, 287, 291, and 294 of SEQ ID NO:2. Within other embodiments, the isolated polypeptide further comprises an affinity tag. Within a related embodiment, the isolated polypeptide comprises an immunoglobulin constant domain.

5 The present invention also provides an isolated protein comprising a first polypeptide operably linked to a second polypeptide, wherein the first polypeptide comprises a sequence of amino acids of the formula $R1_x-R2_y-R3_z$ as disclosed above. The protein modulates cell proliferation, differentiation, metabolism, or migration. Within one embodiment, the protein is a heterodimer. Within related embodiments, the second polypeptide is selected from the group consisting of VEGF, VEGF-B, VEGF-C, VEGF-D, zveg4, PIGF, PDGF-A, and PDGF-B. Within another embodiment, the protein is a homodimer.

15 There is also provided an isolated protein produced by a method comprising the steps of (a) culturing a host cell containing an expression vector comprising the following operably linked elements: a transcription promoter; a DNA segment encoding a polypeptide selected from the group consisting of (i) residues 46-345 of SEQ ID NO:2, (ii) residues 46-234 of SEQ ID NO:2, (iii) residues 164-345 of SEQ ID NO:2, and (iv) residues 235-345 of SEQ ID NO:2; and a transcription terminator, under conditions whereby the DNA segment is expressed; and (b) recovering from the cell the protein product of expression of the DNA construct.

25 Within another aspect of the invention there is provided an isolated polynucleotide of up to approximately 4 kb in length, wherein said polynucleotide encodes a polypeptide as disclosed above. Within one embodiment of the invention, the polynucleotide is DNA.

30 Within a further aspect of the invention there is provided an expression vector comprising the following operably linked elements: (a) a transcription promoter; (b) a DNA polynucleotide as disclosed above; and (c) a transcription terminator. The vector may further comprise a secretory signal sequence operably linked to the DNA polynucleotide.

Also provided by the invention is a cultured cell into which has been introduced an expression vector as disclosed above, wherein the cell expresses the polypeptide encoded by the DNA segment. The cultured cell can be used within a method of producing a polypeptide, the method comprising culturing the cell and recovering the expressed polypeptide.

The proteins provided herein can be combined with a pharmaceutically acceptable vehicle to provide a pharmaceutical composition.

The invention also provides an antibody that specifically binds to an epitope of a polypeptide as disclosed above. Antibodies of the invention include, *inter alia*, monoclonal antibodies and single chain antibodies, and may be linked to a reporter molecule.

The invention further provides a method for detecting a genetic abnormality in a patient, comprising the steps of (a) obtaining a genetic sample from a patient, (b) incubating the genetic sample with a polynucleotide comprising at least 14 contiguous nucleotides of SEQ ID NO:1 or the complement of SEQ ID NO:1, under conditions wherein said polynucleotide will hybridize to complementary polynucleotide sequence, to produce a first reaction product, and (c) comparing the first reaction product to a control reaction product, wherein a difference between the first reaction product and the control reaction product is indicative of a genetic abnormality in the patient.

Within an additional aspect, the invention provides a method of stimulating the growth of fibroblasts or smooth muscle cells comprising applying to the cells an effective amount of a protein as disclosed above.

Within another aspect the invention provides methods for modulating cell growth or other cellular processes. Within one embodiment there is provided a method of stimulating the growth of fibroblasts or smooth muscle cells comprising applying to the cells an effective amount of a protein as disclosed above. Within another embodiment there is provided a method of activating a cell-surface PDGF alpha receptor, comprising exposing a cell comprising a cell-surface PDGF alpha receptor to a polypeptide or protein as disclosed above, whereby the polypeptide or protein binds to and activates the

receptor. Within a further embodiment there is provided a method of inhibiting a PDGF alpha receptor-mediated cellular process, comprising exposing a cell comprising a cell-surface PDGF alpha receptor to a compound that inhibits binding of a polypeptide or protein as disclosed above to the receptor.

5 Within a further method of the invention there is provided a method of inhibiting zveg3 activity in a mammal comprising administering to the mammal an effective amount of a zveg3 antagonist. Within certain embodiments the antagonist is an antibody, a receptor, a ligand-binding receptor fragment, or a receptor IgG-Fc fusion protein.

10 Within another aspect of the invention there is provided an isolated, antisense polynucleotide that is the complement of a polynucleotide encoding a polypeptide comprising a sequence of amino acids of the formula $R1_x-R2_y-R3_z$, wherein R1 comprises a polypeptide of from 100 to 120 residues in length that is at least 90% identical to residues 46-163 of SEQ ID NO:2, and
15 comprises a sequence motif C[KR]Y[DNE][WYF]X{11,15}G[KR][WYF]C (SEQ ID NO:4) corresponding to residues 104-124 of SEQ ID NO:2; R2 is a polypeptide at least 90% identical to residues 164-234 of SEQ ID NO:2; R3 is a polypeptide at least 90% identical in amino acid sequence to residues 235-345 of SEQ ID NO:2 and comprises cysteine residues at positions corresponding to
20 residues 250, 280, 284, 296, 335, and 337 of SEQ ID NO:2, a glycine residue at a position corresponding to residue 282 of SEQ ID NO:2, and a sequence motif CX{18,33}CXGXCX{6,33}CX{20,40}CXC (SEQ ID NO:3) corresponding to residues 250-337 of SEQ ID NO:2; and each of x, y, and z is individually 0 or 1, subject to the limitations that at least one of x and z is 1, and, if x and z are
25 each 1, then y is 1. Within one embodiment the antisense polynucleotide further comprises operably linked transcription promoter and terminator sequences. The antisense polynucleotide can be used within a method of inhibiting zveg3 production in a cell comprising administering to the cell the antisense polynucleotide.

These and other aspects of the invention will become evident upon reference to the following detailed description of the invention and the attached drawings. In the drawings:

Fig. 1 is a Hopp/Woods hydrophilicity profile of the amino acid sequence shown in SEQ ID NO:2. The profile is based on a sliding six-residue window. Buried G, S, and T residues and exposed H, Y, and W residues were ignored. These residues are indicated in the figure by lower case letters.

Fig. 2 is an illustration of the vector pHB12-8 for use in expressing cDNAs in transgenic animals.

Fig. 3 is an illustration of the vector pZMP6/zveg3.

Fig. 4 is an illustration of the vector pZMP11/zv3GF-otPA.

Fig. 5 shows the results of a receptor binding assay for zveg3.

Fig. 6 is an alignment of human (SEQ ID NO:2) and mouse (SEQ ID NO:43) amino acid sequences.

The term "affinity tag" is used herein to denote a polypeptide segment that can be attached to a second polypeptide to provide for purification or detection of the second polypeptide or provide sites for attachment of the second polypeptide to a substrate. In principal, any peptide or protein for which an antibody or other specific binding agent is available can be used as an affinity tag. Affinity tags include a poly-histidine tract, protein A (Nilsson et al., *EMBO J.* 4:1075, 1985; Nilsson et al., *Methods Enzymol.* 198:3, 1991), glutathione S transferase (Smith and Johnson, *Gene* 67:31, 1988), maltose binding protein (Kellerman and Ferenci, *Methods Enzymol.* 90:459-463, 1982; Guan et al., *Gene* 67:21-30, 1987), Glu-Glu affinity tag (Grussenmeyer et al., *Proc. Natl. Acad. Sci. USA* 82:7952-4, 1985; see SEQ ID NO:5), substance P, Flag™ peptide (Hopp et al., *Biotechnology* 6:1204-10, 1988), streptavidin binding peptide, thioredoxin, ubiquitin, cellulose binding protein, T7 polymerase, or other antigenic epitope or binding domain. See, in general, Ford et al., *Protein Expression and Purification* 2: 95-107, 1991. DNAs encoding affinity tags and other reagents are available from commercial

suppliers (e.g., Pharmacia Biotech, Piscataway, NJ; New England Biolabs, Beverly, MA; and Eastman Kodak, New Haven, CT).

The term "allelic variant" is used herein to denote any of two or more alternative forms of a gene occupying the same chromosomal locus.

5 Allelic variation arises naturally through mutation, and may result in phenotypic polymorphism within populations. Gene mutations can be silent (no change in the encoded polypeptide) or may encode polypeptides having altered amino acid sequence. The term allelic variant is also used herein to denote a protein encoded by an allelic variant of a gene.

10 The terms "amino-terminal" and "carboxyl-terminal" are used herein to denote positions within polypeptides. Where the context allows, these terms are used with reference to a particular sequence or portion of a polypeptide to denote proximity or relative position. For example, a certain sequence positioned carboxyl-terminal to a reference sequence within a
15 polypeptide is located proximal to the carboxyl terminus of the reference sequence, but is not necessarily at the carboxyl terminus of the complete polypeptide.

A "beta-strand-like region" is a region of a protein characterized by certain combinations of the polypeptide backbone dihedral angles phi (ϕ) and psi (ψ). Regions wherein ϕ is less than -60° and ψ is greater than 90° are
20 beta-strand-like. Those skilled in the art will recognize that the limits of a β -strand are somewhat imprecise and may vary with the criteria used to define them. See, for example, Richardson and Richardson in Fasman, ed., Prediction of Protein Structure and the Principles of Protein Conformation, Plenum Press, New York, 1989; and Lesk, Protein Architecture: A Practical Approach, Oxford University Press, New York, 1991.
25

A "complement" of a polynucleotide molecule is a polynucleotide molecule having a complementary base sequence and reverse orientation as compared to a reference sequence. For example, the sequence 5'
30 ATGCACGGG 3' is complementary to 5' CCCGTGCAT 3'.

"Corresponding to", when used in reference to a nucleotide or amino acid sequence, indicates the position in a second sequence that aligns with the reference position when two sequences are optimally aligned.

The term "degenerate nucleotide sequence" denotes a sequence
5 of nucleotides that includes one or more degenerate codons (as compared to a reference polynucleotide molecule that encodes a polypeptide). Degenerate codons contain different triplets of nucleotides, but encode the same amino acid residue (i.e., GAU and GAC triplets each encode Asp).

The term "expression vector" is used to denote a DNA molecule,
10 linear or circular, that comprises a segment encoding a polypeptide of interest operably linked to additional segments that provide for its transcription. Such additional segments include promoter and terminator sequences, and may also include one or more origins of replication, one or more selectable markers, an enhancer, a polyadenylation signal, etc. Expression vectors are generally
15 derived from plasmid or viral DNA, or may contain elements of both.

The term "isolated", when applied to a polynucleotide, denotes that the polynucleotide has been removed from its natural genetic milieu and is thus free of other extraneous or unwanted coding sequences, and is in a form suitable for use within genetically engineered protein production systems.
20 Such isolated molecules are those that are separated from their natural environment and include cDNA and genomic clones. Isolated DNA molecules of the present invention are free of other genes with which they are ordinarily associated, but may include naturally occurring 5' and 3' untranslated regions such as promoters and terminators. The identification of associated regions
25 will be evident to one of ordinary skill in the art (see for example, Dynan and Tijan, *Nature* 316:774-78, 1985).

An "isolated" polypeptide or protein is a polypeptide or protein that is found in a condition other than its native environment, such as apart from blood and animal tissue. Within one form, the isolated polypeptide or
30 protein is substantially free of other polypeptides or proteins, particularly other polypeptides or proteins of animal origin. Polypeptides and proteins can be

provided in a highly purified form, i.e. greater than 95% pure or greater than 99% pure. When used in this context, the term "isolated" does not exclude the presence of the same polypeptide or protein in alternative physical forms, such as dimers or alternatively glycosylated or derivatized forms.

5 A "motif" is a series of amino acid positions in a protein sequence for which certain amino acid residues are required. A motif defines the set of possible residues at each such position.

 "Operably linked" means that two or more entities are joined together such that they function in concert for their intended purposes. When
10 referring to DNA segments, the phrase indicates, for example, that coding sequences are joined in the correct reading frame, and transcription initiates in the promoter and proceeds through the coding segment(s) to the terminator. When referring to polypeptides, "operably linked" includes both covalently (e.g., by disulfide bonding) and non-covalently (e.g., by hydrogen bonding,
15 hydrophobic interactions, or salt-bridge interactions) linked sequences, wherein the desired function(s) of the sequences are retained.

 The term "ortholog" denotes a polypeptide or protein obtained from one species that is the functional counterpart of a polypeptide or protein from a different species. Sequence differences among orthologs are the result
20 of speciation.

 A "PDGF alpha receptor-mediated cellular process" is a cellular process that occurs in response to activation of a PDGF alpha receptor. Such processes include, without limitation, cell division, chemotaxis, cell differentiation, and production or release of macromolecules.

25 A "polynucleotide" is a single- or double-stranded polymer of deoxyribonucleotide or ribonucleotide bases read from the 5' to the 3' end. Polynucleotides include RNA and DNA, and may be isolated from natural sources, synthesized *in vitro*, or prepared from a combination of natural and synthetic molecules. Sizes of polynucleotides are expressed as base pairs
30 (abbreviated "bp"), nucleotides ("nt"), or kilobases ("kb"). Where the context allows, the latter two terms may describe polynucleotides that are single-

stranded or double-stranded. When the term is applied to double-stranded molecules it is used to denote overall length and will be understood to be equivalent to the term "base pairs". It will be recognized by those skilled in the art that the two strands of a double-stranded polynucleotide may differ slightly in length and that the ends thereof may be staggered as a result of enzymatic cleavage; thus all nucleotides within a double-stranded polynucleotide molecule may not be paired. Such unpaired ends will in general not exceed 20 nt in length.

A "polypeptide" is a polymer of amino acid residues joined by peptide bonds, whether produced naturally or synthetically. Polypeptides of less than about 10 amino acid residues are commonly referred to as "peptides".

The term "promoter" is used herein for its art-recognized meaning to denote a portion of a gene containing DNA sequences that provide for the binding of RNA polymerase and initiation of transcription. Promoter sequences are commonly, but not always, found in the 5' non-coding regions of genes.

A "protein" is a macromolecule comprising one or more polypeptide chains. A protein may also comprise non-peptidic components, such as carbohydrate groups. Carbohydrates and other non-peptidic substituents may be added to a protein by the cell in which the protein is produced, and will vary with the type of cell. Proteins are defined herein in terms of their amino acid backbone structures; substituents such as carbohydrate groups are generally not specified, but may be present nonetheless.

A "secretory signal sequence" is a DNA sequence that encodes a polypeptide (a "secretory peptide") that, as a component of a larger polypeptide, directs the larger polypeptide through a secretory pathway of a cell in which it is synthesized. The larger polypeptide is commonly cleaved to remove the secretory peptide during transit through the secretory pathway.

A "segment" is a portion of a larger molecule (e.g., polynucleotide or polypeptide) having specified attributes. For example, a DNA segment

encoding a specified polypeptide is a portion of a longer DNA molecule, such as a plasmid or plasmid fragment, that, when read from the 5' to the 3' direction, encodes the sequence of amino acids of the specified polypeptide.

5 Molecular weights and lengths of polymers determined by imprecise analytical methods (e.g., gel electrophoresis) will be understood to be approximate values. When such a value is expressed as "about" X or "approximately" X, the stated value of X will be understood to be accurate to $\pm 20\%$.

10 All references cited herein are incorporated by reference in their entirety.

The present invention is based in part upon the discovery of a novel DNA molecule that encodes a polypeptide comprising a growth factor domain and a CUB domain. The growth factor domain is characterized by an arrangement of cysteine residues and beta strands that is characteristic of the
15 "cystine knot" structure of the PDGF family. The CUB domain shows sequence homology to CUB domains in the neuropilins (Takagi et al., *Neuron* 7:295-307, 1991; Soker et al., *ibid.*), human bone morphogenetic protein-1 (Wozney et al., *Science* 242:1528-1534, 1988), porcine seminal plasma protein and bovine acidic seminal fluid protein (Romero et al., *Nat. Struct. Biol.* 4:783-788, 1997),
20 and *X. laevis* tolloid-like protein (Lin et al., *Dev. Growth Differ.* 39:43-51, 1997). Analysis of the tissue distribution of the mRNA corresponding to this novel DNA showed that expression was widespread in adult human tissues, and that expression occurred up to day 15 in mouse embryo. The polypeptide has been designated "zveg3" in view of its homology to the VEGFs in the growth factor
25 domain.

Structural predictions based on the zveg3 sequence and its homology to other growth factors suggests that the polypeptide can form homomultimers or heteromultimers that act on tissues to control organ development by modulating cell proliferation, migration, differentiation, or
30 metabolism. Zveg3 heteromultimers may comprise a polypeptide from another member of the PDGF/VEGF family of proteins, including VEGF,

VEGF-B, VEGF-C, VEGF-D, zveg4 (SEQ ID NOS:36 and 37), PIGF (Maglione et al., *Proc. Natl. Acad. Sci. USA* 88:9267-9271, 1991), PDGF-A (Murray et al., U.S. Patent No. 4,899,919; Heldin et al., U.S. Patent No. 5,219,759), or PDGF-B (Chiu et al., *Cell* 37:123-129, 1984; Johnsson et al., *EMBO J.* 3:921-928, 1984). Members of this family of polypeptides regulate organ development and regeneration, post-developmental organ growth, and organ maintenance, as well as tissue maintenance and repair processes. These factors are also involved in pathological processes where therapeutic treatments are required, including cancer, rheumatoid arthritis, diabetic retinopathy, ischemic limb disease, peripheral vascular disease, myocardial ischemia, vascular intimal hyperplasia, atherosclerosis, and hemangioma formation. To treat these pathological conditions it will often be required to develop compounds to antagonize the members of the PDGF/VEGF family of proteins, or their respective receptors. This may include the development of neutralizing antibodies, small molecule antagonists, modified forms of the growth factors that maintain receptor binding activity but lack receptor activating activity, soluble receptors, or antisense or ribozyme molecules to block polypeptide production.

SEQ ID NO:2 is the sequence of a representative polypeptide of the present invention. Analysis of the amino acid sequence shown in SEQ ID NO:2 indicates that residues 1 to 14 form a secretory peptide. The CUB domain extends from residue 46 to residue 163. A propeptide-like sequence extends from residue 164 to residue 234, and includes two potential cleavage sites at its carboxyl terminus, a dibasic site at residues 231-232 and a target site for furin or a furin-like protease at residues 231-234. The growth factor domain extends from residue 235 to residue 345. Those skilled in the art will recognize that domain boundaries are somewhat imprecise and can be expected to vary by up to ± 5 residues from the specified positions. Potential proteolytic cleavage sites occur at residues 232 and 234. Processing of recombinant zveg3 produced in BHK cells has been found to occur between residues 225 and 226. Signal peptide cleavage is predicted to occur after

residue 14 (± 3 residues). This analysis suggests that the zveg3 polypeptide chain may be cleaved to produce a plurality of monomeric species as shown in Table 1. Cleavage after Arg-234 is expected to result in subsequent removal of residues 231-234, with possible conversion of Gly-230 to an amide.

- 5 Cleavage after Lys-232 is expected to result in subsequent removal of residue 231, again with possible conversion of Gly-230 to an amide. In addition, it may be advantageous to include up to seven residues of the interdomain region at the carboxyl terminus of the CUB domain. The interdomain region can be truncated at its amino terminus by a like amount. See Table 1.

10

Table 1

<u>Monomer</u>	<u>Residues (SEQ ID NO:2)</u>
Cub domain	15 - 163
	46 - 163
	15 - 170
	46 - 170
CUB domain + interdomain region	15 - 234
	46 - 234
	15 - 229 amide
	15 - 230
Cub domain + interdomain region + growth factor domain	15 - 345
	46 - 345
Growth factor domain	235 - 345
	226 - 345
Growth factor domain + interdomain region	164 - 345
	171 - 345

Also included within the present invention are polypeptides that are at least 90% identical or at least 95% identical to the polypeptides disclosed in Table 1, wherein these additional polypeptides retain certain characteristic sequence motifs as disclosed below.

Zvegf3 polypeptides are designated herein with a subscript indicating the amino acid residues. For example, the CUB domain polypeptides disclosed in Table 1 are designated "zvegf3₁₅₋₁₆₃", "zvegf3₄₆₋₁₆₃", "zvegf3₁₅₋₁₇₀", and "zvegf3₄₆₋₁₇₀".

Higher order structure of zveg3 polypeptides can be predicted by sequence alignment with known homologs and computer analysis using available software (e.g., the Insight II® viewer and homology modeling tools; MSI, San Diego, CA). Analysis of SEQ ID NO:2 predicts that the secondary structure of the growth factor domain is dominated by the cystine knot, which ties together variable beta strand-like regions and loops into a bow tie-like structure. Sequence alignment indicates that Cys residues within the growth factor domain at positions 250, 280, 284, 296, 335, and 337, and Gly 282 are highly conserved within the family. Further analysis suggests pairing (disulfide bond formation) of Cys residues 250 and 296, 280 and 335, and 284 and 337 to form the cystine knot. This arrangement of conserved residues can be represented by the formula $CX_{\{18,33\}}CXGX_{\{6,33\}}CX_{\{20,40\}}CXC$ (SEQ ID NO:3), wherein amino acid residues are represented by the conventional single-letter code, X is any amino acid residue, and {y,z} indicates a region of variable residues (X) from y to z residues in length. A consensus bow tie structure is formed as: amino terminus to cystine knot → beta strand-like region 1 → variable loop 1 → beta strand-like region 2 → cystine knot → beta strand-like region 3 → variable loop 2 → beta strand-like region 4 → cystine knot → beta strand-like region 5 → variable loop 3 → beta strand-like region 6 → cystine knot. Variable loops 1 and 2 form one side of the bow tie, with variable loop 3 forming the other side. The structure of the zveg3 growth factor domain appears to diverge from the consensus structure of other family members in loop 2 and beta strand-like regions 3 and 4, wherein all are abbreviated and essentially replaced by a cysteine cluster comprising residues 285 (Ala) through 295 (Gln), which includes Cys residues at positions 286, 287, 291, and 294 of SEQ ID NO:2. The approximate boundaries of the beta strand-like regions in SEQ ID NO:2 are: region 1, residues 251-259; region 2, residues 275-279; region 5, residues 297-301; region 6, residues 329-334. Loops separate regions 1 and 2, and regions 5 and 6.

The CUB domain of zveg3 is believed to form a beta barrel structure with nine distinct beta strand-like regions. These regions comprise

residues 48-51, 55-59, 72-78, 85-90, 92-94, 107-112, 119-123, 139-146, and 156-163 of SEQ ID NO:2. A multiple alignment of CUB domains of *Xenopus laevis* neuropilin precursor (Takagi et al., *ibid.*), human BMP-1 (Wozney et al., *ibid.*), and *X. laevis* tolloid-like protein (Lin et al., *ibid.*) indicates the presence of
 5 a conserved motif corresponding to residues 104-124 of SEQ ID NO:2. This motif is represented by the formula C[KR]Y[DNE][WYF]X{11,15}G[KR][WYF]C (SEQ ID NO:4), wherein square brackets indicate the allowable residues at a given position and X{y,z} is as defined above.

The proteins of the present invention include proteins comprising
 10 CUB domains homologous to the CUB domain of zveg3. These homologous domains are from 100 to 120 residues in length and comprise a motif of the sequence C[KR]Y[DNE][WYF]X{11,15}G[KR][WYF]C (SEQ ID NO:4) corresponding to residues 104-124 of SEQ ID NO:2. These homologous CUB domains are at least 90% identical to residues 46-163 of SEQ ID NO:2 or at
 15 least 95% identical to residues 46-163 of SEQ ID NO:2.

CUB domain-containing proteins of the present invention may further include a zveg3 interdomain region or homolog thereof. The interdomain region is at least 90% identical to residues 164 to 234 of SEQ ID NO:2.

20 Additional proteins of the present invention comprise the zveg3 growth factor domain or a homolog thereof. These proteins thus comprise a polypeptide segment that is at least 90% or 95% identical to residues 235-345 of SEQ ID NO:2, wherein the polypeptide segment comprises Cys residues at positions corresponding to residues 250, 280, 284, 296, 335, and 337 of SEQ
 25 ID NO:2; a glycine at a position corresponding to residue 284 of SEQ ID NO:2; and the sequence motif CX{18,33}CXGXCX{6,33}CX{20,40}CXC (SEQ ID NO:3) corresponding to residues 250-337 of SEQ ID NO:2.

Additional proteins comprising combinations of the CUB domain, interdomain region, and growth factor domain are shown above in Table 1. In
 30 each case, the invention also includes homologous proteins comprising homologous domains as disclosed above.

Structural analysis and homology predict that zveg3 polypeptides complex with a second polypeptide to form multimeric proteins. These proteins include homodimers and heterodimers. In the latter case, the second polypeptide can be a truncated or other variant zveg3 polypeptide or another polypeptide, such as a PIGF, PDGF-A, PDGF-B, VEGF, VEGF-B, VEGF-C, VEGF-D, or zveg4 polypeptide. Among the dimeric proteins within the present invention are dimers formed by non-covalent association (e.g., hydrophobic interactions) with a second subunit, either a second zveg3 polypeptide or other second subunit, or by covalent association stabilized by intermolecular disulfide bonds between cysteine residues of the component monomers. Within SEQ ID NO:2, the Cys residues at positions 274, 286, 287, 291, 294, and 339 may form intramolecular or intermolecular disulfide bonds. It is likely that residues 274 and 287 form interchain disulfide bonds. In homodimers, the component polypeptides (designated A and B) may be joined in an antiparallel arrangement with a pattern of interchain disulfide bonds A274--B287, A287--B274. The data further suggest that additional intrachain disulfides are formed by the pairing of Cys286 with Cys291, and Cys294 with Cys339, although some or all of these four residues may be involved in interchain pairing.

The present invention thus provides a variety of multimeric proteins comprising a zveg3 polypeptide as disclosed above. These zveg3 polypeptides include zveg3₁₅₋₂₃₄, zveg3₄₆₋₂₃₄, zveg3₁₅₋₂₂₉ amide, zveg3₁₅₋₂₃₀, zveg3₁₅₋₃₄₅, zveg3₄₆₋₃₄₅, and zveg3₂₃₅₋₃₄₅, as well as variants and derivatives of these polypeptides as disclosed herein.

Percent sequence identity is determined by conventional methods. See, for example, Altschul et al., *Bull. Math. Bio.* 48:603-616, 1986, and Henikoff and Henikoff, *Proc. Natl. Acad. Sci. USA* 89:10915-10919, 1992. Briefly, two amino acid sequences are aligned to optimize the alignment scores using a gap opening penalty of 10, a gap extension penalty of 1, and the "BLOSUM62" scoring matrix of Henikoff and Henikoff (*ibid.*) as shown in Table 2 (amino acids are indicated by the standard one-letter codes). The percent identity is then calculated as:

Total number of identical matches

x 100

5

[length of the longer sequence plus the
number of gaps introduced into the longer
sequence in order to align the two
sequences]

Table 2

	A	R	N	D	C	Q	E	G	H	I	L	K	M	F	P	S	T	W	Y	V
A	4																			
R	-1	5																		
N	-2	0	6																	
D	-2	-2	1	6																
C	0	-3	-3	-3	9															
Q	-1	1	0	0	-3	5														
E	-1	0	0	2	-4	2	5													
G	0	-2	0	-1	-3	-2	-2	6												
H	-2	0	1	-1	-3	0	0	-2	8											
I	-1	-3	-3	-3	-1	-3	-3	-4	-3	4										
L	-1	-2	-3	-4	-1	-2	-3	-4	-3	2	4									
K	-1	2	0	-1	-3	1	1	-2	-1	-3	-2	5								
M	-1	-1	-2	-3	-1	0	-2	-3	-2	1	2	-1	5							
F	-2	-3	-3	-3	-2	-3	-3	-3	-1	0	0	-3	0	6						
P	-1	-2	-2	-1	-3	-1	-1	-2	-2	-3	-3	-1	-2	-4	7					
S	1	-1	1	0	-1	0	0	0	-1	-2	-2	0	-1	-2	-1	4				
T	0	-1	0	-1	-1	-1	-1	-2	-2	-1	-1	-1	-1	-2	-1	1	5			
W	-3	-3	-4	-4	-2	-2	-3	-2	-2	-3	-2	-3	-1	1	-4	-3	-2	1	1	
Y	-2	-2	-2	-3	-2	-1	-2	-3	2	-1	-1	-2	-1	3	-3	-2	-2	2	7	
V	0	-3	-3	-3	-1	-2	-2	-3	-3	3	1	-2	1	-1	-2	-2	0	-3	-1	4

The level of identity between amino acid sequences can be determined using the "FASTA" similarity search algorithm disclosed by Pearson and Lipman (*Proc. Natl. Acad. Sci. USA* 85:2444, 1988) and by Pearson (*Meth. Enzymol.* 183:63, 1990). Briefly, FASTA first characterizes sequence similarity by identifying regions shared by the query sequence (e.g., SEQ ID NO:2) and a test sequence that have either the highest density of identities (if the ktup variable is 1) or pairs of identities (if ktup=2), without considering conservative amino acid substitutions, insertions, or deletions. The ten regions with the highest density of identities are then rescored by comparing the similarity of all paired amino acids using an amino acid substitution matrix, and the ends of the regions are "trimmed" to include only those residues that contribute to the highest score. If there are several regions with scores greater than the "cutoff" value (calculated by a predetermined formula based upon the length of the sequence and the ktup value), then the trimmed initial regions are examined to determine whether the regions can be joined to form an approximate alignment with gaps. Finally, the highest scoring regions of the two amino acid sequences are aligned using a modification of the Needleman-Wunsch-Sellers algorithm (Needleman and Wunsch, *J. Mol. Biol.* 48:444, 1970; Sellers, *SIAM J. Appl. Math.* 26:787, 1974), which allows for amino acid insertions and deletions. Preferred parameters for FASTA analysis are: ktup=1, gap opening penalty=10, gap extension penalty=1, and substitution matrix=BLOSUM62. These parameters can be introduced into a FASTA program by modifying the scoring matrix file ("SMATRIX"), as explained in Appendix 2 of Pearson, 1990 (*ibid.*).

FASTA can also be used to determine the sequence identity of nucleic acid molecules using a ratio as disclosed above. For nucleotide sequence comparisons, the ktup value can range between one to six, preferably from three to six, most preferably three, with other parameters set as default.

The present invention includes polypeptides having one or more conservative amino acid changes as compared with the amino acid sequence

of SEQ ID NO:2. The BLOSUM62 matrix (Table 2) is an amino acid substitution matrix derived from about 2,000 local multiple alignments of protein sequence segments, representing highly conserved regions of more than 500 groups of related proteins (Henikoff and Henikoff, *ibid.*). Thus, the BLOSUM62 substitution frequencies can be used to define conservative amino acid substitutions that may be introduced into the amino acid sequences of the present invention. As used herein, the term "conservative amino acid substitution" refers to a substitution represented by a BLOSUM62 value of greater than -1. For example, an amino acid substitution is conservative if the substitution is characterized by a BLOSUM62 value of 0, 1, 2, or 3. Preferred conservative amino acid substitutions are characterized by a BLOSUM62 value of at least one 1 (e.g., 1, 2 or 3), while more preferred conservative amino acid substitutions are characterized by a BLOSUM62 value of at least 2 (e.g., 2 or 3).

The proteins of the present invention can further comprise amino- or carboxyl-terminal extensions, such as an amino-terminal methionine residue, an amino or carboxyl-terminal cysteine residue to facilitate subsequent linking to maleimide-activated keyhole limpet hemocyanin, a small linker peptide of up to about 20-25 residues, or a polypeptide extension that facilitates purification (an affinity tag) as disclosed above. Two or more affinity tags may be used in combination. Polypeptides comprising affinity tags can further comprise a polypeptide linker and/or a proteolytic cleavage site between the zveg3 polypeptide and the affinity tag. Exemplary cleavage sites include thrombin cleavage sites and factor Xa cleavage sites.

The present invention further provides a variety of other polypeptide fusions and related multimeric proteins comprising one or more polypeptide fusions. For example, a zveg3 polypeptide can be prepared as a fusion to a dimerizing protein as disclosed in U.S. Patents Nos. 5,155,027 and 5,567,584. Exemplary dimerizing proteins in this regard include immunoglobulin constant region domains. Dimerization can also be stabilized by fusing a zveg3 polypeptide to a leucine zipper sequence (Riley et al.,

Protein Eng. 9:223-230, 1996; Mohamed et al., *J. Steroid Biochem. Mol. Biol.* 51:241-250, 1994). Immunoglobulin-zveg3 polypeptide fusions and leucine zipper fusions can be expressed in genetically engineered cells to produce a variety of multimeric zveg3 analogs. Auxiliary domains can be fused to zveg3 polypeptides to target them to specific cells, tissues, or macromolecules (e.g., collagen). For example, a zveg3 polypeptide or protein can be targeted to a predetermined cell type by fusing a zveg3 polypeptide to a ligand that specifically binds to a receptor on the surface of the target cell. In this way, polypeptides and proteins can be targeted for therapeutic or diagnostic purposes. A zveg3 polypeptide can be fused to two or more moieties, such as an affinity tag for purification and a targeting domain. Polypeptide fusions can also comprise one or more cleavage sites, particularly between domains. See, Tuan et al., *Connective Tissue Research* 34:1-9, 1996.

Polypeptide fusions of the present invention will generally contain not more than about 1,500 amino acid residues, often not more than about 1,200 residues, often not more than about 1,000 residues, and will in many cases be considerably smaller. For example, a zveg3 polypeptide of 331 residues (residues 15-345 of SEQ ID NO:2) can be fused to *E. coli* β -galactosidase (1,021 residues; see Casadaban et al., *J. Bacteriol.* 143:971-980, 1980), a 10-residue spacer, and a 4-residue factor Xa cleavage site to yield a polypeptide of 1366 residues. In a second example, residues 235-345 of SEQ ID NO:2 can be fused to maltose binding protein (approximately 370 residues), a 4-residue cleavage site, and a 6-residue polyhistidine tag.

A polypeptide comprising the zveg3 growth factor domain (e.g., zveg3₂₃₅₋₃₄₅ or zveg3₁₆₄₋₃₄₅) can be fused to a non-zveg3 CUB domain. Within a related embodiment of the invention, a zveg3 polypeptide comprising zveg3 growth factor and CUB domains is fused to a non-zveg3 CUB domain, such as a CUB-domain comprising neuropilin polypeptide.

The present invention further provides polypeptide fusions comprising the zveg3 CUB domain (e.g., zveg3₄₅₋₁₆₃). The CUB domain, with its homology to neuropilin-1, may be used to target zveg3 or other proteins

containing it to cells having cell-surface semaphorins, including endothelial cells, neuronal cells, lymphocytes, and tumor cells. The zveg3 CUB domain can thus be joined to other moieties, including polypeptides (e.g., other growth factors, antibodies, and enzymes) and non-peptidic moieties (e.g.,
5 radionuclides, contrast agents, and the like), to target them to cells expressing cell-surface semaphorins. The dibasic and furin-like sites between the CUB and growth factor domains of zveg3 may allow for proteolytic release of the growth factor domain or other moiety through existing local proteases within tissues, or by proteases added from exogenous sources. The release of the
10 targetted moiety may provide more localized biological effects.

Proteins comprising the wild-type zveg3 CUB domain and variants thereof may be used to modulate activities mediated by cell-surface semaphorins. While not wishing to be bound by theory, zveg3 may bind to semaphorins via its CUB domain. The observation that semaphorin III is
15 involved in vascular development suggests that members of the vascular growth factor family of proteins may also be involved, especially due to the co-binding activity of VEGF and semaphorin III to neuropilin-1. Zveg3 may thus be used to design agonists and antagonist of neuropilin-semaphorin interactions. For example, the zveg3 sequence disclosed herein provides a
20 starting point for the design of molecules that antagonize semaphorin-stimulated activities, including neurite growth, cardiovascular development, cartilage and limb development, and T and B-cell function. Additional applications include intervention in various pathologies, including rheumatoid arthritis, various forms of cancer, autoimmune disease, inflammation,
25 retinopathies, hemangiomas, ischemic events within tissues including the heart, kidney and peripheral arteries, neuropathies, acute nerve damage, and diseases of the central and peripheral nervous systems.

Amino acid sequence changes are made in zveg3 polypeptides so as to minimize disruption of higher order structure essential to biological
30 activity. In general, conservative amino acid changes are preferred. Changes in amino acid residues will be made so as not to disrupt the cystine knot and

"bow tie" arrangement of loops in the growth factor domain that is characteristic of the protein family. Conserved motifs will also be maintained. The effects of amino acid sequence changes can be predicted by computer modeling as disclosed above or determined by analysis of crystal structure (see, e.g.,
5 Laphorn et al., *ibid.*). A hydrophobicity profile of SEQ ID NO:2 is shown in Fig. 1. Those skilled in the art will recognize that this hydrophobicity will be taken into account when designing alterations in the amino acid sequence of a zveg3 polypeptide, so as not to disrupt the overall profile. Additional guidance in selecting amino acid substitutions is provided by the alignment of mouse and
10 human zveg3 sequences shown in Fig. 6.

The polypeptides of the present invention can also comprise non-naturally occurring amino acid residues. Non-naturally occurring amino acids include, without limitation, *trans*-3-methylproline, 2,4-methanoproline, *cis*-4-hydroxyproline, *trans*-4-hydroxyproline, *N*-methylglycine, *allo*-threonine,
15 methylthreonine, hydroxyethylcysteine, hydroxyethylhomocysteine, nitroglutamine, homoglutamine, pipecolic acid, *tert*-leucine, norvaline, 2-azaphenylalanine, 3-azaphenylalanine, 4-azaphenylalanine, and 4-fluorophenylalanine. Several methods are known in the art for incorporating non-naturally occurring amino acid residues into proteins. For example, an *in*
20 *vitro* system can be employed wherein nonsense mutations are suppressed using chemically aminoacylated suppressor tRNAs. Methods for synthesizing amino acids and aminoacylating tRNA are known in the art. Transcription and translation of plasmids containing nonsense mutations is carried out in a cell-free system comprising an *E. coli* S30 extract and commercially available
25 enzymes and other reagents. Proteins are purified by chromatography. See, for example, Robertson et al., *J. Am. Chem. Soc.* 113:2722, 1991; Ellman et al., *Methods Enzymol.* 202:301, 1991; Chung et al., *Science* 259:806-809, 1993; and Chung et al., *Proc. Natl. Acad. Sci. USA* 90:10145-10149, 1993). In a second method, translation is carried out in *Xenopus* oocytes by
30 microinjection of mutated mRNA and chemically aminoacylated suppressor tRNAs (Turcatti et al., *J. Biol. Chem.* 271:19991-19998, 1996). Within a third

method, *E. coli* cells are cultured in the absence of a natural amino acid that is to be replaced (e.g., phenylalanine) and in the presence of the desired non-naturally occurring amino acid(s) (e.g., 2-azaphenylalanine, 3-azaphenylalanine, 4-azaphenylalanine, or 4-fluorophenylalanine). The non-naturally occurring amino acid is incorporated into the protein in place of its natural counterpart. See, Koide et al., *Biochem.* 33:7470-7476, 1994. Naturally occurring amino acid residues can be converted to non-naturally occurring species by *in vitro* chemical modification. Chemical modification can be combined with site-directed mutagenesis to further expand the range of substitutions (Wynn and Richards, *Protein Sci.* 2:395-403, 1993).

Essential amino acids in the polypeptides of the present invention can be identified according to procedures known in the art, such as site-directed mutagenesis or alanine-scanning mutagenesis (Cunningham and Wells, *Science* 244, 1081-1085, 1989; Bass et al., *Proc. Natl. Acad. Sci. USA* 88:4498-4502, 1991). In the latter technique, single alanine mutations are introduced at every residue in the molecule, and the resultant mutant molecules are tested for biological activity of other properties to identify amino acid residues that are critical to the activity of the molecule.

Multiple amino acid substitutions can be made and tested using known methods of mutagenesis and screening, such as those disclosed by Reidhaar-Olson and Sauer (*Science* 241:53-57, 1988) or Bowie and Sauer (*Proc. Natl. Acad. Sci. USA* 86:2152-2156, 1989). Briefly, these authors disclose methods for simultaneously randomizing two or more positions in a polypeptide, selecting for functional polypeptide, and then sequencing the mutagenized polypeptides to determine the spectrum of allowable substitutions at each position. Other methods that can be used include phage display (e.g., Lowman et al., *Biochem.* 30:10832-10837, 1991; Ladner et al., U.S. Patent No. 5,223,409; Huse, WIPO Publication WO 92/06204) and region-directed mutagenesis (Derbyshire et al., *Gene* 46:145, 1986; Ner et al., *DNA* 7:127, 1988).

Variants of the disclosed zveg3 DNA and polypeptide sequences can be generated through DNA shuffling as disclosed by Stemmer, *Nature* 370:389-391, 1994 and Stemmer, *Proc. Natl. Acad. Sci. USA* 91:10747-10751, 1994. Briefly, variant genes are generated by *in vitro* homologous recombination by random fragmentation of a parent gene followed by reassembly using PCR, resulting in randomly introduced point mutations. This technique can be modified by using a family of parent genes, such as allelic variants or genes from different species, to introduce additional variability into the process. Selection or screening for the desired activity, followed by additional iterations of mutagenesis and assay provides for rapid "evolution" of sequences by selecting for desirable mutations while simultaneously selecting against detrimental changes.

Mutagenesis methods as disclosed above can be combined with high volume or high-throughput screening methods to detect biological activity of zveg3 variant polypeptides, in particular biological activity in modulating cell proliferation or cell differentiation. For example, mitogenesis assays that measure dye incorporation or ³H-thymidine incorporation can be carried out on large numbers of samples, as can cell-based assays that detect expression of a reporter gene (e.g., a luciferase gene). Mutagenesis of the CUB domain can be used to modulate its binding to members of the semaphorin family, including enhancing or inhibiting binding to selected family members. A modified spectrum of binding activity may be desirable for optimizing therapeutic and/or diagnostic utility of proteins comprising a zveg3 CUB domain. Direct binding utilizing labeled CUB protein can be used to monitor changes in CUB domain binding activity to selected semaphorin family members. Semaphorins of interest in this regard include isolated proteins, proteins present in cell membranes, and proteins present on cell-surfaces. The CUB domain can be labeled by a variety of methods including radiolabeling with isotopes, such as ¹²⁵I, conjugation to enzymes such as alkaline phosphatase or horseradish peroxidase, conjugation with biotin, and conjugation with various fluorescent markers including FITC. These and other assays are disclosed in more detail

below. Mutagenized DNA molecules that encode active zveg3 polypeptides can be recovered from the host cells and rapidly sequenced using modern equipment. These methods allow the rapid determination of the importance of individual amino acid residues in a polypeptide of interest, and can be applied
5 to polypeptides of unknown structure.

Using the methods discussed above, one of ordinary skill in the art can identify and/or prepare a variety of polypeptides that are homologous to the zveg3 polypeptides disclosed above in Table 1 and retain the biological properties of the wild-type protein. Such polypeptides can also include
10 additional polypeptide segments as generally disclosed above.

The present invention also provides polynucleotide molecules, including DNA and RNA molecules, that encode the zveg3 polypeptides disclosed above. The polynucleotides of the present invention include the sense strand; the anti-sense strand; and the DNA as double-stranded, having
15 both the sense and anti-sense strand annealed together by hydrogen bonds. A representative DNA sequences encoding zveg3 polypeptides is set forth in SEQ ID NO:1. Additional DNA sequences encoding zveg3 polypeptides can be readily generated by those of ordinary skill in the art based on the genetic code. Counterpart RNA sequences can be generated by substitution of U for
20 T.

Those skilled in the art will readily recognize that, in view of the degeneracy of the genetic code, considerable sequence variation is possible among polynucleotide molecules encoding zveg3 polypeptides. SEQ ID NO:6 is a degenerate DNA sequence that encompasses all DNAs that encode the
25 zveg3 polypeptide of SEQ ID NO: 2. Those skilled in the art will recognize that the degenerate sequence of SEQ ID NO:6 also provides all RNA sequences encoding SEQ ID NO:2 by substituting U for T. Thus, zveg3 polypeptide-encoding polynucleotides comprising nucleotides 1 - 1035, 1 - 489, 43 - 489, 136 - 489, 43 - 702, 136 - 702, 43 - 690, 43 - 1035, 136 - 1035, and 703 - 1035
30 of SEQ ID NO:6 and their RNA equivalents are contemplated by the present invention. Table 3 sets forth the one-letter codes used within SEQ ID NO:6 to

denote degenerate nucleotide positions. "Resolutions" are the nucleotides denoted by a code letter. "Complement" indicates the code for the complementary nucleotide(s). For example, the code Y denotes either C or T, and its complement R denotes A or G, A being complementary to T, and G being complementary to C.

Table 3

Nucleotide	Resolutions	Complement	Resolutions
A	A	T	T
C	C	G	G
G	G	C	C
T	T	A	A
R	A G	Y	C T
Y	C T	R	A G
M	A C	K	G T
K	G T	M	A C
S	C G	S	C G
W	A T	W	A T
H	A C T	D	A G T
B	C G T	V	A C G
V	A C G	B	C G T
D	A G T	H	A C T
N	A C G T	N	A C G T

The degenerate codons used in SEQ ID NO:6, encompassing all possible codons for a given amino acid, are set forth in Table 4, below.

Table 4

Amino Acid	One-Letter Code	Codons	Degenerate Codon
Cys	C	TGC TGT	TGY
Ser	S	AGC AGT TCA TCC TCG TCT	WSN
Thr	T	ACA ACC ACG ACT	CAN
Pro	P	CCA CCC CCG CCT	CCN
Ala	A	GCA GCC GCG GCT	GCN
Gly	G	GGA GGC GGG GGT	GGN
Asn	N	AAC AAT	AAY
Asp	D	GAC GAT	GAY
Glu	E	GAA GAG	GAR
Gln	Q	CAA CAG	CAR
His	H	CAC CAT	CAY
Arg	R	AGA AGG CGA CGC CGG CGT	MGN
Lys	K	AAA AAG	AAR
Met	M	ATG	ATG
Ile	I	ATA ATC ATT	ATH
Leu	L	CTA CTC CTG CTT TTA TTG	YTN
Val	V	GTA GTC GTG GTT	GTN
Phe	F	TTC TTT	TTY
Tyr	Y	TAC TAT	TAY
Trp	W	TGG	TGG
Ter	.	TAA TAG TGA	TRR
Asn Asp	B		RAY
Glu Gln	Z		SAR
Any	X		NNN
Gap	-	---	

One of ordinary skill in the art will appreciate that some ambiguity is introduced in determining a degenerate codon, representative of all possible

codons encoding each amino acid. For example, the degenerate codon for serine (WSN) can, in some circumstances, encode arginine (AGR), and the degenerate codon for arginine (MGN) can, in some circumstances, encode serine (AGY). A similar relationship exists between codons encoding
5 phenylalanine and leucine. Thus, some polynucleotides encompassed by the degenerate sequences may encode variant amino acid sequences, but one of ordinary skill in the art can easily identify such variant sequences by reference to the amino acid sequence of SEQ ID NO: 2. Variant sequences can be readily tested for functionality as described herein.

10 Within certain embodiments of the invention the isolated polynucleotides will hybridize to similar sized regions of SEQ ID NO:1, or a sequence complementary thereto, under stringent conditions. In general, stringent conditions are selected to be about 5°C lower than the thermal melting point (T_m) for the specific sequence at a defined ionic strength and pH.
15 The T_m is the temperature (under defined ionic strength and pH) at which 50% of the target sequence hybridizes to a perfectly matched probe. Typical stringent conditions are those in which the salt concentration is up to about 0.03 M at pH 7 and the temperature is at least about 60°C.

As previously noted, the isolated polynucleotides of the present
20 invention include DNA and RNA. Methods for preparing DNA and RNA are well known in the art. Complementary DNA (cDNA) clones are prepared from RNA that is isolated from a tissue or cell that produces large amounts of zveg3 RNA. Such tissues and cells are identified by Northern blotting (Thomas, *Proc. Natl. Acad. Sci. USA* 77:5201, 1980), and include thyroid, spinal cord, and
25 adrenal gland. Total RNA can be prepared using guanidine HCl extraction followed by isolation by centrifugation in a CsCl gradient (Chirgwin et al., *Biochemistry* 18:52-94, 1979). Poly (A)⁺ RNA is prepared from total RNA using the method of Aviv and Leder (*Proc. Natl. Acad. Sci. USA* 69:1408-1412, 1972). Complementary DNA (cDNA) is prepared from poly(A)⁺ RNA using
30 known methods. In the alternative, genomic DNA can be isolated. For some applications (e.g., expression in transgenic animals) it may be preferable to use

a genomic clone, or to modify a cDNA clone to include at least one genomic intron. Methods for identifying and isolating cDNA and genomic clones are well known and within the level of ordinary skill in the art, and include the use of the sequence disclosed herein, or parts thereof, for probing or priming a library.

- 5 Polynucleotides encoding zveg3 polypeptides are identified and isolated by, for example, hybridization or polymerase chain reaction ("PCR", Mullis, U.S. Patent 4,683,202). Expression libraries can be probed with antibodies to zveg3, receptor fragments, or other specific binding partners.

- Those skilled in the art will recognize that the sequences
10 disclosed in SEQ ID NOS:1 and 2 represent a single allele of human zveg3. Allelic variants of these sequences can be cloned by probing cDNA or genomic libraries from different individuals according to standard procedures. Alternatively spliced forms of zveg3 are also expected to exist.

- The zveg3 polynucleotide sequence disclosed herein can be
15 used to isolate polynucleotides encoding other zveg3 proteins. Such other polynucleotides include allelic variants, alternatively spliced cDNAs and counterpart polynucleotides from other species (orthologs). These orthologous polynucleotides can be used, *inter alia*, to prepare the respective orthologous proteins. Other species of interest include, but are not limited to, mammalian,
20 avian, amphibian, reptile, fish, insect and other vertebrate and invertebrate species. Of particular interest are zveg3 polynucleotides and proteins from other mammalian species, including non-human primate, murine, porcine, ovine, bovine, canine, feline, and equine polynucleotides and proteins. Non-human ztb1 polypeptides and polynucleotides, as well as antagonists thereof
25 and other related molecules, can be used, *inter alia*, in veterinary medicine. Orthologs of human zveg3 can be cloned using information and compositions provided by the present invention in combination with conventional cloning techniques. For example, a cDNA can be cloned using mRNA obtained from a tissue or cell type that expresses zveg3 as disclosed herein. Suitable sources
30 of mRNA can be identified by probing Northern blots with probes designed from the sequences disclosed herein. A library is then prepared from mRNA of

a positive tissue or cell line. A zveg3-encoding cDNA can then be isolated by a variety of methods, such as by probing with a complete or partial human cDNA or with one or more sets of degenerate probes based on the disclosed sequences. Hybridization will generally be done under low stringency
5 conditions, wherein washing is carried out in 1 x SSC with an initial wash at 40°C and with subsequent washes at 5°C higher intervals until background is suitably reduced. A cDNA can also be cloned using the polymerase chain reaction, or PCR (Mullis, U.S. Patent No. 4,683,202), using primers designed from the representative human zveg3 sequence disclosed herein. Within an
10 additional method, the cDNA library can be used to transform or transfect host cells, and expression of the cDNA of interest can be detected with an antibody to zveg3 polypeptide. Similar techniques can also be applied to the isolation of genomic clones.

For any zveg3 polypeptide, including variants and fusion
15 proteins, one of ordinary skill in the art can readily generate a fully degenerate polynucleotide sequence encoding that variant using the information set forth in Tables 3 and 4, above.

Conserved regions of zveg3, identified by alignment with sequences of other family members, can be used to identify related
20 polynucleotides and proteins. For instance, reverse transcription-polymerase chain reaction (RT-PCR) and other techniques known in the art can be used to amplify sequences encoding the conserved motifs present in zveg3 from RNA obtained from a variety of tissue sources. In particular, highly degenerate primers as shown below in Table 5 (designed from an alignment of zveg3 with
25 PDGF A and B chains, VEGF, VEGF-B, VEGF-C, and VEGF-D) are useful for cloning polynucleotides encoding homologous growth factor domains. Primers shown in Table 6, designed from an alignment of zveg3 with *X. laevis* neuropilin precursor, human BMP-1, and *X. laevis* tolloid-like protein, are useful for cloning polynucleotides encoding CUB domains. The primers of Tables 6
30 and 7 can thus be used to obtain additional polynucleotides encoding homologs of the zveg3 sequence of SEQ ID NO:1 and NO:2.

Table 5

zveg3 residues 279-284

5 degenerate: MGN TGY GGN GGN AAY TG (SEQ ID NO:7)
 consensus: MGN TGY DSN GGN WRY TG (SEQ ID NO:8)
 complement: CAR YWN CCN SHR CAN CK (SEQ ID NO:9)

zveg3 residues 270-275

10 degenerate: TTY TGG CCN GGN TGY YT (SEQ ID NO:10)
 consensus: NTN DDN CCN NSN TGY BT (SEQ ID NO:11)
 complement: AVR CAN SNN GGN HHN AN (SEQ ID NO:12)

zveg3 residues 332-337

15 degenerate CAY GAR GAR TGY GAY TG (SEQ ID NO:13)
 consensus: CAY NNN NVN TGY VVN TG (SEQ ID NO:14)
 complement: CAN BBR CAN BNN NNR TG (SEQ ID NO:15)

zveg3 residues 250-255

20 degenerate: TGY ACN CCN MGN AAY TT (SEQ ID NO:16)
 consensus: TGY HNN MCN MKN RMN DH (SEQ ID NO:17)
 complement: DHN KYN MKN GKN NDR CA (SEQ ID NO:18)

Table 6

zveg3 residues 104-109

consensus: TGY AAR TAY GAY TWY GT (SEQ ID NO:19)

complement: ACR WAR TCR TAY TTR CA (SEQ ID NO:20)

5

zveg3 residues 120-125

consensus: YWN GGN MRN TDB TGY GG (SEQ ID NO:21)

complement: CCR CAV HAN YKN CCN WR (SEQ ID NO:22)

10 zveg3 residues 63-68

consensus: TDB CCN MAN DVN TAY CC (SEQ ID NO:23)

complement: GGR TAN BHN TKN GGV HA (SEQ ID NO:24)

15 Zveg3 polynucleotide sequences disclosed herein can also be used as probes or primers to clone 5' non-coding regions of a zveg3 gene, including promoter sequences. These flanking sequences can be used to direct the expression of zveg3 and other recombinant proteins. In addition, 5' flanking sequences can be used as targetting sites for regulatory constructs to activate or increase expression of endogenous zveg3 genes as disclosed by

20 Treco et al., U.S. Patent No. 5,641,670.

The polynucleotides of the present invention can also be prepared by automated synthesis. The current method of choice is the phosphoramidite method. If chemically synthesized, double-stranded DNA is required, each complementary strand is made separately. The production of

25 short, double-stranded segments (60 to 80 bp) is technically straightforward and can be accomplished by synthesizing the complementary strands and then annealing them. Longer segments (typically >300 bp) are assembled in modular form from single-stranded fragments that are from 20 to 100 nucleotides in length. Automated synthesis of polynucleotides is within the

30 level of ordinary skill in the art, and suitable equipment and reagents are available from commercial suppliers. See, in general, Glick and Pasternak,

Molecular Biotechnology, Principles & Applications of Recombinant DNA, ASM Press, Washington, D.C., 1994; Itakura et al., *Ann. Rev. Biochem.* 53: 323-56, 1984; and Climie et al., *Proc. Natl. Acad. Sci. USA* 87:633-7, 1990.

The polypeptides of the present invention, including full-length
5 polypeptides, biologically active fragments, and fusion polypeptides can be produced in genetically engineered host cells according to conventional techniques. Suitable host cells are those cell types that can be transformed or transfected with exogenous DNA and grown in culture, and include bacteria, fungal cells, and cultured higher eukaryotic cells (including cultured cells of
10 multicellular organisms). Techniques for manipulating cloned DNA molecules and introducing exogenous DNA into a variety of host cells are disclosed by Sambrook et al., *Molecular Cloning: A Laboratory Manual*, 2nd ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1989, and Ausubel et al., eds., *Current Protocols in Molecular Biology*, Green and Wiley and Sons,
15 NY, 1993.

In general, a DNA sequence encoding a zveg3 polypeptide is operably linked to other genetic elements required for its expression, generally including a transcription promoter and terminator, within an expression vector. The vector will also commonly contain one or more selectable markers and one
20 or more origins of replication, although those skilled in the art will recognize that within certain systems selectable markers may be provided on separate vectors, and replication of the exogenous DNA may be provided by integration into the host cell genome. Selection of promoters, terminators, selectable markers, vectors and other elements is a matter of routine design within the
25 level of ordinary skill in the art. Many such elements are described in the literature and are available through commercial suppliers.

To direct a zveg3 polypeptide into the secretory pathway of a host cell, a secretory signal sequence (also known as a leader sequence, prepro sequence or pre sequence) is provided in the expression vector. The
30 secretory signal sequence may be that of zveg3, or may be derived from another secreted protein (e.g., t-PA; see, U.S. Patent No. 5,641,655) or

synthesized *de novo*. The secretory signal sequence is operably linked to the zveg3 DNA sequence, i.e., the two sequences are joined in the correct reading frame and positioned to direct the newly synthesized polypeptide into the secretory pathway of the host cell. Secretory signal sequences are
5 commonly positioned 5' to the DNA sequence encoding the polypeptide of interest, although certain signal sequences may be positioned elsewhere in the DNA sequence of interest (see, e.g., Welch et al., U.S. Patent No. 5,037,743; Holland et al., U.S. Patent No. 5,143,830).

Expression of zveg3 polypeptides via a host cell secretory
10 pathway is expected to result in the production of multimeric proteins. As noted above, such multimers include both homomultimers and heteromultimers, the latter including proteins comprising only zveg3 polypeptides and proteins including zveg3 and heterologous polypeptides. For example, a heteromultimer comprising a zveg3 polypeptide and a
15 polypeptide from a related family member (e.g., VEGF, VEGF-B, VEGF-C, VEGF-D, zveg4, PIGF, PDGF-A, or PDGF-B) can be produced by co-expression of the two polypeptides in a host cell. Sequences encoding these other family members are known. See, for example, Dvorak et al, *ibid.*; Olofsson et al, *ibid.*; Hayward et al., *ibid.*; Joukov et al., *ibid.*; Oliviero et al.,
20 *ibid.*; Achen et al., *ibid.*; Maglione et al., *ibid.*; Heldin et al., U.S. Patent No. 5,219,759; and Johnsson et al., *ibid.* If a mixture of proteins results from expression, individual species are isolated by conventional methods. Monomers, dimers, and higher order multimers are separated by, for example, size exclusion chromatography. Heteromultimers can be separated from
25 homomultimers by immunoaffinity chromatography using antibodies specific for individual dimers or by sequential immunoaffinity steps using antibodies specific for individual component polypeptides. See, in general, U.S. Patent No. 5,094,941. Multimers may also be assembled *in vitro* upon incubation of component polypeptides under suitable conditions. In general, *in vitro*
30 assembly will include incubating the protein mixture under denaturing and reducing conditions followed by refolding and reoxidation of the polypeptides to

from homodimers and heterodimers. Recovery and assembly of proteins expressed in bacterial cells is disclosed below.

Cultured mammalian cells are suitable hosts for use within the present invention. Methods for introducing exogenous DNA into mammalian host cells include calcium phosphate-mediated transfection (Wigler et al., *Cell* 5 14:725, 1978; Corsaro and Pearson, *Somatic Cell Genetics* 7:603, 1981; Graham and Van der Eb, *Virology* 52:456, 1973), electroporation (Neumann et al., *EMBO J.* 1:841-845, 1982), DEAE-dextran mediated transfection (Ausubel et al., *ibid.*), and liposome-mediated transfection (Felgner et al., *Proc. Natl. Acad. Sci. USA* 84:7413-7, 1987; Mackey et al., *Proc. Natl. Acad. Sci. USA* 10 85:8027-31, 1988; Hawley-Nelson et al., *Focus* 15:73, 1993; Ciccarone et al., *Focus* 15:80, 1993). The production of recombinant polypeptides in cultured mammalian cells is disclosed by, for example, Levinson et al., U.S. Patent No. 4,713,339; Hagen et al., U.S. Patent No. 4,784,950; Palmiter et al., U.S. Patent 15 No. 4,579,821; and Ringold, U.S. Patent No. 4,656,134. Suitable cultured mammalian cells include the COS-1 (ATCC No. CRL 1650), COS-7 (ATCC No. CRL 1651), BHK (ATCC No. CRL 1632), BHK 570 (ATCC No. CRL 10314), 293 (ATCC No. CRL 1573; Graham et al., *J. Gen. Virol.* 36:59-72, 1977) and Chinese hamster ovary (e.g. CHO-K1; ATCC No. CCL 61) cell lines. Additional 20 suitable cell lines are known in the art and available from public depositories such as the American Type Culture Collection, Rockville, Maryland. Strong transcription promoters include promoters from SV-40 or cytomegalovirus. See, e.g., U.S. Patent No. 4,956,288. Other suitable promoters include those from metallothionein genes (U.S. Patent Nos. 4,579,821 and 4,601,978) and 25 the adenovirus major late promoter. Expression vectors for use in mammalian cells include pZP-1 and pZP-9, which have been deposited with the American Type Culture Collection, Rockville, MD USA under accession numbers 98669 and 98668, respectively.

Drug selection is generally used to select for cultured mammalian 30 cells into which foreign DNA has been inserted. Such cells are commonly referred to as "transfectants". Cells that have been cultured in the presence of

the selective agent and are able to pass the gene of interest to their progeny are referred to as "stable transfectants." An exemplary selectable marker is a gene encoding resistance to the antibiotic neomycin. Selection is carried out in the presence of a neomycin-type drug, such as G-418 or the like. Selection systems can also be used to increase the expression level of the gene of interest, a process referred to as "amplification." Amplification is carried out by culturing transfectants in the presence of a low level of the selective agent and then increasing the amount of selective agent to select for cells that produce high levels of the products of the introduced genes. An exemplary amplifiable selectable marker is dihydrofolate reductase, which confers resistance to methotrexate. Other drug resistance genes (e.g. hygromycin resistance, multi-drug resistance, puromycin acetyltransferase) can also be used. Alternative markers that introduce an altered phenotype, such as green fluorescent protein, or cell surface proteins such as CD4, CD8, Class I MHC, placental alkaline phosphatase can be used to sort transfected cells from untransfected cells by such means as FACS sorting or magnetic bead separation technology.

Other higher eukaryotic cells can also be used as hosts, including insect cells, plant cells and avian cells. The use of *Agrobacterium rhizogenes* as a vector for expressing genes in plant cells has been reviewed by Sinkar et al., *J. Biosci. (Bangalore)* 11:47-58, 1987. Transformation of insect cells and production of foreign polypeptides therein is disclosed by Guarino et al., U.S. Patent No. 5,162,222 and WIPO publication WO 94/06463.

Insect cells can be infected with recombinant baculovirus, commonly derived from *Autographa californica* nuclear polyhedrosis virus (AcNPV). See, King and Possee, The Baculovirus Expression System: A Laboratory Guide, London, Chapman & Hall; O'Reilly et al., Baculovirus Expression Vectors: A Laboratory Manual, New York, Oxford University Press., 1994; and Richardson, Ed., Baculovirus Expression Protocols. Methods in Molecular Biology, Humana Press, Totowa, NJ, 1995. Recombinant baculovirus can also be produced through the use of a transposon-based system described by Luckow et al. (*J. Virol.* 67:4566-4579, 1993). This

system, which utilizes transfer vectors, is commercially available in kit form (Bac-to-Bac™ kit; Life Technologies, Rockville, MD). The transfer vector (e.g., pFastBac1™; Life Technologies) contains a Tn7 transposon to move the DNA encoding the protein of interest into a baculovirus genome maintained in *E. coli* as a large plasmid called a "bacmid." See, Hill-Perkins and Possee, *J. Gen. Virol.* 71:971-976, 1990; Bonning et al., *J. Gen. Virol.* 75:1551-1556, 1994; and Chazenbalk and Rapoport, *J. Biol. Chem.* 270:1543-1549, 1995. In addition, transfer vectors can include an in-frame fusion with DNA encoding a polypeptide extension or affinity tag as disclosed above. Using techniques known in the art, a transfer vector containing a zveg3-encoding sequence is transformed into *E. coli* host cells, and the cells are screened for bacmids which contain an interrupted lacZ gene indicative of recombinant baculovirus. The bacmid DNA containing the recombinant baculovirus genome is isolated, using common techniques, and used to transfect *Spodoptera frugiperda* cells, such as Sf9 cells. Recombinant virus that expresses zveg3 protein is subsequently produced. Recombinant viral stocks are made by methods commonly used the art.

For protein production, the recombinant virus is used to infect host cells, typically a cell line derived from the fall armyworm, *Spodoptera frugiperda* (e.g., Sf9 or Sf21 cells) or *Trichoplusia ni* (e.g., High Five™ cells; Invitrogen, Carlsbad, CA). See, in general, Glick and Pasternak, Molecular Biotechnology: Principles and Applications of Recombinant DNA, ASM Press, Washington, D.C., 1994. See also, U.S. Patent No. 5,300,435. Serum-free media are used to grow and maintain the cells. Suitable media formulations are known in the art and can be obtained from commercial suppliers. The cells are grown up from an inoculation density of approximately $2-5 \times 10^5$ cells to a density of $1-2 \times 10^6$ cells, at which time a recombinant viral stock is added at a multiplicity of infection (MOI) of 0.1 to 10, more typically near 3. Procedures used are generally described in available laboratory manuals (e.g., King and Possee, *ibid.*; O'Reilly et al., *ibid.*; Richardson, *ibid.*).

Fungal cells, including yeast cells, can also be used within the present invention. Yeast species of particular interest in this regard include *Saccharomyces cerevisiae*, *Pichia pastoris*, and *Pichia methanolica*. Methods for transforming *S. cerevisiae* cells with exogenous DNA and producing recombinant polypeptides therefrom are disclosed by, for example, Kawasaki, U.S. Patent No. 4,599,311; Kawasaki et al., U.S. Patent No. 4,931,373; Brake, U.S. Patent No. 4,870,008; Welch et al., U.S. Patent No. 5,037,743; and Murray et al., U.S. Patent No. 4,845,075. Transformed cells are selected by phenotype determined by the selectable marker, commonly drug resistance or the ability to grow in the absence of a particular nutrient (e.g., leucine). An exemplary vector system for use in *Saccharomyces cerevisiae* is the *POT1* vector system disclosed by Kawasaki et al. (U.S. Patent No. 4,931,373), which allows transformed cells to be selected by growth in glucose-containing media. Suitable promoters and terminators for use in yeast include those from glycolytic enzyme genes (see, e.g., Kawasaki, U.S. Patent No. 4,599,311; Kingsman et al., U.S. Patent No. 4,615,974; and Bitter, U.S. Patent No. 4,977,092) and alcohol dehydrogenase genes. See also U.S. Patents Nos. 4,990,446; 5,063,154; 5,139,936 and 4,661,454. Transformation systems for other yeasts, including *Hansenula polymorpha*, *Schizosaccharomyces pombe*, *Kluyveromyces lactis*, *Kluyveromyces fragilis*, *Ustilago maydis*, *Pichia pastoris*, *Pichia methanolica*, *Pichia guillermondii* and *Candida maltosa* are known in the art. See, for example, Gleeson et al., *J. Gen. Microbiol.* 132:3459-3465, 1986; Cregg, U.S. Patent No. 4,882,279; and Raymond et al., *Yeast* 14, 11-23, 1998. Aspergillus cells may be utilized according to the methods of McKnight et al., U.S. Patent No. 4,935,349. Methods for transforming *Acremonium chrysogenum* are disclosed by Sumino et al., U.S. Patent No. 5,162,228. Methods for transforming *Neurospora* are disclosed by Lambowitz, U.S. Patent No. 4,486,533. Production of recombinant proteins in *Pichia methanolica* is disclosed in U.S. Patents No. 5,716,808, 5,736,383, 5,854,039, and 5,888,768.

Prokaryotic host cells, including strains of the bacteria *Escherichia coli*, *Bacillus* and other genera are also useful host cells within the

present invention. Techniques for transforming these hosts and expressing foreign DNA sequences cloned therein are well known in the art (see, e.g., Sambrook et al., *ibid.*). When expressing a zveg3 polypeptide in bacteria such as *E. coli*, the polypeptide may be retained in the cytoplasm, typically as insoluble granules, or may be directed to the periplasmic space by a bacterial secretion sequence. In the former case, the cells are lysed, and the granules are recovered and denatured using, for example, guanidine isothiocyanate or urea. The denatured polypeptide can then be refolded and dimerized by diluting the denaturant, such as by dialysis against a solution of urea and a combination of reduced and oxidized glutathione, followed by dialysis against a buffered saline solution. In the alternative, the protein may be recovered from the cytoplasm in soluble form and isolated without the use of denaturants. The protein is recovered from the cell as an aqueous extract in, for example, phosphate buffered saline. To capture the protein of interest, the extract is applied directly to a chromatographic medium, such as an immobilized antibody or heparin-Sepharose column. Secreted polypeptides can be recovered from the periplasmic space in a soluble and functional form by disrupting the cells (by, for example, sonication or osmotic shock) to release the contents of the periplasmic space and recovering the protein, thereby obviating the need for denaturation and refolding.

Production of zveg3 fusion proteins in prokaryotic host cells is of particular interest. An exemplary such fusion protein comprises a zveg3 polypeptide fused to maltose binding protein. Such fusions may further comprise additional sequences, such as polyhistidine to provide for affinity purification of the polypeptide fusion. An enzymatic cleavage site (e.g., a thrombin cleavage site) may also be included to allow for separation of zveg3 and non-zveg3 components of the fusion.

Transformed or transfected host cells are cultured according to conventional procedures in a culture medium containing nutrients and other components required for the growth of the chosen host cells. A variety of suitable media, including defined media and complex media, are known in the

art and generally include a carbon source, a nitrogen source, essential amino acids, vitamins and minerals. Media may also contain such components as growth factors or serum, as required. The growth medium will generally select for cells containing the exogenously added DNA by, for example, drug
5 selection or deficiency in an essential nutrient which is complemented by the selectable marker carried on the expression vector or co-transfected into the host cell. *P. methanolica* cells, for example, are cultured in a medium comprising adequate sources of carbon, nitrogen and trace nutrients at a temperature of about 25°C to 35°C. Liquid cultures are provided with sufficient
10 aeration by conventional means, such as shaking of small flasks or sparging of fermentors.

Zveg3 polypeptides or fragments thereof can also be prepared through chemical synthesis according to methods known in the art, including exclusive solid phase synthesis, partial solid phase methods, fragment
15 condensation or classical solution synthesis. See, for example, Merrifield, *J. Am. Chem. Soc.* 85:2149, 1963; Stewart et al., Solid Phase Peptide Synthesis (2nd edition), Pierce Chemical Co., Rockford, IL, 1984; Bayer and Rapp, *Chem. Pept. Prot.* 3:3, 1986; and Atherton et al., Solid Phase Peptide Synthesis: A Practical Approach, IRL Press, Oxford, 1989.

20 Non-covalent complexes comprising a zveg3 polypeptide can be prepared by incubating a zveg3 polypeptide and a second polypeptide (e.g., a zveg3 polypeptide or another peptide of the PDGF/VEGF family) at near-physiological pH. In a typical reaction, polypeptides at a concentration of about 0.1-0.5 µg/µl are incubated at pH≈7.4 in a weak buffer (e.g., 0.01 M phosphate
25 or acetate buffer); sodium chloride may be included at a concentration of about 0.1 M. At 37°C the reaction is essentially complete with 4-24 hours. See, for example, Weintraub et al., *Endocrinology* 101:225-235, 1997.

Covalent complexes can also be made by isolating the desired component polypeptides and combining them *in vitro*. Covalent complexes
30 that can be prepared in this manner include homodimers of zveg3 polypeptides, heterodimers of two different zveg3 polypeptides, and

heterodimers of a zveg3 polypeptide and a polypeptide from another family member of the VEGF/PDGF family of proteins. The two polypeptides are mixed together under denaturing and reducing conditions, followed by renaturation of the proteins by removal of the denaturants. Removal can be
5 done by, for example, dialysis or size exclusion chromatography to provide for buffer exchange. When combining two different polypeptides, the resulting renaturated proteins may form homodimers of the individual components as well as heterodimers of the two polypeptide components. See, Cao et al., *J. Biol. Chem.* 271:3154-3162, 1996.

10 Depending upon the intended use, polypeptides and proteins of the present invention can be purified to $\geq 80\%$ purity, to $\geq 90\%$ purity, to $\geq 95\%$ purity, or to a pharmaceutically pure state, that is greater than 99.9% pure with respect to contaminating macromolecules, particularly other proteins and nucleic acids, and free of infectious and pyrogenic agents.

15 Expressed recombinant zveg3 proteins (including chimeric polypeptides and multimeric proteins) are purified by conventional protein purification methods, typically by a combination of chromatographic techniques. See, in general, Affinity Chromatography: Principles & Methods, Pharmacia LKB Biotechnology, Uppsala, Sweden, 1988; and Scopes, Protein
20 Purification: Principles and Practice, Springer-Verlag, New York, 1994. Proteins comprising a polyhistidine affinity tag (typically about 6 histidine residues) are purified by affinity chromatography on a nickel chelate resin. See, for example, Houchuli et al., *Bio/Technol.* 6: 1321-1325, 1988. Furthermore, the growth factor domain itself binds to nickel resin at pH 7.0-8.0
25 and 25 mM Na phosphate, 0.25 M NaCl. Bound protein can be eluted with a descending pH gradient down to pH 5.0 or an imidazole gradient. Proteins comprising a glu-glu tag can be purified by immunoaffinity chromatography according to conventional procedures. See, for example, Grussenmeyer et al., *ibid.* Maltose binding protein fusions are purified on an amylose column
30 according to methods known in the art. As disclosed in more detail below, zveg3 growth factor domain protein can be purified using a combination of

chromatography on a strong cation exchanger followed by hydrophobic interaction chromatography. When the protein is produced in BHK cells, insulin-like growth factor binding protein 4 (IGFBP4) co-purifies with the zveg3 under these conditions. Further purification can be obtained using reverse-phase HPLC, anion exchange on a quaternary amine strong cation exchanger at low ionic strength and pH from 7.0 to 9.0, or hydrophobic interaction chromatography on a phenyl ether resin. It has also been found that zveg3 binds to various dye matrices (e.g., BLUE1, BLUE 2, ORANGE 1, ORANGE 3, and RED3 from Lexton Scientific, Signal Hill, CA) in PBS at pH 6-8, from which the bound protein can be eluted in 1-2M NaCl in 20 mM boric acid buffer at pH 8.8. Protein eluted from RED3 may be passed over RED2 (Lexton Scientific) to remove remaining contaminants.

Using methods known in the art, zveg3 proteins can be prepared as monomers or multimers; glycosylated or non-glycosylated; pegylated or non-pegylated; and may or may not include an initial methionine amino acid residue.

The invention further provides polypeptides that comprise an epitope-bearing portion of a protein as shown in SEQ ID NO:2. An "epitope" is a region of a protein to which an antibody can bind. See, for example, Geysen et al., *Proc. Natl. Acad. Sci. USA* 81:3998-4002, 1984. Epitopes can be linear or conformational, the latter being composed of discontinuous regions of the protein that form an epitope upon folding of the protein. Linear epitopes are generally at least 6 amino acid residues in length. Relatively short synthetic peptides that mimic part of a protein sequence are routinely capable of eliciting an antiserum that reacts with the partially mimicked protein. See, Sutcliffe et al., *Science* 219:660-666, 1983. Antibodies that recognize short, linear epitopes are particularly useful in analytic and diagnostic applications that employ denatured protein, such as Western blotting (Tobin, *Proc. Natl. Acad. Sci. USA* 76:4350-4356, 1979). Antibodies to short peptides may also recognize proteins in native conformation and will thus be useful for monitoring

protein expression and protein isolation, and in detecting zveg3 proteins in solution, such as by ELISA or in immunoprecipitation studies.

Antigenic, epitope-bearing polypeptides of the present invention are useful for raising antibodies, including monoclonal antibodies, that specifically bind to a zveg3 protein. Antigenic, epitope-bearing polypeptides contain a sequence of at least six, often at least nine, more often from 15 to about 30 contiguous amino acid residues of a zveg3 protein (e.g., SEQ ID NO:2). Polypeptides comprising a larger portion of a zveg3 protein, i.e. from 30 to 50 residues up to the entire sequence are included. It is preferred that the amino acid sequence of the epitope-bearing polypeptide is selected to provide substantial solubility in aqueous solvents, that is the sequence includes relatively hydrophilic residues, and hydrophobic residues are substantially avoided. Such regions include residues 43-48, 96-101, 97-102, 260-265, and 330-335 of SEQ ID NO:2. As noted above, it is generally preferred to use somewhat longer peptides as immunogens, such as a peptide comprising residues 80-104, 299-314, and 299-326 of SEQ ID NO:2. The latter peptide can be prepared with an additional N-terminal cys residue to facilitate coupling.

As used herein, the term "antibodies" includes polyclonal antibodies, affinity-purified polyclonal antibodies, monoclonal antibodies, and antigen-binding fragments, such as F(ab')₂ and Fab proteolytic fragments. Genetically engineered intact antibodies or fragments, such as chimeric antibodies, Fv fragments, single chain antibodies and the like, as well as synthetic antigen-binding peptides and polypeptides, are also included. Non-human antibodies may be humanized by grafting non-human CDRs onto human framework and constant regions, or by incorporating the entire non-human variable domains (optionally "cloaking" them with a human-like surface by replacement of exposed residues, wherein the result is a "veneered" antibody). In some instances, humanized antibodies may retain non-human residues within the human variable region framework domains to enhance proper binding characteristics. Through humanizing antibodies, biological half-life may be increased, and the potential for adverse immune reactions upon

administration to humans is reduced. Monoclonal antibodies can also be produced in mice that have been genetically altered to produce antibodies that have a human structure.

Methods for preparing and isolating polyclonal and monoclonal
5 antibodies are well known in the art. See, for example, Cooligan, et al. (eds.),
Current Protocols in Immunology, National Institutes of Health, John Wiley and
Sons, Inc., 1995; Sambrook et al., Molecular Cloning: A Laboratory Manual,
second edition, Cold Spring Harbor, NY, 1989; and Hurrell, J. G. R. (ed.),
Monoclonal Hybridoma Antibodies: Techniques and Applications, CRC Press,
10 Inc., Boca Raton, FL, 1982. As would be evident to one of ordinary skill in the
art, polyclonal antibodies can be generated from inoculating a variety of warm-
blooded animals such as horses, cows, goats, sheep, dogs, chickens, rabbits,
mice, and rats with a zveg3 polypeptide or a fragment thereof. The
immunogenicity of a zveg3 polypeptide may be increased through the use of
15 an adjuvant, such as alum (aluminum hydroxide) or Freund's complete or
incomplete adjuvant. Polypeptides useful for immunization also include fusion
polypeptides, such as fusions of zveg3 or a portion thereof with an
immunoglobulin polypeptide or with maltose binding protein. If the polypeptide
portion is "haptten-like", such portion may be advantageously joined or linked to
20 a macromolecular carrier (such as keyhole limpet hemocyanin (KLH), bovine
serum albumin (BSA), or tetanus toxoid) for immunization.

Alternative techniques for generating or selecting antibodies
include *in vitro* exposure of lymphocytes to zveg3 protein or peptide, and
selection of antibody display libraries in phage or similar vectors (for instance,
25 through use of immobilized or labeled zveg3 protein or peptide). Techniques
for creating and screening such random peptide display libraries are known in
the art (Ladner et al., US Patent NO. 5,223,409; Ladner et al., US Patent NO.
4,946,778; Ladner et al., US Patent NO. 5,403,484 and Ladner et al., US
Patent NO. 5,571,698), and random peptide display libraries and kits for
30 screening such libraries are available commercially, for instance from Clontech
Laboratories (Palo Alto, CA), Invitrogen Inc. (San Diego, CA), New England

Biolabs, Inc. (Beverly, MA), and Pharmacia LKB Biotechnology Inc. (Piscataway, NJ). Random peptide display libraries can be screened using the zveg3 sequences disclosed herein to identify proteins which bind to zveg3. These "binding proteins", which interact with zveg3 polypeptides, can be used
5 for tagging cells or for isolating homologous polypeptides by affinity purification, or they can be directly or indirectly conjugated to drugs, toxins, radionuclides, and the like. These binding proteins can also be used in analytical methods, such as for screening expression libraries and neutralizing activity; within diagnostic assays, such as for determining circulating levels of polypeptides;
10 for detecting or quantitating soluble polypeptides as markers of underlying pathology or disease; and as zveg3 antagonists to block zveg3 binding and signal transduction *in vitro* and *in vivo*.

Antibodies are determined to be specifically binding if they bind to a zveg3 polypeptide, peptide or epitope with an affinity at least 10-fold greater
15 than the binding affinity to control (non-zveg3) polypeptide or protein. In this regard, a "non-zveg3 polypeptide" includes the related molecules VEGF, VEGF-B, VEGF-C, VEGF-D, PIGF, PDGF-A, and PDGF-B, but excludes zveg3 polypeptides from non-human species. Due to the high level of amino acid sequence identity expected between zveg3 orthologs, antibodies specific
20 for human zveg3 may also bind to zveg3 from other species. The binding affinity of an antibody can be readily determined by one of ordinary skill in the art, for example, by Scatchard analysis (Scatchard, G., *Ann. NY Acad. Sci.* 51: 660-672, 1949). Methods for screening and isolating specific antibodies are well known in the art. See, for example, Paul (ed.), Fundamental Immunology,
25 Raven Press, 1993; Getzoff et al., *Adv. in Immunol.* 43:1-98, 1988; Goding, J.W. (ed.), Monoclonal Antibodies: Principles and Practice, Academic Press Ltd., 1996; Benjamin et al., *Ann. Rev. Immunol.* 2:67-101, 1984.

A variety of assays known to those skilled in the art can be utilized to detect antibodies which specifically bind to zveg3 proteins or
30 peptides. Exemplary assays are described in detail in Antibodies: A Laboratory Manual, Harlow and Lane (Eds.), Cold Spring Harbor Laboratory Press, 1988.

Representative examples of such assays include: concurrent immunoelectrophoresis, radioimmunoassay, radioimmuno-precipitation, enzyme-linked immunosorbent assay (ELISA), dot blot or Western blot assay, inhibition or competition assay, and sandwich assay. In addition, antibodies
5 can be screened for binding to wild-type versus mutant zveg3 protein or polypeptide.

Antibodies to zveg3 may be used for tagging cells that express zveg3; for isolating zveg3 by affinity purification; for diagnostic assays for determining circulating levels of zveg3 polypeptides; for detecting or
10 quantitating soluble zveg3 as a marker of underlying pathology or disease; in analytical methods employing FACS; for screening expression libraries; for generating anti-idiotypic antibodies; and as neutralizing antibodies or as antagonists to block zveg3 activity *in vitro* and *in vivo*. Suitable direct tags or labels include radionuclides, enzymes, substrates, cofactors, inhibitors,
15 fluorescent markers, chemiluminescent markers, magnetic particles and the like; indirect tags or labels may feature use of biotin-avidin or other complement/anti-complement pairs as intermediates. Antibodies may also be directly or indirectly conjugated to drugs, toxins, radionuclides and the like, and these conjugates used for *in vivo* diagnostic or therapeutic applications.
20 Moreover, antibodies to zveg3 or fragments thereof may be used *in vitro* to detect denatured zveg3 or fragments thereof in assays, for example, Western Blots or other assays known in the art. Antibodies can also be used to target an attached therapeutic or diagnostic moiety to cells expressing zveg3 or receptors for zveg3.

25 For some applications (e.g., certain therapeutic applications) it is preferred to use neutralizing antibodies. As used herein, the term "neutralizing antibody" denotes an antibody that inhibits at least 50% of the biological activity of the cognate antigen when the antibody is added at a 1000-fold molar excess. Those of skill in the art will recognize that greater neutralizing activity
30 is sometimes desirable, and antibodies that provide 50% inhibition at a 100-fold or 10-fold molar excess may be advantageously employed.

Activity of zveg3 proteins and antagonists thereof can be measured *in vitro* using cultured cells or *in vivo* by administering molecules of the claimed invention to an appropriate animal model. Target cells for use in zveg3 activity assays include vascular cells (especially endothelial cells and smooth muscle cells), hematopoietic (myeloid and lymphoid) cells, liver cells (including hepatocytes, fenestrated endothelial cells, Kupffer cells, and Ito cells), fibroblasts (including human dermal fibroblasts and lung fibroblasts), neurite cells (including astrocytes, glial cells, dendritic cells, and PC-12 cells), Schwann cells, fetal lung cells, articular synoviocytes, pericytes, chondrocytes, oligodendrocytes, osteoblasts, and other cells expressing PDGF alpha receptors.

Zveg3 proteins can be analyzed for receptor binding activity by a variety of methods well known in the art, including receptor competition assays (Bowen-Pope and Ross, *Methods Enzymol.* 109:69-100, 1985), use of soluble receptors, and use of receptors produced as IgG fusion proteins (U.S. Patent No. 5,750,375). Receptor binding assays can be performed on cell lines that contain known cell-surface receptors for evaluation. The receptors can be naturally present in the cell, or can be recombinant receptors expressed by genetically engineered cells. Cell types that are able to bind zveg3 can be identified through the use of zveg3-toxin conjugates, such as conjugates of a zveg3 protein and saporin. Binding of the zveg3-toxin conjugate by cells, either in tissue culture, in organ cultures, or *in vivo* settings will allow for the incorporation of the conjugate into the cell. Once inside the cell saporin has a toxic effect on the cell, thereby killing it. This activity can be used to identify cell types that are able to bind and internalize zveg3. In addition to allowing for the identification of responsive cell types, toxin conjugates can be used in *in vivo* studies to identify organs and tissues where zveg3 has a biological activity by looking for pathology within the animal following injection of the conjugate.

Activity of zveg3 proteins can be measured *in vitro* using cultured cells. Mitogenic activity can be measured using known assays, including ^3H -

thymidine incorporation assays (as disclosed by, e.g., Raines and Ross, *Methods Enzymol.* 109:749-773, 1985 and Wahl et al., *Mol. Cell Biol.* 8:5016-5025, 1988), dye incorporation assays (as disclosed by, for example, Mosman, *J. Immunol. Meth.* 65:55-63, 1983 and Raz et al., *Acta Trop.* 68:139-147, 1997) or cell counts. Suitable mitogenesis assays measure incorporation of ³H-thymidine into (1) 20% confluent cultures to look for the ability of zveg3 proteins to further stimulate proliferating cells, and (2) quiescent cells held at confluence for 48 hours to look for the ability of zveg3 proteins to overcome contact-induced growth inhibition. Suitable dye incorporation assays include measurement of the incorporation of the dye Alamar blue (Raz et al., *ibid.*) into target cells. See also, Gospodarowicz et al., *J. Cell. Biol.* 70:395-405, 1976; Ewton and Florini, *Endocrinol.* 106:577-583, 1980; and Gospodarowicz et al., *Proc. Natl. Acad. Sci. USA* 86:7311-7315, 1989. Cell differentiation can be assayed using suitable precursor cells that can be induced to differentiate into a more mature phenotype. For example, endothelial cells and hematopoietic cells are derived from a common ancestral cell, the hemangioblast (Choi et al., *Development* 125:725-732, 1998). Mesenchymal stem cells can also be used to measure the ability of zveg3 protein to stimulate differentiation into osteoblasts. Differentiation is indicated by the expression of osteocalcin, the ability of the cells to mineralize, and the expression of alkaline phosphatase, all of which can be measured by routine methods known in the art. Effects of zveg3 proteins on tumor cell growth and metastasis can be analyzed using the Lewis lung carcinoma model, for example as described by Cao et al., *J. Exp. Med.* 182:2069-2077, 1995. Activity of zveg3 proteins on cells of neural origin can be analyzed using assays that measure effects on neurite growth.

Zveg3 activity may also be detected using assays designed to measure zveg3-induced production of one or more additional growth factors or other macromolecules. Such assays include those for determining the presence of hepatocyte growth factor (HGF), epidermal growth factor (EGF), transforming growth factor alpha (TGF α), interleukin-6 (IL-6), VEGF, acidic fibroblast growth factor (aFGF), and angiogenin. Suitable assays include

mitogenesis assays using target cells responsive to the macromolecule of interest, receptor-binding assays, competition binding assays, immunological assays (e.g., ELISA), and other formats known in the art. Metalloprotease secretion is measured from treated primary human dermal fibroblasts, synoviocytes and chondrocytes. The relative levels of collagenase, gelatinase and stromalysin produced in response to culturing in the presence of a zveg3 protein is measured using zymogram gels (Loita and Stetler-Stevenson, *Cancer Biology* 1:96-106, 1990). Procollagen/collagen synthesis by dermal fibroblasts and chondrocytes in response to a test protein is measured using ³H-proline incorporation into nascent secreted collagen. ³H-labeled collagen is visualized by SDS-PAGE followed by autoradiography (Unemori and Amento, *J. Biol. Chem.* 265: 10681-10685, 1990). Glycosaminoglycan (GAG) secretion from dermal fibroblasts and chondrocytes is measured using a 1,9-dimethylmethylene blue dye binding assay (Farndale et al., *Biochim. Biophys. Acta* 883:173-177, 1986). Collagen and GAG assays are also carried out in the presence of IL-1 β or TGF- β to examine the ability of zveg3 protein to modify the established responses to these cytokines.

Monocyte activation assays are carried out (1) to look for the ability of zveg3 proteins to further stimulate monocyte activation, and (2) to examine the ability of zveg3 proteins to modulate attachment-induced or endotoxin-induced monocyte activation (Fuhlbrigge et al., *J. Immunol.* 138: 3799-3802, 1987). IL-1 β and TNF α levels produced in response to activation are measured by ELISA (Biosource, Inc. Camarillo, CA). Monocyte/macrophage cells, by virtue of CD14 (LPS receptor), are exquisitely sensitive to endotoxin, and proteins with moderate levels of endotoxin-like activity will activate these cells.

Hematopoietic activity of zveg3 proteins can be assayed on various hematopoietic cells in culture. Suitable assays include primary bone marrow or peripheral blood leukocyte colony assays, and later stage lineage-restricted colony assays, which are known in the art (e.g., Holly et al., WIPO Publication WO 95/21920). Marrow cells plated on a suitable semi-solid

medium (e.g., 50% methylcellulose containing 15% fetal bovine serum, 10% bovine serum albumin, and 0.6% PSN antibiotic mix) are incubated in the presence of test polypeptide, then examined microscopically for colony formation. Known hematopoietic factors are used as controls. Mitogenic activity of zveg3 polypeptides on hematopoietic cell lines can be measured using ^3H -thymidine incorporation assays, dye incorporation assays, or cell counts (Raines and Röss, *Methods Enzymol.* 109:749-773, 1985 and Foster et al., U.S. Patent No. 5,641,655). For example, cells are cultured in multi-well microtiter plates. Test samples and ^3H -thymidine are added, and the cells are incubated overnight at 37°C. Contents of the wells are transferred to filters, dried, and counted to determine incorporation of label. Cell proliferation can also be measured using a colorimetric assay based on the metabolic breakdown of 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyl tetrazolium bromide (MTT) (Mosman, *ibid.*). Briefly, a solution of MTT is added to 100 μl of assay cells, and the cells are incubated at 37° C. After 4 hours, 200 μl of 0.04 N HCl in isopropanol is added, the solution is mixed, and the absorbance of the sample is measured at 570 nm.

Cell migration is assayed essentially as disclosed by Kähler et al. (*Arteriosclerosis, Thrombosis, and Vascular Biology* 17:932-939, 1997). A protein is considered to be chemotactic if it induces migration of cells from an area of low protein concentration to an area of high protein concentration. The assay is performed using modified Boyden chambers with a polystyrene membrane separating the two chambers (Transwell; Corning Costar Corp.). The test sample, diluted in medium containing 1% BSA, is added to the lower chamber of a 24-well plate containing Transwells. Cells are then placed on the Transwell insert that has been pretreated with 0.2% gelatin. Cell migration is measured after 4 hours of incubation at 37°C. Non-migrating cells are wiped off the top of the Transwell membrane, and cells attached to the lower face of the membrane are fixed and stained with 0.1% crystal violet. Stained cells are then extracted with 10% acetic acid and absorbance is measured at 600 nm. Migration is then calculated from a standard calibration curve.

Smooth muscle cell (SMC) migration can be measured in the aortic explant assay of Kenagy et al. (*Circulation* 96:3555-3560, 1997). In a typical protocol, explants are prepared from baboon thoracic aortas, and the inner media is isolated and chopped into 1-mm² pieces. The explants are placed in tissue culture flasks containing DMEM supplemented with 5 µg/ml transferrin, 6 µg/ml insulin, 1 mg/ml ovalbumin, and the test compound. The number of migrating cells is determined daily.

Cell adhesion activity is assayed essentially as disclosed by LaFleur et al. (*J. Biol. Chem.* 272:32798-32803, 1997). Briefly, microtiter plates are coated with the test protein, non-specific sites are blocked with BSA, and cells (such as smooth muscle cells, leukocytes, or endothelial cells) are plated at a density of approximately 10⁴ - 10⁵ cells/well. The wells are incubated at 37°C (typically for about 60 minutes), then non-adherent cells are removed by gentle washing. Adhered cells are quantitated by conventional methods (e.g., by staining with crystal violet, lysing the cells, and determining the optical density of the lysate). Control wells are coated with a known adhesive protein, such as fibronectin or vitronectin.

Assays for angiogenic activity are also known in the art. For example, the effect of zveg3 proteins on primordial endothelial cells in angiogenesis can be assayed in the chick chorioallantoic membrane angiogenesis assay (Leung, *Science* 246:1306-1309, 1989; Ferrara, *Ann. NY Acad. Sci.* 752:246-256, 1995). Briefly, a small window is cut into the shell of an eight-day old fertilized egg, and a test substance is applied to the chorioallantoic membrane. After 72 hours, the membrane is examined for neovascularization. Other suitable assays include microinjection of early stage quail (*Coturnix coturnix japonica*) embryos as disclosed by Drake et al. (*Proc. Natl. Acad. Sci. USA* 92:7657-7661, 1995); the rodent model of corneal neovascularization disclosed by Muthukkaruppan and Auerbach (*Science* 205:1416-1418, 1979), wherein a test substance is inserted into a pocket in the cornea of an inbred mouse; and the hamster cheek pouch assay (Höckel et al., *Arch. Surg.* 128:423-429, 1993). Induction of vascular permeability, which

is indicative of angiogenic activity, is measured in assays designed to detect leakage of protein from the vasculature of a test animal (e.g., mouse or guinea pig) after administration of a test compound (Miles and Miles, *J. Physiol.* 118:228-257, 1952; Feng et al., *J. Exp. Med.* 183:1981-1986, 1996). *In vitro* assays for angiogenic activity include the tridimensional collagen gel matrix model (Pepper et al. *Biochem. Biophys. Res. Comm.* 189:824-831, 1992 and Ferrara et al., *Ann. NY Acad. Sci.* 732:246-256, 1995), which measures the formation of tube-like structures by microvascular endothelial cells; and matrigel models (Grant et al., "Angiogenesis as a component of epithelial-mesenchymal interactions" in Goldberg and Rosen, *Epithelial-Mesenchymal Interaction in Cancer*, Birkhäuser Verlag, 1995, 235-248; Baatout, *Anticancer Research* 17:451-456, 1997), which are used to determine effects on cell migration and tube formation by endothelial cells seeded in matrigel, a basement membrane extract enriched in laminin. Angiogenesis assays can be carried out in the presence and absence of VEGF to assess possible combinatorial effects. VEGF can be used as a control within *in vivo* assays.

Zvegf3 activity can also be measured using assays that measure axon guidance and growth. Of particular interest are assays that indicate changes in neuron growth patterns, for example those disclosed in Hastings, WIPO Publication WO 97/29189 and Walter et al., *Development* 101:685-96, 1987. Assays to measure the effects on neuron growth are well known in the art. For example, the C assay (e.g., Raper and Kapfhammer, *Neuron* 4:21-9, 1990 and Luo et al., *Cell* 75:217-27, 1993) can be used to determine collapsing activity of zvegf3 on growing neurons. Other methods that can assess zvegf3-induced effects on neurite extension are also known. See, Goodman, *Annu. Rev. Neurosci.* 19:341-77, 1996. Conditioned media from cells expressing a zvegf3 protein, a zvegf3 agonist, or a zvegf3 antagonist, or aggregates of such cells, can be placed in a gel matrix near suitable neural cells, such as dorsal root ganglia (DRG) or sympathetic ganglia explants, which have been co-cultured with nerve growth factor. Compared to control cells, zvegf3-induced changes in neuron growth can be measured (as disclosed by, for example,

Messersmith et al., *Neuron* 14:949-59, 1995 and Puschel et al., *Neuron* 14:941-8, 1995). Likewise neurite outgrowth can be measured using neuronal cell suspensions grown in the presence of molecules of the present invention. See, for example, O'Shea et al., *Neuron* 7:231-7, 1991 and DeFreitas et al.,
5 *Neuron* 15:333-43, 1995.

The biological activities of zveg3 proteins can be studied in non-human animals by administration of exogenous protein, by expression of zveg3-encoding polynucleotides, and by suppression of endogenous zveg3 expression through antisense or knock-out techniques. Zveg3 proteins can be
10 administered or expressed individually, in combination with other zveg3 proteins, or in combination with non-veg3 proteins, including other growth factors (e.g., other VEGFs, PIGFs, or PDGFs). For example, a combination of zveg3 polypeptides (e.g., a combination of zveg3₁₅₋₁₆₃, zveg3₁₅₋₂₃₀, and zveg3₂₃₅₋₃₄₅) can be administered to a test animal or expressed in the animal.
15 Test animals are monitored for changes in such parameters as clinical signs, body weight, blood cell counts, clinical chemistry, histopathology, and the like.

Effects of zveg3 and zveg3 antagonists on liver and kidney fibrosis can be tested in known animal models, such as the db/db mouse model disclosed by Cohen et al., *Diabetologia* 39:270-274, 1996 and Cohen et al., *J. Clin. Invest.* 95:2338-2345, 1995 or transgenic animal models (Imai et al., *Contrib. Nephrol.* 107:205-215, 1994).
20

Effects on fibrosis can also be assayed in a mouse model using bleomycin. The chemotherapy agent bleomycin is a known causative agent of pulmonary fibrosis in humans and can induce interstitial lung disease in mice, including an increase in the number of fibroblasts, enhanced collagen
25 deposition, and dysregulated matrix remodeling. C57Bl/6 mice are administered bleomycin by osmotic minipump for 1 week. There follows a period of inflammation, with cutaneous toxicity beginning approximately 4-7 days after bleomycin administration and continuing for about a week, after
30 which the mice appear to regain health. About 3-4 weeks after the finish of bleomycin delivery, the mice are sacrificed, and the lungs are examined

histologically for signs of fibrosis. Scoring is based on the extent of lung fibrotic lesions and their severity. Serum is assayed for lactic dehydrogenase, an intracellular enzyme that is released into the circulation upon general cell death or injury. Lung tissue is assayed for hydroxyproline as a measure of collagen
5 deposition.

Stimulation of coronary collateral growth can be measured in known animal models, including a rabbit model of peripheral limb ischemia and hind limb ischemia and a pig model of chronic myocardial ischemia (Ferrara et al., *Endocrine Reviews* 18:4-25, 1997). Zveg3 proteins are assayed in the
10 presence and absence of VEGFs, angiopoietins, and basic FGF to test for combinatorial effects. These models can be modified by the use of adenovirus or naked DNA for gene delivery as disclosed in more detail below, resulting in local expression of the test protein(s).

Efficacy of zveg3 polypeptides in promoting wound healing can
15 be assayed in animal models. One such model is the linear skin incision model of Mustoe et al. (*Science* 237:1333, 1987). In a typical procedure, a 6-cm incision is made in the dorsal pelt of an adult rat, then closed with wound clips. Test substances and controls (in solution, gel, or powder form) are applied before primary closure. Administration will often be limited to a single
20 application, although additional applications can be made on succeeding days by careful injection at several sites under the incision. Wound breaking strength is evaluated between 3 and 21 days post wounding. In a second model, multiple, small, full-thickness excisions are made on the ear of a rabbit. The cartilage in the ear splints the wound, removing the variable of wound
25 contraction from the evaluation of closure. Experimental treatments and controls are applied. The geometry and anatomy of the wound site allow for reliable quantification of cell ingrowth and epithelial migration, as well as quantitative analysis of the biochemistry of the wounds (e.g., collagen content). See, Mustoe et al., *J. Clin. Invest.* 87:694, 1991. The rabbit ear model can be
30 modified to create an ischemic wound environment, which more closely resembles the clinical situation (Ahn et al., *Ann. Plast. Surg.* 24:17, 1990).

Within a third model, healing of partial-thickness skin wounds in pigs or guinea pigs is evaluated (LeGrand et al., *Growth Factors* 8:307, 1993). Experimental treatments are applied daily on or under dressings. Seven days after wounding, granulation tissue thickness is determined. This model is particularly useful for dose-response studies, as it is more quantitative than other *in vivo* models of wound healing. A full thickness excision model can also be employed. Within this model, the epidermis and dermis are removed down to the panniculus carnosus in rodents or the subcutaneous fat in pigs. Experimental treatments are applied topically on or under a dressing, and can be applied daily if desired. The wound closes by a combination of contraction and cell ingrowth and proliferation. Measurable endpoints include time to wound closure, histologic score, and biochemical parameters of wound tissue. Impaired wound healing models are also known in the art (e.g., Cromack et al., *Surgery* 113:36, 1993; Pierce et al., *Proc. Natl. Acad. Sci. USA* 86:2229, 1989; Greenhalgh et al., *Amer. J. Pathol.* 136:1235, 1990). Delay or prolongation of the wound healing process can be induced pharmacologically by treatment with steroids, irradiation of the wound site, or by concomitant disease states (e.g., diabetes). Linear incisions or full-thickness excisions are most commonly used as the experimental wound. Endpoints are as disclosed above for each type of wound. Subcutaneous implants can be used to assess compounds acting in the early stages of wound healing (Broadley et al., *Lab. Invest.* 61:571, 1985; Sprugel et al., *Amer. J. Pathol.* 129: 601, 1987). Implants are prepared in a porous, relatively non-inflammatory container (e.g., polyethylene sponges or expanded polytetrafluoroethylene implants filled with bovine collagen) and placed subcutaneously in mice or rats. The interior of the implant is empty of cells, producing a "wound space" that is well-defined and separable from the preexisting tissue. This arrangement allows the assessment of cell influx and cell type as well as the measurement of vasculogenesis/angiogenesis and extracellular matrix production.

Expression of zveg3 proteins in animals provides models for study of the biological effects of overproduction or inhibition of protein activity in

vivo. Zveg3f3-encoding polynucleotides can be introduced into test animals, such as mice, using viral vectors or naked DNA, or transgenic animals can be produced. In general, a zveg3f3 protein is expressed with a secretory peptide. Suitable secretory peptides include the zveg3f3 secretory peptide (e.g., residues
5 1-14 of SEQ ID NO:2) and heterologous secretory peptides. An exemplary heterologous secretory peptide is that of human tissue plasminogen activator (t-PA). The t-PA secretory peptide may be modified to reduce undesired proteolytic cleavage as disclosed in U.S. Patent No. 5,641,655.

Proteins of the present invention can be assayed *in vivo* using
10 viral delivery systems. Exemplary viruses include adenovirus, herpesvirus, retroviruses, vaccinia virus, and adeno-associated virus (AAV). For review, see Becker et al., *Meth. Cell Biol.* 43:161-89, 1994; and Douglas and Curiel, *Science & Medicine* 4:44-53, 1997. Adenovirus (reviewed by Becker et al., *Meth. Cell Biol.* 43:161-89, 1994; and Douglas and Curiel, *Science & Medicine*
15 4:44-53, 1997) offers several advantages. Adenovirus can (i) accommodate relatively large DNA inserts; (ii) be grown to high-titer; (iii) infect a broad range of mammalian cell types; and (iv) be used with many different promoters including ubiquitous, tissue specific, and regulatable promoters. Because adenoviruses are stable in the bloodstream, they can be administered by
20 intravenous injection. If the adenoviral delivery system has an E1 gene deletion, the virus cannot replicate in the host cells. However, the host's tissue (e.g., liver) will express and process (and, if a secretory signal sequence is present, secrete) the heterologous protein. Secreted proteins will enter the circulation in the highly vascularized liver, and effects on the infected animal
25 can be determined. Adenoviral vectors containing various deletions of viral genes can be used in an attempt to reduce or eliminate immune responses to the vector. Such adenoviruses are E1 deleted, and in addition contain deletions of E2A or E4 (Lusky et al., *J. Virol.* 72:2022-2032, 1998; Raper et al., *Human Gene Therapy* 9:671-679, 1998). In addition, deletion of E2b is
30 reported to reduce immune responses (Amalfitano, et al., *J. Virol.* 72:926-933, 1998). Generation of so-called "gutless" adenoviruses where all viral

transcription units are deleted is particularly advantageous for insertion of large inserts of heterologous DNA. For review, see Yeh and Perricaudet, *FASEB J.* 11:615-623, 1997. Retroviral vectors are disclosed, for example, by Anderson et al., U.S. Patent No. 5,399,346; Mann et al., *Cell* 33:153, 1983; Temin et al.,
5 U.S. Patent No. 4,650,764; Temin et al., U.S. Patent No. 4,980,289; Markowitz et al., *J. Virol.* 62:1120, 1988; Temin et al., U.S. Patent No. 5,124,263; Dougherty et al., WIPO publication WO 95/07358; and Kuo et al., *Blood* 82:845, 1993.

In an alternative method, a vector can be introduced by liposome-mediated transfection, a technique that provides certain practical advantages,
10 including the molecular targeting of liposomes to specific cells. Directing transfection to particular cell types is particularly advantageous in a tissue with cellular heterogeneity, such as the pancreas, liver, kidney, and brain. Lipids may be chemically coupled to other molecules for the purpose of targeting.
15 Targeted peptides (e.g., hormones or neurotransmitters), proteins such as antibodies, or non-peptide molecules can be coupled to liposomes chemically.

Within another embodiment target cells are removed from the animal, and DNA is introduced as a naked DNA plasmid. The transformed cells are then re-implanted into the body of the animal. Naked DNA vectors
20 can be introduced into the desired host cells by methods known in the art, e.g., transfection, electroporation, microinjection, transduction, cell fusion, DEAE dextran, calcium phosphate precipitation, use of a gene gun or use of a DNA vector transporter. See, e.g., Wu et al., *J. Biol. Chem.* 267:963-7, 1992; Wu et al., *J. Biol. Chem.* 263:14621-4, 1988.

25 Mice engineered to express the *zveg3* gene, referred to as "transgenic mice," and mice that exhibit a complete absence of *zveg3* gene function, referred to as "knockout mice," can also be generated (Snouwaert et al., *Science* 257:1083, 1992; Lowell et al., *Nature* 366:740-42, 1993; Capecchi, *Science* 244:1288-1292, 1989; Palmiter et al., *Ann. Rev. Genet.* 20:465-499,
30 1986). Transgenesis experiments can be performed using normal mice or mice with genetic disease or other altered phenotypes. Transgenic mice that

over-express zveg3, either ubiquitously or under a tissue-specific or tissue-restricted promoter, can be used to determine whether or not over-expression causes a phenotypic change. Suitable promoters include metallothionein, albumin (Pinkert et al., *Genes Dev.* 1(3):268-76, 1987), and K-14 keratinocyte (Vassar et al., *Proc. Natl. Acad. Sci. USA* 86(5):1563-1567, 1989) gene promoters. The metallothionein-1 (MT-1) promoter provides expression in liver and other tissues, often leading to high levels of circulating protein. Over-expression of a wild-type zveg3 polypeptide, polypeptide fragment or a mutant thereof may alter normal cellular processes, resulting in a phenotype that identifies a tissue in which zveg3 expression is functionally relevant and may indicate a therapeutic target for the zveg3, its agonists or antagonists. For example, a transgenic mouse can be engineered to over-express a full-length zveg3 sequence, which may result in a phenotype that shows similarity with human diseases. Similarly, knockout zveg3 mice can be used to determine where zveg3 is absolutely required *in vivo*. The phenotype of knockout mice is predictive of the *in vivo* effects of zveg3 antagonists. Knockout mice can also be used to study the effects of zveg3 proteins in models of disease, including, for example, cancer, atherosclerosis, rheumatoid arthritis, ischemia, and cardiovascular disease. The human zveg3 cDNA can be used to isolate murine zveg3 mRNA, cDNA and genomic DNA as disclosed above, which are subsequently used to generate knockout mice. These mice may be employed to study the zveg3 gene and the protein encoded thereby in an *in vivo* system, and can be used as *in vivo* models for corresponding human diseases. Moreover, transgenic mice expressing zveg3 antisense polynucleotides or ribozymes directed against zveg3, described herein, can be used analogously to knockout mice described above.

The functional relationship between zveg3 and PDGF ligands can be investigated through knockin experiments. For example, the PDGF A or PDGF B gene in an animal can be replaced by knocking a zveg3 gene or cDNA into the genomic locus of a PDGF ligand in embryonic stem (ES) cells, which are then used to generate the knockin mice, in which zveg3 protein is

expressed by the (replaced) PDGF genomic locus. Such knockin mice can be used to address a number of questions, such as whether zveg3 can substitute PDGF function during development; whether zveg3 has unique functions; and, using cell lines established from the mice, the mechanism of zveg3 signal transduction. See, for example, Wang et al., *Development* 124:2507-2513, 1997; Zhuang et al., *Mol. Cell Biol.* 18:3340-3349, 1998; Geng et al., *Cell* 97:767-777, 1999. In a further application of this technology, individual domains of zveg3 can be specifically deleted or modified by knocking in a modified zveg3 sequence or a specific spliced version of zveg3, thereby providing a transgenic model for the study of functional domains of the protein. See, for example, Zhuang et al. (*ibid.*) and Baudoin et al. (*Genes Dev.* 12:1202-1216, 1998). In another application, tissues and cell types that express the zveg3 gene can be identified by knockin of a sensitive reporter gene (e.g., *LacZ*) into the zveg3 locus. See, for example, Monroe et al., *Immunity* 11:201-212, 1999; Zhuang et al., *ibid.*; Geng et al., *ibid.*

Antisense methodology can be used to inhibit zveg3 gene transcription to examine the effects of such inhibition *in vivo*. Polynucleotides that are complementary to a segment of a zveg3-encoding polynucleotide (e.g., a polynucleotide as set forth in SEQ ID NO:1) are designed to bind to zveg3-encoding mRNA and to inhibit translation of such mRNA. Such antisense oligonucleotides can also be used to inhibit expression of zveg3 polypeptide-encoding genes in cell culture.

Those skilled in the art will recognize that the assays disclosed herein can be readily adapted to study the activity of zveg3 proteins, anti-zveg3 antibodies and other antagonists, and test substances derived from a variety of sources.

Zveg3 proteins may be used therapeutically to stimulate tissue development or repair, or cellular differentiation or proliferation. Zveg3 has been found to bind to PDGF alpha receptor and to stimulate alpha receptor-mediated cellular processes. The protein can therefore be used as a PDGF alpha receptor agonist. Specific applications include, without limitation: the

treatment of full-thickness skin wounds, including venous stasis ulcers and other chronic, non-healing wounds, particularly in cases of compromised wound healing due to diabetes mellitus, connective tissue disease, smoking, burns, and other exacerbating conditions; fracture repair; skin grafting; within
5 reconstructive surgery to promote neovascularization and increase skin flap survival; to establish vascular networks in transplanted cells and tissues, such as transplanted islets of Langerhans; to treat female reproductive tract disorders, including acute or chronic placental insufficiency (an important factor causing perinatal morbidity and mortality) and prolonged bleeding; to promote
10 the growth of tissue damaged by periodontal disease; to promote the repair of damaged liver tissue; in the treatment of acute and chronic lesions of the gastrointestinal tract, including duodenal ulcers, which are characterized by a deficiency of microvessels; to promote angiogenesis and prevent neuronal degeneration due to chronic cerebral ischemia; to accelerate the formation of
15 collateral blood vessels in ischemic limbs; to promote vessel repair and development of collateral circulation following myocardial infarction so as to limit ischemic injury; and to stimulate hematopoiesis. The polypeptides are also useful additives in tissue adhesives for promoting revascularization of the healing tissue.

20 Of particular interest is the use of zveg3 or zveg3 antagonists for the treatment or repair of liver damage, including damage due to chronic liver disease, including chronic active hepatitis (including hepatitis C) and many other types of cirrhosis. Widespread, massive necrosis, including destruction of virtually the entire liver, can be caused by, *inter alia*, fulminant viral hepatitis;
25 overdoses of the analgesic acetaminophen; exposure to other drugs and chemicals such as halothane, monoamine oxidase inhibitors, agents employed in the treatment of tuberculosis, phosphorus, carbon tetrachloride, and other industrial chemicals. Conditions associated with ultrastructural lesions that do not necessarily produce obvious liver cell necrosis include Reye's syndrome in
30 children, tetracycline toxicity, and acute fatty liver of pregnancy. Cirrhosis, a diffuse process characterized by fibrosis and a conversion of normal

architecture into structurally abnormal nodules, can come about for a variety reasons including alcohol abuse, post necrotic cirrhosis (usually due to chronic active hepatitis), biliary cirrhosis, pigment cirrhosis, cryptogenic cirrhosis, Wilson's disease, and alpha-1-antitrypsin deficiency. Zveg3 may also be
5 useful for the treatment of hepatic chronic passive congestion (CPC) and central hemorrhagic necrosis (CHN), which are two circulatory changes representing a continuum encountered in right-sided heart failure. Other circulatory disorders that may be treated with zveg3 include hepatic vein thrombosis, portal vein thrombosis, and cardiac sclerosis. In cases of liver
10 fibrosis it may be beneficial to administer a zveg3 antagonist to suppress the activation of stellate cells, which have been implicated in the production of extracellular matrix in fibrotic liver (Li and Friedman, *J. Gastroenterol. Hepatol.* 14:618-633, 1999).

Zveg3 polypeptides can be administered alone or in combination
15 with other vasculogenic or angiogenic agents, including VEGF. For example, basic and acidic FGFs and VEGF have been found to play a role in the development of collateral circulation, and the combined use of zveg3 with one or more of these factors may be advantageous. VEGF has also been implicated in the survival of transplanted islet cells (Gorden et al.
20 *Transplantation* 63:436-443, 1997; Pepper, *Arteriosclerosis, Throm. and Vascular Biol.* 17:605-619, 1997). Basic FGF has been shown to induce angiogenesis and accelerate healing of ulcers in experimental animals (reviewed by Folkman, *Nature Medicine* 1:27-31, 1995). VEGF has been shown to promote vessel re-endothelialization and to reduce intimal
25 hyperplasia in animal models of restenosis (Asahara et al., *Circulation* 91:2802-2809, 1995; Callow et al., *Growth Factors* 10:223-228, 1994); efficacy of zveg3 polypeptides can be tested in these and other known models. When using zveg3 in combination with an additional agent, the two compounds can be administered simultaneously or sequentially as appropriate for the specific
30 condition being treated.

Zveg3 proteins may be used either alone or in combination with other hematopoietic factors such as IL-3, G-CSF, GM-CSF, or stem cell factor to enhance expansion and mobilization of endothelial precursor stem cells. Cells that can be expanded in this manner include cells isolated from bone marrow or cells isolated from blood. Zveg3 proteins may also be given directly to an individual to enhance endothelial stem cell production and differentiation within the treated individual. The stem cells, either developed within the patient, or provided back to a patient, may then play a role in modulating areas of ischemia within the body, thereby providing a therapeutic effect. These cells may also be useful in enhancing re-endothelialization of areas devoid of endothelial coverage, such as vascular grafts, vascular stents, and areas where the endothelial coverage has been damaged or removed (e.g., areas of angioplasty). Zveg3 proteins may also be used in combination with other growth and differentiation factors such as angiopoietin-1 (Davis et al., *Cell* 87:1161-1169, 1996) to help create and stabilize new vessel formation in areas requiring neovascularization, including areas of ischemia (cardiac or peripheral ischemia), organ transplants, wound healing, and tissue grafting.

Zveg3 proteins, agonists and antagonists may be used to modulate neurite growth and development and demarcate nervous system structures. As such, Zveg3 proteins, agonists, or antagonists may be useful in the treatment of peripheral neuropathies by increasing spinal cord and sensory neurite outgrowth, and as part of a therapeutic treatment for the regeneration of neurite outgrowths following strokes, brain damage caused by head injuries, and paralysis caused by spinal injuries. Application may also be made in treating neurodegenerative diseases such as multiple sclerosis, Alzheimer's disease and Parkinson's disease. Application may also be made in mediating development and innervation pattern of stomach tissue.

For pharmaceutical use, zveg3 proteins are formulated for topical or parenteral, particularly intravenous or subcutaneous, delivery according to conventional methods. In general, pharmaceutical formulations will include a zveg3 polypeptide in combination with a pharmaceutically

acceptable vehicle, such as saline, buffered saline, 5% dextrose in water, or the like. Formulations may further include one or more excipients, preservatives, solubilizers, buffering agents, albumin to prevent protein loss on vial surfaces, etc. Methods of formulation are well known in the art and are disclosed, for example, in *Remington: The Science and Practice of Pharmacy*, 5 Gennaro, ed., Mack Publishing Co., Easton, PA, 19th ed., 1995. Zveg3 will ordinarily be used in a concentration of about 10 to 100 µg/ml of total volume, although concentrations in the range of 1 ng/ml to 1000 µg/ml may be used. For topical application, such as for the promotion of wound healing, the protein 10 will be applied in the range of 0.1-10 µg/cm² of wound area, with the exact dose determined by the clinician according to accepted standards, taking into account the nature and severity of the condition to be treated, patient traits, etc. Determination of dose is within the level of ordinary skill in the art. The therapeutic formulations will generally be administered over the period required 15 for neovascularization, typically from one to several months and, in treatment of chronic conditions, for a year or more. Dosing is daily or intermittently over the period of treatment. Intravenous administration will be by bolus injection or infusion over a typical period of one to several hours. Sustained release formulations can also be employed. In general, a therapeutically effective 20 amount of zveg3 is an amount sufficient to produce a clinically significant change in the treated condition, such as a clinically significant reduction in time required by wound closure, a significant reduction in wound area, a significant improvement in vascularization, a significant reduction in morbidity, or a significantly increased histological score.

25 Proteins of the present invention are useful for modulating the proliferation, differentiation, migration, or metabolism of responsive cell types, which include both primary cells and cultured cell lines. Of particular interest in this regard are liver cells, hematopoietic cells (including stem cells and mature myeloid and lymphoid cells), endothelial cells, neuronal cells, and 30 mesenchymal cells (including fibroblasts and smooth muscle cells). Zveg3 polypeptides are added to tissue culture media for these cell types at a

concentration of about 10 pg/ml to about 1000 ng/ml. Those skilled in the art will recognize that zveg3 proteins can be advantageously combined with other growth factors in culture media.

Within the laboratory research field, zveg3 proteins can also be
5 used as molecular weight standards; as reagents in assays for determining circulating levels of the protein, such as in the diagnosis of disorders characterized by over- or under-production of zveg3 protein; or as standards in the analysis of cell phenotype.

Zveg3 proteins can also be used to identify inhibitors of their
10 activity. Test compounds are added to the assays disclosed above to identify compounds that inhibit the activity of zveg3 protein. In addition to those assays disclosed above, samples can be tested for inhibition of zveg3 activity within a variety of assays designed to measure receptor binding or the stimulation/inhibition of zveg3-dependent cellular responses. For example,
15 zveg3-responsive cell lines can be transfected with a reporter gene construct that is responsive to a zveg3-stimulated cellular pathway. Reporter gene constructs of this type are known in the art, and will generally comprise a zveg3-activated serum response element (SRE) operably linked to a gene encoding an assayable protein, such as luciferase. Candidate compounds,
20 solutions, mixtures or extracts are tested for the ability to inhibit the activity of zveg3 on the target cells as evidenced by a decrease in zveg3 stimulation of reporter gene expression. Assays of this type will detect compounds that directly block zveg3 binding to cell-surface receptors, as well as compounds that block processes in the cellular pathway subsequent to receptor-ligand
25 binding. In the alternative, compounds or other samples can be tested for direct blocking of zveg3 binding to receptor using zveg3 tagged with a detectable label (e.g., ¹²⁵I, biotin, horseradish peroxidase, FITC, and the like). Within assays of this type, the ability of a test sample to inhibit the binding of labeled zveg3 to the receptor is indicative of inhibitory activity, which can be
30 confirmed through secondary assays. Receptors used within binding assays may be cellular receptors or isolated, immobilized receptors.

The activity of zveg3 proteins can be measured with a silicon-based biosensor microphysiometer that measures the extracellular acidification rate or proton excretion associated with receptor binding and subsequent physiologic cellular responses. An exemplary such device is the Cytosensor™
5 Microphysiometer manufactured by Molecular Devices, Sunnyvale, CA. A variety of cellular responses, such as cell proliferation, ion transport, energy production, inflammatory response, regulatory and receptor activation, and the like, can be measured by this method. See, for example, McConnell et al.,
10 *Science* 257:1906-1912, 1992; Pitchford et al., *Meth. Enzymol.* 228:84-108, 1997; Arimilli et al., *J. Immunol. Meth.* 212:49-59, 1998; and Van Liefde et al.,
Eur. J. Pharmacol. 346:87-95, 1998. The microphysiometer can be used for assaying adherent or non-adherent eukaryotic or prokaryotic cells. By measuring extracellular acidification changes in cell media over time, the microphysiometer directly measures cellular responses to various stimuli,
15 including zveg3 proteins, their agonists, and antagonists. The microphysiometer can be used to measure responses of a zveg3-responsive eukaryotic cell, compared to a control eukaryotic cell that does not respond to zveg3 polypeptide. Zveg3-responsive eukaryotic cells comprise cells into which a receptor for zveg3 has been transfected creating a cell that is
20 responsive to zveg3, as well as cells naturally responsive to zveg3 such as cells derived from vascular or neural tissue. Differences, measured by a change, for example, an increase or diminution in extracellular acidification, in the response of cells exposed to zveg3 polypeptide, relative to a control not exposed to zveg3, are a direct measurement of zveg3-modulated cellular
25 responses. Moreover, such zveg3-modulated responses can be assayed under a variety of stimuli. The present invention thus provides methods of identifying agonists and antagonists of zveg3 proteins, comprising providing cells responsive to a zveg3 polypeptide, culturing a first portion of the cells in the absence of a test compound, culturing a second portion of the cells in the
30 presence of a test compound, and detecting a change, for example, an increase or diminution, in a cellular response of the second portion of the cells

as compared to the first portion of the cells. The change in cellular response is shown as a measurable change in extracellular acidification rate. Culturing a third portion of the cells in the presence of a zveg3 protein and the absence of a test compound provides a positive control for the zveg3-responsive cells and
5 a control to compare the agonist activity of a test compound with that of the zveg3 polypeptide. Antagonists of zveg3 can be identified by exposing the cells to zveg3 protein in the presence and absence of the test compound, whereby a reduction in zveg3-stimulated activity is indicative of antagonist activity in the test compound.

10 Zveg3 proteins can also be used to identify cells, tissues, or cell lines that respond to a zveg3-stimulated pathway. The microphysiometer, described above, can be used to rapidly identify ligand-responsive cells, such as cells responsive to zveg3 proteins. Cells are cultured in the presence or absence of zveg3 polypeptide. Those cells that elicit a measurable change in
15 extracellular acidification in the presence of zveg3 are responsive to zveg3. Responsive cells can then be used to identify antagonists and agonists of zveg3 polypeptide as described above.

Inhibitors of zveg3 activity (zveg3 antagonists) include anti-zveg3 antibodies, soluble zveg3 receptors (including soluble PDGF alpha
20 receptor; see, e.g., Herren et al., J. Biol. Chem. 268:15088-15095, 1993), anti-receptor antibodies, and other peptidic and non-peptidic agents, including ribozymes, small molecule inhibitors, and angiogenically or mitogenically inactive receptor-binding fragments of zveg3 polypeptides. Such antagonists can be used to block the mitogenic, chemotactic, or angiogenic effects of
25 zveg3. These antagonists may therefore be useful in reducing the growth of solid tumors by inhibiting neovascularization of the developing tumor, by directly blocking tumor cell growth, or by promoting apoptosis through the inhibition of PDGF alpha receptor-mediated processes. For example, experimental evidence indicates that zveg3 is produced by glioblastoma cells,
30 suggesting that inhibition of zveg3 activity may be useful in the treatment of these tumors. Other uses of zveg3 antagonists include treating diabetic

retinopathy, psoriasis, arthritis, and scleroderma; and reducing fibrosis, including scar formation, keloids, liver fibrosis, lung fibrosis (e.g., silicosis, asbestosis), kidney fibrosis (including diabetic nephropathy), and glomerulosclerosis. Inhibitors of zveg3 may also be useful in the treatment of proliferative vascular disorders wherein zveg3 activity is pathogenic. Such disorders may include atherosclerosis and intimal hyperplastic restenosis following angioplasty, endarterectomy, vascular grafting, organ transplant, or vascular stent emplacement. These conditions involve complex growth factor-mediated responses wherein certain factors may be beneficial to the clinical outcome and others may be pathogenic.

Inhibitors of zveg3 may also prove useful in the treatment of ocular neovascularization, including diabetic retinopathy and age-related macular degeneration. Experimental evidence suggests that these conditions result from the expression of angiogenic factors induced by hypoxia in the retina.

Zveg3 antagonists are also of interest in the treatment of inflammatory disorders, such as rheumatoid arthritis and psoriasis. In rheumatoid arthritis, studies suggest that VEGF plays an important role in the formation of pannus, an extensively vascularized tissue that invades and destroys cartilage. Psoriatic lesions are hypervascular and overexpress the angiogenic polypeptide IL-8.

Zveg3 antagonists may also prove useful in the treatment of infantile hemangiomas, which exhibit overexpression of VEGF and bFGF during the proliferative phase.

Inhibitors are formulated for pharmaceutical use as generally disclosed above, taking into account the precise chemical and physical nature of the inhibitor and the condition to be treated. The relevant determinations are within the level of ordinary skill in the formulation art. Other angiogenic and vasculogenic factors, including VEGF and bFGF, have been implicated in pathological neovascularization. In such instances it may be advantageous to combine a zveg3 inhibitor with one or more inhibitors of these other factors.

The polypeptides, nucleic acids, and antibodies of the present invention may be used in diagnosis or treatment of disorders associated with cell loss or abnormal cell proliferation, including cancer, impaired or excessive vasculogenesis or angiogenesis, and diseases of the nervous system.

5 Labeled zveg3 polypeptides may be used for imaging tumors or other sites of abnormal cell proliferation. In view of the binding of zveg3 to the PDGF alpha receptor, zveg3 antagonists may be useful in the treatment of tumors that express this receptor (e.g., glioblastomas, malignant melanomas). Because angiogenesis in adult animals is generally limited to wound healing and the
10 female reproductive cycle, it is a very specific indicator of pathological processes. Angiogenesis is indicative of, for example, developing solid tumors, retinopathies, and arthritis.

Zveg3 polypeptides and anti-zveg3 antibodies can be directly or indirectly conjugated to drugs, toxins, radionuclides and the like, and these
15 conjugates used for *in vivo* diagnostic or therapeutic applications. For instance, polypeptides or antibodies of the present invention may be used to identify or treat tissues or organs that express a corresponding anti-complementary molecule (receptor or antigen, respectively, for instance). More specifically, zveg3 polypeptides or anti-zveg3 antibodies, or bioactive
20 fragments or portions thereof, can be coupled to detectable or cytotoxic molecules and delivered to a mammal having cells, tissues, or organs that express the anti-complementary molecule. For example, the CUB domain of zveg3 can be used to target peptidic and non-peptidic moieties to semaphorins as disclosed above.

25 Suitable detectable molecules can be directly or indirectly attached to the polypeptide or antibody, and include radionuclides, enzymes, substrates, cofactors, inhibitors, fluorescent markers, chemiluminescent markers, magnetic particles, and the like. Suitable cytotoxic molecules can be directly or indirectly attached to the polypeptide or antibody, and include
30 bacterial or plant toxins (for instance, diphtheria toxin, *Pseudomonas* exotoxin, ricin, abrin, saporin, and the like), as well as therapeutic radionuclides, such as

iodine-131, rhenium-188 or yttrium-90. These can be either directly attached to the polypeptide or antibody, or indirectly attached according to known methods, such as through a chelating moiety. Polypeptides or antibodies can also be conjugated to cytotoxic drugs, such as adriamycin. For indirect
5 attachment of a detectable or cytotoxic molecule, the detectable or cytotoxic molecule may be conjugated with a member of a complementary/anticomplementary pair, where the other member is bound to the polypeptide or antibody portion. For these purposes, biotin/streptavidin is an exemplary complementary/anticomplementary pair.

10 In another embodiment, polypeptide-toxin fusion proteins or antibody/fragment-toxin fusion proteins may be used for targeted cell or tissue inhibition or ablation, such as in cancer therapy. Of particular interest in this regard are conjugates of a zveg3 polypeptide and a cytotoxin, which can be used to target the cytotoxin to a tumor or other tissue that is undergoing
15 undesired cell replication and modification.

In another embodiment, zveg3-cytokine fusion proteins or antibody/fragment-cytokine fusion proteins may be used for enhancing *in vitro* cytotoxicity (for instance, that mediated by monoclonal antibodies against tumor targets) and for enhancing *in vivo* killing of target tissues (for example,
20 blood and bone marrow cancers). See, generally, Hornick et al., *Blood* 89:4437-4447, 1997). In general, cytokines are toxic if administered systemically. The described fusion proteins enable targeting of a cytokine to a desired site of action, such as a cell having binding sites for zveg3, thereby providing an elevated local concentration of cytokine. Suitable cytokines for
25 this purpose include, for example, interleukin-2 and granulocyte-macrophage colony-stimulating factor (GM-CSF). Such fusion proteins may be used to cause cytokine-induced killing of tumors and other tissues exhibiting undesired cell replication.

In yet another embodiment, a zveg3 polypeptide or anti-zveg3
30 antibody can be conjugated with a radionuclide, particularly with a beta-emitting or gamma-emitting radionuclide, and used to reduce restenosis. For

instance, iridium-192 impregnated ribbons placed into stented vessels of patients until the required radiation dose was delivered resulted in decreased tissue growth in the vessel and greater luminal diameter than the control group, which received placebo ribbons. Further, revascularisation and stent
5 thrombosis were significantly lower in the treatment group. Similar results are predicted with targeting of a bioactive conjugate containing a radionuclide, as described herein.

The bioactive polypeptide or antibody conjugates described herein can be delivered intravenously, intra-arterially or intraductally, or may be
10 introduced locally at the intended site of action.

Polynucleotides encoding zveg3 polypeptides are useful within gene therapy applications where it is desired to increase or inhibit zveg3 activity. For example, Isner et al., *The Lancet* (ibid.) reported that VEGF gene therapy promoted blood vessel growth in an ischemic limb. Additional
15 applications of zveg3 gene therapy include stimulation of wound healing, repopulation of vascular grafts, stimulation of neurite growth, and inhibition of cancer growth and metastasis.

The present invention also provides polynucleotide reagents for diagnostic use. For example, a zveg3 gene, a probe comprising zveg3 DNA
20 or RNA, or a subsequence thereof can be used to determine if the zveg3 gene is present on chromosome 4 of a human patient or if a mutation has occurred. Detectable chromosomal aberrations at the zveg3 gene locus include, but are not limited to, aneuploidy, gene copy number changes, insertions, deletions, restriction site changes and rearrangements. Such aberrations can be
25 detected using polynucleotides of the present invention by employing molecular genetic techniques, such as restriction fragment length polymorphism (RFLP) analysis, short tandem repeat (STR) analysis employing PCR techniques, and other genetic linkage analysis techniques known in the art (Sambrook et al., *ibid.*; Ausubel et. al., *ibid.*; A.J. Marian, *Chest* 108:255-
30 265, 1995).

Radiation hybrid mapping is a somatic cell genetic technique developed for constructing high-resolution, contiguous maps of mammalian chromosomes (Cox et al., *Science* 250:245-250, 1990). Partial or full knowledge of a gene's sequence allows one to design PCR primers suitable for use with chromosomal radiation hybrid mapping panels. Commercially available radiation hybrid mapping panels that cover the entire human genome, such as the Stanford G3 RH Panel and the GeneBridge 4 RH Panel (Research Genetics, Inc., Huntsville, AL), are available. These panels enable rapid, PCR-based chromosomal localizations and ordering of genes, sequence-tagged sites (STSs), and other nonpolymorphic and polymorphic markers within a region of interest. This technique allows one to establish directly proportional physical distances between newly discovered genes of interest and previously mapped markers. The precise knowledge of a gene's position can be useful for a number of purposes, including: 1) determining relationships between short sequences and obtaining additional surrounding genetic sequences in various forms, such as YACs, BACs or cDNA clones; 2) providing a possible candidate gene for an inheritable disease which shows linkage to the same chromosomal region; and 3) cross-referencing model organisms, such as mouse, which may aid in determining what function a particular gene might have.

The invention is further illustrated by the following non-limiting examples.

Example 1

Clones comprising portions of the *zveg3* coding sequence were identified in public and proprietary databases of expressed sequence tags (ESTs). A first clone, corresponding to an EST in a proprietary database, was obtained and sequenced. It contained a 2350-bp insert with an open reading frame of about 800 bp. The 5' end of the ORF was missing. A second clone, corresponding to an EST in a public database, was then sequenced, but it didn't extend the sequence obtained from the first clone. A third clone, from a

proprietary database, was then sequenced. This clone contained approximately 156 bp more than the first clone, but was also missing the 5' end.

Northern blots were performed using a series of Northern blots (Multiple Tissue Northern Blots, Clontech Laboratories, Inc., Palo Alto, CA). An approximately 400-bp DNA probe, based directly on the identified EST, was generated by digestion of a clone corresponding to the first proprietary EST with EcoRI and BglII. The DNA probe was gel-purified using a spin column containing a silica gel membrane (QIAquick™ Gel Extraction Kit; Qiagen, Inc., Valencia, CA). The probe was radioactively labeled with ³²P using a commercially available random-prime labeling kit (Rediprime™ II, Amersham Corp., Arlington Heights, IL) according to the manufacturer's specifications. The probe was purified using a push column (NucTrap® column; Stratagene, La Jolla, CA; see U.S. Patent No. 5,336,412). A commercially available hybridization solution (ExpressHyb™ Hybridization Solution; Clontech Laboratories, Inc., Palo Alto, CA) was used for prehybridization and as a hybridizing solution for the Northern blots. Hybridization took place overnight at 65°C, and the blots were then washed 4 times in 2X SSC and 0.1% SDS at room temperature, followed by two washes in 0.1X SSC and 0.1% SDS at 50°C and one wash in 0.1X SSC and 0.1% SDS at 56°C. The Northern blots were then exposed to film overnight at -80°C and for three days at -80°C. One transcript size was seen in all tissues at approximately 4.0 kb. Signal intensity was highest in thyroid, spinal cord and adrenal gland. Signal intensity was average in heart, kidney, pancreas, prostate, ovary, stomach, and trachea. Signal intensity was weak in all other tissues.

Northern blot analysis was performed using Mouse Multiple Tissue Blots obtained from Clontech Laboratories and Invitrogen (Carlsbad, CA). An approximately 400 bp DNA probe, based directly on the identified EST, was generated by digestion of a clone corresponding to the first proprietary EST with EcoRI and BglII. The DNA probe was gel purified using a spin column as disclosed above. The probe was radioactively labeled with ³²P

by random priming as disclosed above. The probe was purified using a NucTrap® push column. Hybridization conditions were as disclosed above. The blots were washed four times in 2X SSC, 0.1% SDS, then twice in 1X SSC at 50°C, then exposed to film overnight at -80°C and for three days at -80°C.

5 One transcript size of approximately 3.0 kb was seen in 7-day embryo. Three-day exposure showed bands of less intensity in 11-, 15-, and 17-day embryo. The intensity decreased with the age of the embryo. Dot blots showed a spot in 17-day embryo, and with-three day exposure in submaxillary gland. A potential band of approximately 2.0 kb was seen in testis after three days.

10 Southern blot analysis was performed using a pre-made Southern blot containing EcoRI-digested genomic DNA from nine different eukaryotic species (ZOO-BLOT from Clontech Laboratories). An approximately 400 bp DNA probe, based directly on the first proprietary database EST, was generated by digestion of the corresponding clone with
15 EcoRI and BglII. The DNA probe was gel purified using a spin column. The probe was radioactively labeled with ³²P by random priming and purified using a push column. Hybridization and wash conditions were as disclosed above for the Mouse Multiple Tissue Blots. The Northern blots were then exposed to film overnight at -80°C and for three days at -80°C. Strong bands were seen in
20 rabbit, mouse, rat, and monkey.

Example 2

A human salivary gland library was screened for a full-length clone of zveg3 by PCR. This library was an arrayed library representing 9.6 X
25 10⁵ clones made in the vector pZP5x. The vector pZP5x is the same as vector pZP-9 (deposited with American Type Culture Collection, 10801 University Blvd., Manassas, VA under Accession Number 98668), but contains a cytomegalovirus promoter instead of a metallothionein promoter between the Asp718 and BamHI sites. The plasmid thus comprises a dihydrofolate
30 reductase gene under control of the SV40 early promoter and SV40 polyadenylation site, and a cloning site to insert the gene of interest under

control of the CMV promoter and the human growth hormone (hGH) gene polyadenylation site. The working plate containing 80 pools of 12,000 colonies each was screened by PCR using oligonucleotide primers ZC19,045 (SEQ ID NO:25) and ZC19,047 (SEQ ID NO:26) with an annealing temperature of 60°C for 35 cycles. There were two strong positives, pools 58 (T-8 F1-F12) and 77 (T-7 H1-H12). The corresponding pools in the transfer plate were then screened by PCR using the same conditions. Two positives were obtained at the transfer level. The positives were T-7 H11 and T-8 F10. 5' RACE reactions were done on the transfer plate pools, and the fragments were sequenced to check zveg3 sequence and determine if a full-length clone was present. For PCR, oligonucleotide primers ZC12,700 (SEQ ID NO:27) and ZC19,045 (SEQ ID NO:25) were used at an annealing temperature of 61°C for 5 cycles, then 55°C for 30 cycles. Sequencing showed that the pool T-7 H11 had a frameshift. Transfer plate 8 pool F10 sequence appeared to be correct, so this pool of DNA was used in filter lifts.

Pool F10 from transfer plate 8 was plated and filter lifted using nylon membranes (Hybond-N™; Amersham Corporation). Approximately 1200 colonies per plate on each of 5 filters were lifted for a total of approximately 6000 colonies. The filters were marked with a hot needle for orientation, then denatured for 6 minutes in 0.5 M NaOH and 1.5 M Tris-HCl, pH 7.2. The filters were then neutralized in 1.5 M NaCl and 0.5 M Tris-HCl, pH 7.2 for 6 minutes. The DNA was affixed to the filters using a UV crosslinker (Stratalinker®, Stratagene, La Jolla, CA) at 1200 joules. The filters were prewashed at 65°C in prewash buffer consisting of 0.25X SSC, 0.25% SDS, and 1mM EDTA. The solution was changed a total of three times over a 45-minute period to remove cell debris. Filters were prehybridized for approximately 3 hours at 65°C in 25 ml of ExpressHyb™. The probe was generated using an approximately 400-bp fragment produced by digestion of the first proprietary database clone with EcoRI and BglII and gel-purified using a spin column as disclosed above. The probe was radioactively labeled with ³²P by random priming as disclosed above and purified using a push column. ExpressHyb™ solution was used for the

hybridizing solution for the filters. Hybridization took place overnight at 65°C. Blots were rinsed 2X in 65°C solution 1 (2X SSC, 0.1% SDS), then washed 4 times in solution 1 at 65°C. The filters were exposed to film overnight at -80°C. There were 14 positives on the filters. 85 clones were picked from the positive areas and screened by PCR using oligonucleotide primers ZC19,045 (SEQ ID NO:25) and ZC19,047 (SEQ ID NO:26) and an annealing temperature of 60°C. Thirteen positives were obtained and streaked out for individual clones. Twenty-four colonies were picked and checked by PCR as previously described. Six positives were obtained, two of which were sequenced. Both sequences were the same and full length. The sequence is shown in SEQ ID NO:1.

Example 3

The human zveg3 gene was mapped to chromosome 4 using the commercially available GeneBridge 4 Radiation Hybrid Panel (Research Genetics, Inc., Huntsville, AL). The GeneBridge 4 Radiation Hybrid Panel contains PCRable DNAs from each of 93 radiation hybrid clones, plus two control DNAs (the HFL donor and the A23 recipient). A publicly available WWW server (<http://www-genome.wi.mit.edu/cgi-bin/contig/rhmapper.pl>) allows mapping relative to the Whitehead Institute/MIT Center for Genome Research's radiation hybrid map of the human genome (the "WICGR" radiation hybrid map), which was constructed with the GeneBridge 4 Radiation Hybrid Panel.

For the mapping of Zveg3 with the GeneBridge 4 RH Panel, 20- μ l reactions were set up in a PCRable 96-well microtiter plate (Stratagene, La Jolla, CA) and used in a thermal cycler (RoboCycler® Gradient 96; Stratagene). Each of the 95 PCR reactions contained 2 μ l 10X PCR reaction buffer (Clontech Laboratories, Inc., Palo Alto, CA), 1.6 μ l dNTPs mix (2.5 mM each, PERKIN-ELMER, Foster City, CA), 1 μ l sense primer ZC 20,368 (SEQ ID NO:28), 1 μ l antisense primer ZC 20,369 (SEQ ID NO:29), 2 μ l of a density increasing agent and tracking dye (RediLoad, Research Genetics, Inc., Huntsville, AL), 0.4 μ l of a commercially available DNA polymerase/antibody

mix (50X Advantage™ KlenTaq Polymerase Mix, obtained from Clontech Laboratories, Inc.), 25 ng of DNA from an individual hybrid clone or control, and x µl ddH₂O for a total volume of 20 µl. The reaction mixtures were overlaid with an equal amount of mineral oil and sealed. The PCR cycycler conditions were as follows: an initial 4-minute denaturation at 94°C; 35 cycles of 45 seconds denaturation at 94°C, 45 seconds annealing at 56°C, and 75 seconds extension at 72°C; followed by a final extension of 7 minutes at 72°C. The reaction products were separated by electrophoresis on a 2% agarose gel (Life Technologies, Gaithersburg, MD).

The results showed that zveg3 maps 3.56 cR_3000 from the framework marker CHLC.GATA72A08 on the chromosome 4 WICGR radiation hybrid map. Proximal and distal framework markers were CHLC.GATA72A08 and WI-3936, respectively. The use of surrounding markers positions the zveg3 gene in the 4q28.3 region on the integrated LDB chromosome 4 map (The Genetic Location Database, University of Southampton, WWW server: http://cedar.genetics.soton.ac.uk/public_html/).

Using substantially the same methods, the mouse zveg3 gene was mapped to chromosome 3, linked to the framework marker D3Mit212 located at 39.7 cM. This region is syntenic with the human zveg3 locus.

Example 4

A PCR panel was screened for mouse zveg3 DNA. The panel contained 8 cDNA samples from brain, bone marrow, 15-day embryo, testis, salivary gland, placenta, 15-day embryo (Clontech Laboratories), and 17-day embryo (Clontech Laboratories) libraries.

PCR mixtures contained oligonucleotide primers zc21,222 (SEQ ID NO:38) and zc21,224 (SEQ ID NO:39). The reaction was run at an annealing temperature of 66°C with an extension time of 2 minutes for a total of 35 cycles using Ex Taq™ DNA polymerase (PanVera, Madison, WI) plus antibody. DNA samples found to be positive for zveg3 by PCR and confirmed by sequencing included mouse 15-day embryo library total pool cDNA, mouse

15-day embryo (Clontech Laboratories) and 17-day embryo (both obtained from Clontech Laboratories), mouse salivary gland library total pool cDNA, and mouse testis library total pool cDNA. Fragments of about 600 bp from each of the mouse 15-day embryo library total pool cDNA, mouse 15-day embryo
5 mcDNA, and mouse 17-day embryo mcDNA PCR products were sequenced. Sequence from the mouse 17-day embryo mcDNA and mouse 15-day embryo library total pool cDNA products confirmed the fragments to be mouse zveg3 DNA.

The mouse 15-day embryo library was screened for full-length
10 zveg3 DNA. This library was an arrayed library representing 9.6×10^5 clones in the pCMV•SPORT 2 vector (Life Technologies, Gaithersburg, MD). The working plate, containing 80 pools of 12,000 colonies each, was screened by PCR using oligonucleotide primers zc21,223 (SEQ ID NO:40) and zc21,224 (SEQ ID NO:39) with an annealing temperature of 66°C for 35 cycles.
15 Eighteen positives were obtained. Fragments from four pools (A2, A10, B2, and C4) were sequenced; all were confirmed to encode zveg3. Additional rounds of screening using the same reaction conditions and pools from the working and source plates identified one positive pool (5D).

Positive colonies were screened by hybridization. Pool 5D from
20 original source plate #5 was plated at about 250 colonies per plate and transferred to nylon membranes (Hybond-N™; Amersham Corporation, Arlington Heights, IL). Five filters were lifted for a total of ~1250 colonies. The filters were marked with a hot needle for orientation, then denatured for 6 minutes in 0.5 M NaOH and 1.5 M Tris-HCl, pH 7.2. The filters were then
25 neutralized in 1.5 M NaCl and 0.5 M Tris-HCl, pH 7.2 for 6 minutes. The DNA was fixed to the filters using a UV crosslinker (Stratalinker®, Stratagene, La Jolla, CA) at 1200 joules. A probe was generated by PCR using oligonucleotide primers zc21,223 (SEQ ID NO:40) and zc21,224 (SEQ ID NO:39), and a mouse 15-day embryo template at an annealing temperature of
30 66°C for 35 cycles. The PCR fragment was gel purified using a spin column containing a silica gel membrane (QIAquick™ Gel Extraction Kit; Qiagen, Inc.,

Valencia, CA). The DNA was radioactively labeled with ^{32}P using a commercially available kit (Rediprime™ II random-prime labeling system; Amersham Corp., Arlington Heights, IL) according to the manufacturer's specifications. The probe was purified using a commercially available push
5 column (NucTrap® column; Stratagene, La Jolla, CA; see U.S. Patent No. 5,336,412). The filters were prewashed at 65°C in prewash buffer consisting of 0.25X SSC, 0.25% SDS and 1mM EDTA. The solution was changed a total of three times over a 45-minute period to remove cell debris. Filters were prehybridized overnight at 65°C in 25 ml of a hybridization solution
10 (ExpressHyb™ Hybridization Solution; Clontech Laboratories, Inc., Palo Alto, CA), then hybridized overnight at 65° C in the same solution. Filters were rinsed twice at 65°C in pre-wash buffer (0.25X SSC, 0.25% SDS, and 1mM EDTA), then washed twice in pre-wash buffer at 65°C. Filters were exposed to film for 2 days at -80°C. There were 10 positives on the filters. 3 clones were
15 picked from the positive areas, streaked out, and 15 individual colonies from these three positives were screened by PCR using primers zc21,223 (SEQ ID NO:40) and zc21,334 (SEQ ID NO:41) at an annealing temp of 66°C. Two positives were recovered and sequenced. Both sequences were found to be the same and encoded full-length mouse zveg3 (SEQ ID NO:42).

20 The amino acid sequence is highly conserved between mouse and human zveg3s, with an overall amino acid sequence identity of 87%. The secretory peptide, CUB domain, inter-domain, and growth factor domain have 82%, 92%, 79% and 94% amino acid identity, respectively.

25 Example 5

Northern blotting was performed using Mouse Multiple Tissue Blots from Origene, Rockville, MD and Clontech, Palo Alto, CA. An approximately 800-bp DNA probe was generated by PCR using primer zc21,223 (SEQ ID NO:40) for the 5' end and primer zc21,224 (SEQ ID NO:39)
30 for the 3' end. The reaction was run for 35 cycles at an annealing temperature of 66°C using Ex Taq™ DNA polymerase (PanVera). The reaction product was

gel purified using a spin column containing a silica gel membrane and labeled with ^{32}P using a commercially available labeling kit (Multiprime™ DNA labeling system, Amersham Corp.) according to the manufacturer's specifications. The labeled probe was purified using a push column. A commercially available hybridization solution (ExpressHyb™ Hybridization Solution; Clontech Laboratories, Inc.) was used for prehybridization and as a hybridizing solution for the Northern blots. Hybridization took place overnight at 65°C, then the blots were washed 4 times in 2X SCC and 0.05% SDS at room temperature, followed by two washes in 0.1X SSC and 0.1% SDS at 50°C. The blots were then exposed to film two days and overnight at -80°C. Multiple transcript sizes were observed. Transcripts of ~3.5 and ~4.0 kb were seen in 7, 11, 15 and 17-day embryo, with the strongest at 7 days and tapering off at 17 days. A ~3.0 kb transcript was seen in kidney, liver, brain, and possibly testis. A ~1.0 kb transcript was seen in testis, muscle, and spleen. Dot blots corresponded to the northern blots.

Example 6

To make transgenic animals expressing zveg3 genes requires adult, fertile males (studs) (B6C3f1, 2-8 months of age (Taconic Farms, Germantown, NY)), vasectomized males (duds) (B6D2f1, 2-8 months, (Taconic Farms)), prepubescent fertile females (donors) (B6C3f1, 4-5 weeks, (Taconic Farms)) and adult fertile females (recipients) (B6D2f1, 2-4 months, (Taconic Farms)).

The donors are acclimated for 1 week, then injected with approximately 8 IU/mouse of Pregnant Mare's Serum gonadotrophin (Sigma, St. Louis, MO) I.P., and 46-47 hours later, 8 IU/mouse of human Chorionic Gonadotropin (hCG (Sigma)) I.P. to induce superovulation. Donors are mated with studs subsequent to hormone injections. Ovulation generally occurs within 13 hours of hCG injection. Copulation is confirmed by the presence of a vaginal plug the morning following mating.

Fertilized eggs are collected under a surgical scope (Leica MZ12 Stereo Microscope, Leica, Wetzlar, Germany). The oviducts are collected and eggs are released into urinanalysis slides containing hyaluronidase (Sigma Chemical Co.). Eggs are washed once in hyaluronidase, and twice in
 5 Whitten's W640 medium (Table 7; all reagents available from Sigma Chemical Co.) that has been incubated with 5% CO₂, 5% O₂, and 90% N₂ at 37°C. The eggs are stored in a 37°C/5% CO₂ incubator until microinjection.

Table 7

	<u>mgs/200 ml</u>	<u>mgs/500 ml</u>
NaCl	1280	3200
KCl	72	180
KH ₂ PO ₄	32	80
MgSO ₄ · 7H ₂ O	60	150
Glucose	200	500
Ca ²⁺ Lactate	106	265
Benzylpenicillin	15	37.5
Streptomycin SO ₄	10	25
NaHCO ₃	380	950
Na Pyruvate	5	12.5
H ₂ O	200 ml	500 ml
500 mM EDTA	100 µl	250 µl
5% Phenol Red	200 µl	500 µl
BSA	600	1500

10

Zveg3 cDNA is inserted into the expression vector pHB12-8 (see Fig. 2). Vector pHB12-8 was derived from p2999B4 (Palmiter et al., *Mol. Cell Biol.* 13:5266-5275, 1993) by insertion of a rat insulin II intron (ca. 200 bp) and polylinker (Fse I/Pme I/Asc I) into the Nru I site. The vector comprises a
 15 mouse metallothionein (MT-1) promoter (ca. 750 bp) and human growth hormone (hGH) untranslated region and polyadenylation signal (ca. 650 bp)

flanked by 10 kb of MT-1 5' flanking sequence and 7 kb of MT-1 3' flanking sequence. The cDNA is inserted between the insulin II and hGH sequences.

10-20 micrograms of plasmid DNA is linearized, gel-purified, and resuspended in 10 mM Tris pH 7.4, 0.25 mM EDTA pH 8.0, at a final
5 concentration of 5-10 nanograms per microliter for microinjection.

Plasmid DNA is microinjected into harvested eggs contained in a drop of W640 medium overlaid by warm, CO₂-equilibrated mineral oil. The DNA is drawn into an injection needle (pulled from a 0.75mm ID, 1mm OD borosilicate glass capillary) and injected into individual eggs. Each egg is
10 penetrated with the injection needle into one or both of the haploid pronuclei.

Picoliters of DNA are injected into the pronuclei, and the injection needle is withdrawn without coming into contact with the nucleoli. The procedure is repeated until all the eggs are injected. Successfully microinjected eggs are transferred into an organ tissue-culture dish with
15 pregassed W640 medium for storage overnight in a 37°C/5% CO₂ incubator.

The following day, 2-cell embryos are transferred into pseudopregnant recipients. The recipients are identified by the presence of copulation plugs, after copulating with vasectomized duds. Recipients are anesthetized and shaved on the dorsal left side and transferred to a surgical
20 microscope. A small incision is made in the skin and through the muscle wall in the middle of the abdominal area outlined by the ribcage, the saddle, and the hind leg, midway between knee and spleen. The reproductive organs are exteriorized onto a small surgical drape. The fat pad is stretched out over the surgical drape, and a baby serrefine (Roboz, Rockville, MD) is attached to the
25 fat pad and left hanging over the back of the mouse, preventing the organs from sliding back in.

With a fine transfer pipette containing mineral oil followed by alternating W640 and air bubbles, 12-17 healthy 2-cell embryos from the previous day's injection are transferred into the recipient. The swollen ampulla
30 is located and holding the oviduct between the ampulla and the bursa, and a

nick in the oviduct is made with a 28 g needle close to the bursa, making sure not to tear the ampulla or the bursa.

The pipette is transferred into the nick in the oviduct, and the embryos are blown in, allowing the first air bubble to escape the pipette. The fat pad is gently pushed into the peritoneum, and the reproductive organs are allowed to slide in. The peritoneal wall is closed with one suture, and the skin is closed with a wound clip. The mice recuperate on a 37°C slide warmer for a minimum of 4 hours.

The recipients are returned to cages in pairs, and allowed 19-21 days gestation. After birth, 19-21 days postpartum is allowed before weaning. The weanlings are sexed and placed into separate sex cages, and a 0.5 cm biopsy (used for genotyping) is snipped off the tail with clean scissors.

Genomic DNA is prepared from the tail snips using a commercially available kit (DNeasy™ 96 Tissue Kit; Qiagen, Valencia, CA) following the manufacturer's instructions. Genomic DNA is analyzed by PCR using primers designed to the human growth hormone (hGH) 3' UTR portion of the transgenic vector. The use of a region unique to the human sequence (identified from an alignment of the human and mouse growth hormone 3' UTR DNA sequences) ensures that the PCR reaction does not amplify the mouse sequence. Primers zc17,251 (SEQ ID NO:30) and zc17,252 (SEQ ID NO:31) amplify a 368-base-pair fragment of hGH. In addition, primers zc17,156 (SEQ ID NO:32) and zc17,157 (SEQ ID NO:33), which hybridize to vector sequences and amplify the cDNA insert, may be used along with the hGH primers. In these experiments, DNA from animals positive for the transgene will generate two bands, a 368-base-pair band corresponding to the hGH 3' UTR fragment and a band of variable size corresponding to the cDNA insert.

Once animals are confirmed to be transgenic (TG), they are back-crossed into an inbred strain by placing a TG female with a wild-type male, or a TG male with one or two wild-type female(s). As pups are born and weaned, the sexes are separated, and their tails snipped for genotyping.

To check for expression of a transgene in a live animal, a partial hepatectomy is performed. A surgical prep is made of the upper abdomen directly below the xiphoid process. Using sterile technique, a small 1.5-2 cm incision is made below the sternum, and the left lateral lobe of the liver is exteriorized. Using 4-0 silk, a tie is made around the lower lobe securing it outside the body cavity. An atraumatic clamp is used to hold the tie while a second loop of absorbable Dexon (American Cyanamid, Wayne, N.J.) is placed proximal to the first tie. A distal cut is made from the Dexon tie, and approximately 100 mg of the excised liver tissue is placed in a sterile petri dish. The excised liver section is transferred to a 14-ml polypropylene round bottom tube, snap frozen in liquid nitrogen, and stored on dry ice. The surgical site is closed with suture and wound clips, and the animal's cage is placed on a 37°C heating pad for 24 hours post-operatively. The animal is checked daily post-operatively, and the wound clips are removed 7-10 days after surgery.

Analysis of the mRNA expression level of each transgene is done using an RNA solution hybridization assay or real-time PCR on an ABI Prism 7700 (PE Applied Biosystems, Inc., Foster City, CA) following the manufacturer's instructions.

An adenovirus vector was prepared using a liver-specific albumin gene enhancer and basal promoter (designated "AEO promoter"). The albumin promoter construct (designated pAEO) was constructed by inserting a 2.2 kb Not1/EcoRV fragment from pALBdelta2L (Pinkert et al., *Genes Dev.* 1:268-276, 1987) and an 850 bp Nrul/Not1 DNA segment comprising the rat insulin II intron, an FseI/PmeI/Ascl polylinker, and the human growth hormone poly A sequence into a commercially available phagemid vector (pBluescript® KS(+); Stratagene, La Jolla, CA). For microinjection, the plasmid is digested with Not1 to liberate the expression cassette.

An additional adenovirus vector was constructed using an epithelial cell-specific keratin gene (K14) promoter (Vassar et al., *Proc. Natl. Acad. Sci. USA* 86:1563-1567, 1989). The 1038-bp open reading frame encoding full-length human zveg3 was amplified by PCR so as to introduce an

optimized initiation codon and flanking 5' *PmeI* and 3' *Ascl* sites using the primers ZC20,180 (SEQ ID NO:34) and ZC20,181 (SEQ ID NO:35). The resulting *PmeI/Ascl* fragment was subcloned into the polylinker of pKFO114, a basal keratinocyte-restricted transgenic vector comprising the human keratin 14 (K14) promoter (an approximately 2.3 Kb fragment amplified from human genomic DNA [obtained from Clontech Laboratories, Inc.] based on the sequence of Staggars et al., "Sequence of the promoter for the epidermal keratin gene, K14", GenBank accession #U11076, 1994), followed by a heterologous intron (a 294-bp *BstXI/PstI* fragment from pIRES1hyg (Clontech Laboratories, Inc.; see, Huang and Gorman, *Nucleic Acids Res.* 18:937-947, 1990), a *PmeI/Ascl* polylinker, and the human growth hormone gene polyadenylation signal (a 627 bp *SmaI/EcoRI* fragment; see, Seeburg, *DNA* 1:239-249, 1982). The transgene insert was separated from the plasmid backbone by *NotI* digestion and agarose gel purification, and fertilized ova from matings of B6C3F1Tac mice or inbred FVB/NTac mice were microinjected and implanted into pseudopregnant females essentially as described by Malik et al., *Molec. Cell. Biol.* 15:2349-2358, 1995. Transgenic founders were identified by PCR on genomic tail DNA using primers specific for the human growth hormone poly A signal (ZC17,252, SEQ ID NO:31; and ZC17,251, SEQ ID NO:30) to amplify a 368-bp diagnostic product. Transgenic lines were initiated by breeding founders with C57BL/6Tac or FVB/NTac mice.

Transgenic mice were generated essentially as disclosed above using MT-1, K14, and AEO promoters. Four MT-1/zvegf3 transgenic mice were generated. In one animal (female) approximately 800 molecules zvegf3 mRNA/cell were produced in the liver after zinc induction. This animal had enlargement of the liver and spleen. Also observed were proliferation of hepatic sinusoidal cells and extra-medullary hematopoiesis. One K14/zvegf3 transgenic mouse (female) showed a low level of expression with low body weight, low hematocrit, and low platelet count. One AEO/zvegf3 transgenic mouse (male) with a low level of expression exhibited liver sinusoidal cell proliferation.

Example 7

An expression plasmid containing all or part of a polynucleotide encoding zveg3 is constructed via homologous recombination. A fragment of zveg3 cDNA is isolated by PCR using the polynucleotide sequence of SEQ ID NO: 1 with flanking regions at the 5' and 3' ends corresponding to the vector sequences flanking the zveg3 insertion point. The primers for PCR each include from 5' to 3' end: 40 bp of flanking sequence from the vector and 17 bp corresponding to the amino and carboxyl termini from the open reading frame of zveg3.

Ten μ l of the 100 μ l PCR reaction is run on a 0.8% low-melting-temperature agarose (SeaPlaque GTG®; FMC BioProducts, Rockland, ME) gel with 1 x TBE buffer for analysis. The remaining 90 μ l of PCR reaction is precipitated with the addition of 5 μ l 1 M NaCl and 250 μ l of absolute ethanol. The plasmid pZMP6, which has been cut with SmaI, is used for recombination with the PCR fragment. Plasmid pZMP6 was constructed from pZP9 (deposited at the American Type Culture Collection, 10801 University Boulevard, Manassas, VA 20110-2209, under Accession No. 98668) with the yeast genetic elements taken from pRS316 (deposited at the American Type Culture Collection, 10801 University Boulevard, Manassas, VA 20110-2209, under Accession No. 77145), an internal ribosome entry site (IRES) element from poliovirus, and the extracellular domain of CD8 truncated at the C-terminal end of the transmembrane domain. pZMP6 is a mammalian expression vector containing an expression cassette having the cytomegalovirus immediate early promoter, multiple restriction sites for insertion of coding sequences, a stop codon, and a human growth hormone terminator. The plasmid also contains an *E. coli* origin of replication; a mammalian selectable marker expression unit comprising an SV40 promoter, enhancer and origin of replication, a DHFR gene, and the SV40 terminator; as well as the URA3 and CEN-ARS sequences required for selection and replication in *S. cerevisiae*.

One hundred microliters of competent yeast (*S. cerevisiae*) cells are independently combined with 10 μ l of the various DNA mixtures from above and transferred to a 0.2-cm electroporation cuvette. The yeast/DNA mixtures are electropulsed using power supply settings of 0.75 kV (5 kV/cm), ∞ ohms, 25 μ F. To each cuvette is added 600 μ l of 1.2 M sorbitol, and the yeast is plated in two 300- μ l aliquots onto two URA-D plates and incubated at 30°C. After about 48 hours, the Ura⁺ yeast transformants from a single plate are resuspended in 1 ml H₂O and spun briefly to pellet the yeast cells. The cell pellet is resuspended in 1 ml of lysis buffer (2% Triton X-100, 1% SDS, 100 mM NaCl, 10 mM Tris, pH 8.0, 1 mM EDTA). Five hundred microliters of the lysis mixture is added to an Eppendorf tube containing 300 μ l acid-washed glass beads and 200 μ l phenol-chloroform, vortexed for 1 minute intervals two or three times, and spun for 5 minutes in an Eppendorf centrifuge at maximum speed. Three hundred microliters of the aqueous phase is transferred to a fresh tube, and the DNA is precipitated with 600 μ l ethanol (EtOH), followed by centrifugation for 10 minutes at 4°C. The DNA pellet is resuspended in 10 μ l H₂O.

Transformation of electrocompetent *E. coli* host cells (Electromax DH10B™ cells; obtained from Life Technologies, Inc., Gaithersburg, MD) is done with 0.5-2 ml yeast DNA prep and 40 μ l of cells. The cells are electropulsed at 1.7 kV, 25 μ F, and 400 ohms. Following electroporation, 1 ml SOC (2% Bacto™ Tryptone (Difco, Detroit, MI), 0.5% yeast extract (Difco), 10 mM NaCl, 2.5 mM KCl, 10 mM MgCl₂, 10 mM MgSO₄, 20 mM glucose) is plated in 250- μ l aliquots on four LB AMP plates (LB broth (Lennox), 1.8% Bacto™ Agar (Difco), 100 mg/L Ampicillin).

Individual clones harboring the correct expression construct for zveg3 are identified by restriction digest to verify the presence of the zveg3 insert and to confirm that the various DNA sequences have been joined correctly to one another. The inserts of positive clones are subjected to sequence analysis. Larger scale plasmid DNA is isolated using a commercially available kit (QIAGEN Plasmid Maxi Kit, Qiagen, Valencia, CA) according to

manufacturer's instructions. The correct construct is designated pZMP6/zveg3 (Fig. 3).

Example 8

5 CHO DG44 cells (Chasin et al., *Som. Cell. Molec. Genet.* 12:555-666, 1986) are plated in 10-cm tissue culture dishes and allowed to grow to approximately 50% to 70% confluency overnight at 37°C, 5% CO₂, in Ham's F12/FBS media (Ham's F12 medium, Life Technologies), 5% fetal bovine serum (Hyclone, Logan, UT), 1% L-glutamine (JRH Biosciences, Lenexa, KS),
10 1% sodium pyruvate (Life Technologies). The cells are then transfected with the plasmid pZMP6/zveg3 by liposome-mediated transfection using a 3:1 (w/w) liposome formulation of the polycationic lipid 2,3-dioleoyloxy-N-[2(sperminecarboxamido)ethyl]-N,N-dimethyl-1-propaniminium-trifluoroacetate and the neutral lipid dioleoyl phosphatidylethanolamine in membrane-filtered
15 water (Lipofectamine™ Reagent, Life Technologies), in serum free (SF) media formulation (Ham's F12, 10 mg/ml transferrin, 5 mg/ml insulin, 2 mg/ml fetuin, 1% L-glutamine and 1% sodium pyruvate). Plasmid pZMP6/Zveg3 is diluted into 15-ml tubes to a total final volume of 640 µl with SF media. 35 µl of Lipofectamine™ is mixed with 605 µl of SF medium. The Lipofectamine™
20 mixture is added to the DNA mixture and allowed to incubate approximately 30 minutes at room temperature. Five ml of SF media is added to the DNA:Lipofectamine™ mixture. The cells are rinsed once with 5 ml of SF media, aspirated, and the DNA:Lipofectamine™ mixture is added. The cells are incubated at 37°C for five hours, then 6.4 ml of Ham's F12/10% FBS, 1%
25 PSN media is added to each plate. The plates are incubated at 37°C overnight, and the DNA:Lipofectamine™ mixture is replaced with fresh 5% FBS/Ham's media the next day. On day 3 post-transfection, the cells are split into T-175 flasks in growth medium. On day 7 postransfection, the cells are stained with FITC-anti-CD8 monoclonal antibody (Pharmingen, San Diego, CA)
30 followed by anti-FITC-conjugated magnetic beads (Miltenyi Biotec). The CD8-positive cells are separated using commercially available columns (mini-MACS

columns; Miltenyi Biotec) according to the manufacturer's directions and put into DMEM/Ham's F12/5% FBS without nucleosides but with 50 nM methotrexate (selection medium).

Cells are plated for subcloning at a density of 0.5, 1 and 5 cells per well in 96-well dishes in selection medium and allowed to grow out for approximately two weeks. The wells are checked for evaporation of medium and brought back to 200 μ l per well as necessary during this process. When a large percentage of the colonies in the plate are near confluency, 100 μ l of medium is collected from each well for analysis by dot blot, and the cells are fed with fresh selection medium. The supernatant is applied to a nitrocellulose filter in a dot blot apparatus, and the filter is treated at 100°C in a vacuum oven to denature the protein. The filter is incubated in 625 mM Tris-glycine, pH 9.1, 5mM β -mercaptoethanol, at 65°C, 10 minutes, then in 2.5% non-fat dry milk Western A Buffer (0.25% gelatin, 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 5 mM EDTA, 0.05% Igepal CA-630) overnight at 4°C on a rotating shaker. The filter is incubated with the antibody-HRP conjugate in 2.5% non-fat dry milk Western A buffer for 1 hour at room temperature on a rotating shaker. The filter is then washed three times at room temperature in PBS plus 0.01% Tween 20, 15 minutes per wash. The filter is developed with chemiluminescence reagents (ECL™ direct labelling kit; Amersham Corp., Arlington Heights, IL) according to the manufacturer's directions and exposed to film (Hyperfilm ECL, Amersham) for approximately 5 minutes. Positive clones are trypsinized from the 96-well dish and transferred to 6-well dishes in selection medium for scaleup and analysis by Western blot.

25

Example 9

Full-length zveg3 protein was produced in BHK cells transfected with pZMP6/zveg3 (Example 7). BHK 570 cells (ATCC CRL-10314) were plated in 10-cm tissue culture dishes and allowed to grow to approximately 50 to 70% confluence overnight at 37°C, 5% CO₂, in DMEM/FBS media (DMEM, Gibco/BRL High Glucose; Life Technologies), 5% fetal bovine serum (Hyclone,

Logan, UT), 1 mM L-glutamine (JRH Biosciences, Lenexa, KS), 1 mM sodium pyruvate (Life Technologies). The cells were then transfected with pZMP6/zveg3 by liposome-mediated transfection (using Lipofectamine™; Life Technologies), in serum free (SF) media (DMEM supplemented with 10 mg/ml transferrin, 5 mg/ml insulin, 2 mg/ml fetuin, 1% L-glutamine and 1% sodium pyruvate). The plasmid was diluted into 15-ml tubes to a total final volume of 640 µl with SF media. 35 µl of the lipid mixture was mixed with 605 µl of SF medium, and the mixture was allowed to incubate approximately 30 minutes at room temperature. Five milliliters of SF media was added to the DNA:lipid mixture. The cells were rinsed once with 5 ml of SF media, aspirated, and the DNA:lipid mixture was added. The cells were incubated at 37°C for five hours, then 6.4 ml of DMEM/10% FBS, 1% PSN media was added to each plate. The plates were incubated at 37°C overnight, and the DNA:lipid mixture was replaced with fresh 5% FBS/DMEM media the next day. On day 5 post-transfection, the cells were split into T-162 flasks in selection medium (DMEM + 5% FBS, 1% L-Gln, 1% NaPyr, 1 µM methotrexate). Approximately 10 days post-transfection, two 150-mm culture dishes of methotrexate-resistant colonies from each transfection were trypsinized, and the cells are pooled and plated into a T-162 flask and transferred to large-scale culture.

20

Example 10

A mammalian cell expression vector for the growth factor domain of zveg3 was constructed essentially as disclosed in Example 7. The coding sequence for the growth factor domain (residues 235-345 of SEQ ID NO:2), joined to a sequence encoding an optimized t-PA secretory signal sequence (U.S. Patent No. 5,641,655) was joined to the linearized pZMP11 vector downstream of the CMV promoter. The plasmid pZMP11 is a mammalian expression vector containing an expression cassette having the CMV immediate early promoter, a consensus intron from the variable region of mouse immunoglobulin heavy chain locus, Kozak sequences, multiple restriction sites for insertion of coding sequences, a stop codon, and a human

30

growth hormone terminator. The plasmid also contains an IRES element from poliovirus, the extracellular domain of CD8 truncated at the C-terminal end of the transmembrane domain, an *E. coli* origin of replication, a mammalian selectable marker expression unit having an SV40 promoter, enhancer and
5 origin of replication, a DHFR gene, the SV40 terminator, and the URA3 and CEN-ARS sequences required for selection and replication in *S. cerevisiae*. The resulting vector, designated pZMP11/zv3GF-otPA, is shown in Fig. 4.

BHK 570 cells were transfected with pZMP11/zv3GF-otPA and cultured essentially as disclosed in Example 9.

10

Example 11

For construction of adenovirus vectors, the protein coding region of human zveg3 was amplified by PCR using primers that added PmeI and Ascl restriction sites at the 5' and 3' termini respectively. PCR primers
15 ZC20,180 (SEQ ID NO:34) and ZC20,181 (SEQ ID NO:35) were used with a full-length zveg3 cDNA template in a PCR reaction as follows: one cycle at 95°C for 5 minutes; followed by 15 cycles at 95°C for 1 min., 61°C for 1 min., and 72°C for 1.5 min.; followed by 72°C for 7 min.; followed by a 4°C soak. The PCR reaction product was loaded onto a 1.2 % low-melting-temperature
20 agarose gel in TAE buffer (0.04 M Tris-acetate, 0.001 M EDTA). The zveg3 PCR product was excised from the gel and purified using a commercially available kit comprising a silica gel membrane spin column (QIAquick™ PCR Purification Kit and gel cleanup kit; Qiagen, Inc.) as per kit instructions. The PCR product was then digested with PmeI and Ascl, phenol/chloroform
25 extracted, EtOH precipitated, and rehydrated in 20 ml TE (Tris/EDTA pH 8). The 1038 bp zveg3 fragment was then ligated into the PmeI-Ascl sites of the transgenic vector pTG12-8 (also known as pHB12-8; see Example 6) and transformed into *E. coli* DH10B™ competent cells by electroporation. Clones containing zveg3 were identified by plasmid DNA miniprep followed by
30 digestion with PmeI and Ascl. A positive clone was sequenced to insure that

there were no deletions or other anomalies in the construct. The sequence of zveg3 cDNA was confirmed.

DNA was prepared using a commercially available kit (Maxi Kit, Qiagen, Inc.), and the 1038bp zveg3 cDNA was released from the pTG12-8 vector using PmeI and AseI enzymes. The cDNA was isolated on a 1% low melting temperature agarose gel and was excised from the gel. The gel slice was melted at 70°C, and the DNA was extracted twice with an equal volume of Tris-buffered phenol and precipitated with EtOH. The DNA was resuspended in 10 µl H₂O.

10 The zveg3 cDNA was cloned into the EcoRV-AseI sites of a modified pAdTrack-CMV (He, T.-C. et al., *Proc. Natl. Acad. Sci. USA* 95:2509-2514, 1998). This construct contains the green fluorescent protein (GFP) marker gene. The CMV promoter driving GFP expression was replaced with the SV40 promoter, and the SV40 polyadenylation signal was replaced with the
15 human growth hormone polyadenylation signal. In addition, the native polylinker was replaced with FseI, EcoRV, and AseI sites. This modified form of pAdTrack-CMV was named pZyTrack. Ligation was performed using a commercially available DNA ligation and screening kit (Fast-Link™ kit; Epicentre Technologies, Madison, WI). Clones containing zveg3 were
20 identified by digestion of mini prep DNA with FseI and AseI. In order to linearize the plasmid, approximately 5 µg of the resulting pZyTrack zveg3 plasmid was digested with PmeI. Approximately 1 µg of the linearized plasmid was cotransformed with 200 ng of supercoiled pAdEasy (He et al., *ibid.*) into *E. coli* BJ5183 cells (He et al., *ibid.*). The co-transformation was done using a
25 Bio-Rad Gene Pulser at 2.5 kV, 200 ohms and 25 µFa. The entire co-transformation mixture was plated on 4 LB plates containing 25 µg/ml kanamycin. The smallest colonies were picked and expanded in LB/kanamycin, and recombinant adenovirus DNA was identified by standard DNA miniprep procedures. Digestion of the recombinant adenovirus DNA with
30 FseI and AseI confirmed the presence of the zveg3 insert. The recombinant adenovirus miniprep DNA was transformed into *E. coli* DH10B™ competent

cells, and DNA was prepared using a Maxi Kit (Qiagen, Inc.) according to kit instructions.

Approximately 5 µg of recombinant adenoviral DNA was digested with PacI enzyme (New England Biolabs) for 3 hours at 37°C in a reaction volume of 100 µl containing 20-30U of PacI. The digested DNA was extracted twice with an equal volume of phenol/chloroform and precipitated with ethanol. The DNA pellet was resuspended in 10µl distilled water. A T25 flask of QBI-293A cells (Quantum Biotechnologies, Inc. Montreal, Qc. Canada), inoculated the day before and grown to 60-70% confluence, were transfected with the PacI digested DNA. The PacI-digested DNA was diluted up to a total volume of 50 µl with sterile HBS (150mM NaCl, 20mM HEPES). In a separate tube, 20 µl of 1mg/ml N-[1-(2,3-Dioleoyloxy)propyl]-N,N,N-trimethyl-ammonium salts (DOTAP) (Boehringer Mannheim, Indianapolis, IN) was diluted to a total volume of 100 µl with HBS. The DNA was added to the DOTAP, mixed gently by pipeting up and down, and left at room temperature for 15 minutes. The media was removed from the 293A cells and washed with 5 ml serum-free minimum essential medium (MEM) alpha containing 1mM sodium pyruvate, 0.1 mM MEM non-essential amino acids, and 25mM HEPES buffer (reagents obtained from Life Technologies, Gaithersburg, MD). 5 ml of serum-free MEM was added to the 293A cells and held at 37°C. The DNA/lipid mixture was added drop-wise to the T25 flask of 293A cells, mixed gently, and incubated at 37°C for 4 hours. After 4 h the media containing the DNA/lipid mixture was aspirated off and replaced with 5 ml complete MEM containing 5% fetal bovine serum. The transfected cells were monitored for GFP expression and formation of foci (viral plaques).

Seven days after transfection of 293A cells with the recombinant adenoviral DNA, the cells expressed the GFP protein and started to form foci (viral "plaques"). The crude viral lysate was collected using a cell scraper to collect all of the 293A cells. The lysate was transferred to a 50-ml conical tube. To release most of the virus particles from the cells, three freeze/thaw cycles were done in a dry ice/ethanol bath and a 37° waterbath.

The crude lysate was amplified (Primary (1°) amplification) to obtain a working "stock" of zveg3 rAdV lysate. Ten 10-cm plates of nearly confluent (80-90%) 293A cells were set up 20 hours previously, 200 µl of crude rAdV lysate added to each 10-cm plate and monitored for 48 to 72 hours looking for CPE under the white light microscope and expression of GFP under the fluorescent microscope. When all of the 293A cells showed CPE (Cytopathic Effect) this 1° stock lysate was collected and freeze/thaw cycles performed as described under Crude rAdV Lysate.

Secondary (2°) amplification of zveg3 rAdV was obtained as follows: Twenty 15-cm tissue culture dishes of 293A cells were prepared so that the cells were 80-90% confluent. All but 20 ml of 5% MEM media was removed, and each dish was inoculated with 300-500 µl of the 1° amplified rAdv lysate. After 48 hours the 293A cells were lysed from virus production, the lysate was collected into 250-ml polypropylene centrifuge bottles, and the rAdV was purified.

NP-40 detergent was added to a final concentration of 0.5% to the bottles of crude lysate in order to lyse all cells. Bottles were placed on a rotating platform for 10 minutes agitating as fast as possible without the bottles falling over. The debris was pelleted by centrifugation at 20,000 X G for 15 minutes. The supernatant was transferred to 250-ml polycarbonate centrifuge bottles, and 0.5 volume of 20% PEG8000/2.5 M NaCl solution was added. The bottles were shaken overnight on ice. The bottles were centrifuged at 20,000 X G for 15 minutes and, the supernatant was discarded into a bleach solution. Using a sterile cell scraper, the white, virus/PEG precipitate from 2 bottles was resuspended in 2.5 ml PBS. The resulting virus solution was placed in 2-ml microcentrifuge tubes and centrifuged at 14,000 X G in the microcentrifuge for 10 minutes to remove any additional cell debris. The supernatant from the 2-ml microcentrifuge tubes was transferred into a 15-ml polypropylene snapcap tube and adjusted to a density of 1.34 g/ml with CsCl. The volume of the virus solution was estimated, and 0.55 g/ml of CsCl was added. The CsCl was dissolved, and 1 ml of this solution weighed 1.34 g. The solution was

transferred to 3.2-ml, polycarbonate, thick-walled centrifuge tubes and spun at 348,000 X G for 3-4 hours at 25°C. The virus formed a white band. Using wide-bore pipette tips, the virus band was collected.

The virus from the gradient had a large amount of CsCl which had to be removed before it was used on cells. Commercially available ion-exchange columns (PD-10 columns prepacked with Sephadex® G-25M; Pharmacia Biotech, Piscataway, NJ) were used to desalt the virus preparation. The column was equilibrated with 20 ml of PBS. The virus was loaded and allowed to run into the column. 5 ml of PBS was added to the column, and fractions of 8-10 drops were collected. The optical densities of 1:50 dilutions of each fraction were determined at 260 nm on a spectrophotometer. A clear absorbance peak was present between fractions 7-12. These fractions were pooled, and the optical density (OD) of a 1:25 dilution was determined. OD was converted to virus concentration using the formula: (OD at 260nm)(25)(1.1×10^{12}) = virions/ml. The OD of a 1:25 dilution of the zvegf3 rAdV was 0.145, giving a virus concentration of 4×10^{12} virions/ml.

To store the virus, glycerol was added to the purified virus to a final concentration of 15%, mixed gently but effectively, and stored in aliquots at -80°C.

A protocol developed by Quantum Biotechnologies, Inc. (Montreal, Canada) was followed to measure recombinant virus infectivity. Briefly, two 96-well tissue culture plates were seeded with 1×10^4 293A cells per well in MEM containing 2% fetal bovine serum for each recombinant virus to be assayed. After 24 hours 10-fold dilutions of each virus from 1×10^{-2} to 1×10^{-14} were made in MEM containing 2% fetal bovine serum. 100 µl of each dilution was placed in each of 20 wells. After 5 days at 37°C, wells were read either positive or negative for Cytopathic Effect (CPE) and a value for "Plaque Forming Units/ml" (PFU) was calculated.

TCID₅₀ formulation used was as per Quantum Biotechnologies, Inc., above. The titer (T) was determined from a plate where virus was diluted

from 10^{-2} to 10^{-14} , and read 5 days after the infection. At each dilution a ratio (R) of positive wells for CPE per the total number of wells was determined.

To calculate titer of the undiluted virus sample: the factor, "F" = $1+d(S-0.5)$; where "S" is the sum of the ratios (R); and "d" is Log10 of the dilution series, for example, "d" is equal to 1 for a ten-fold dilution series. The titer of the undiluted sample is $T = 10^{(1+F)} = \text{TCID}_{50}/\text{ml}$. To convert $\text{TCID}_{50}/\text{ml}$ to pfu/ml, 0.7 is subtracted from the exponent in the calculation for titer (T).

The zveg3 adenovirus had a titer of 1.8×10^{10} pfu/ml.

10 Example 12

Treatment of mice with zveg3-adenovirus led to changes in liver and spleen. The livers were pale and very enlarged, with enlarged vessels at the tips of the lobes. The livers also showed sinusoidal cell proliferation. Changes were also seen in hepatocytes (hypertrophy, degeneration, and necrosis) and were most likely non-specific effects of adenovirus infection. Splenic change consisted of increased extramedullary hematopoiesis, which was correlated with enlarged splenic size.

Example 13

20 Data from adenovirus-treated and transgenic mice were consistent with increased hematopoiesis and/or angiogenesis in the test animals. A study was therefore undertaken to test whether adenovirally delivered zveg3 stimulated cells in the hematopoietic or lymphoid compartments to proliferate, as determined by examination of peripheral blood, histological examination, and incorporation of bromodeoxy uridine (BrdU) into tissues.

Mice (male, C57Bl, 7 weeks old) were divided into three groups. On day 0, parental or zveg3 adenovirus was administered to the first (n=11) and second (n=12) groups, respectively, via the tail vein, with each mouse receiving a dose of $\sim 1 \times 10^{11}$ particles in ~ 0.1 ml volume. The third group (n=8) received no treatment. Each mouse was given two intraperitoneal doses of 3

mg of freshly made BrdU solution at approximately 24 and 12 hours prior to sacrifice. On days 2, 4, 6, 8, and 10, two mice from each treatment group and one or two untreated mice were sacrificed, and tissues and blood were harvested. Samples were analyzed for complete blood count (CBC) and serum chemistry, and slides were prepared for manual differential blood and marrow progenitor cell analysis. One femur, lung, heart, thymus, liver, kidney, spleen, pancreas, duodenum, and mesenteric lymph nodes were submitted for standard histology and assessment of BrdU incorporation. The lining of the duodenum served as the control tissue for BrdU incorporation.

10 In addition, two mice that received approximately half the dose of zvegf3 adenovirus particles and one mouse that received the full dose of parental adenovirus were sacrificed and analyzed as described above on day 16.

15 A piece of liver from each mouse was saved for mRNA assay of adenovirus protein to examine the time course of expression of the adenovirus preparations.

Beginning on day 6, most of the animals treated with either adenovirus had visibly enlarged livers and spleens compared to the untreated mice. The livers of the zvegf3 adenovirus-treated mice tended to look more pale than animals treated with the parental virus. Proliferation of sinusoidal cells was observed in liver. Visual inspection suggested that these cells were stellate cells and/or fibroblasts. Spleen color was the same in both groups. Most of the animals that received the zvegf3 adenovirus had paler femur shafts, with the marrow lighter in color.

25 Peripheral blood CBCs showed a possible difference in platelet counts, but not in RBC or WBC counts between zvegf3 and parental virus-treated animals. In comparison to the untreated and parental virus-treated groups, the zvegf3 group had lower platelet counts on days 2, 4, 6, and 8, but not on day 10. The mean platelet volume (average size of individual platelets) 30 in the zvegf3 group also tended to be greater, consistent with a relative increase in the larger, immature platelet population.

BrdU labeling showed increased cell proliferation in kidney, mainly in the medulla and to a lesser extent in the cortex. Proliferating cells appeared to be interstitial cells, which may have included fibroblasts and/or mesangial cells.

5

Example 14

Zveg3 was assayed in an aortic ring outgrowth assay (Nicosia and Ottinetti, *Laboratory Investigation* 63(1):115, 1990; Villaschi and Nicosia, *Am. J. Pathology* 143(1):181-190, 1993). Thoracic aortas were isolated from
10 1-2 month old SD male rats and transferred to petri dishes containing HANK's buffered salt solution. The aortas were flushed with additional HANK's buffered salt solution to remove blood, and adventitial tissue surrounding the aorta was carefully removed. Cleaned aortas were transferred to petri dishes containing EBM basal media, serum free (Clonetics, San Diego, CA). Aortic
15 rings were obtained by slicing approximately 1-mm sections using a scalpel blade. The ends of the aortas used to hold the aorta in place were not used. The rings were rinsed in fresh EBM basal media and placed individually in a wells of a 24-well plate coated with Matrigel (Becton Dickinson, Bedford, MA). The rings were overlayed with an additional 50 µl Matrigel and placed at 37°C
20 for 30 minutes to allow the matrix to gel. Test samples were diluted in EBM basal serum-free media supplemented with 100 units/ml penicillin, 100 µg/ml streptomycin and HEPES buffer and added at 1 ml/well. Background control was EBM basal serum-free media alone. Basic FGF (R&D Systems, Minneapolis, MN) at 20 ng/ml was used as a positive control. Zveg33
25 pZyTrack adenovirus (Example 11) was added to wells, assuming a cell count of 500,000 cells and a multiplicity of infection of 5000 particles/cell. A null ZyTrack adenovirus (zPar) was used as a control. Samples were added in a minimum of quadruplets. Rings were incubated for 5-7 days at 37°C and analyzed for growth. Aortic outgrowth was scored by multiple, blinded
30 observers using 0 as no growth and 4 as maximum growth. Zveg3 adenovirus produced a significant increase in outgrowth as compared to controls, and was

comparable to other potent growth factors (e.g., bFGF). In additional experiments, purified zveg3 growth factor domain also caused a significant increase in outgrowth at concentrations down to approximately 50 ng/ml.

5 Zveg3-responsive cells were stained for alpha smooth muscle actin (characteristic of SMCs), von Willebrand factor (characteristic of endothelial cells), type I collagen (characteristic of fibroblasts), and vimentin (stains all three cell types). The observed staining patterns indicated that the cells were fibroblasts and smooth muscle cells, with the possible inclusion of pericytes.

10

Example 15

Thirty mice (male, c57BL6) were each injected subcutaneously with 2.5×10^5 Lewis lung carcinoma cells (obtained from American Type Culture Collection, Manassas, VA). Three days after implantation of cells, the
15 mice were split into three groups of ten and were injected with either saline, zveg3 adenovirus (1×10^{11} particles), or control adenovirus (1×10^{11} particles). Growth of tumors was monitored by dimensional measurement on day 14 and by gross tumor weight at the time of sacrifice (day 21). Lungs, liver and tumor were examined by histological methods. Tumor size was significantly lower in
20 the zveg3-treated group compared to the control adenovirus group ($p < 0.007$), but not significantly different from the saline-treated group ($p = 0.6$). The incidence of metastasis was low and did not differ among the groups.

Example 16

25 Myeloproliferative activity of zveg3 is tested in a mouse model of myelosuppression. On day 0, mice (male C57Bl) in the myelosuppressed groups are administered 450cGy of radiation and 1.2 mg of Carboplatin. A second set of mice remain untreated. Between days 0 and 7, purified zveg3, thrombopoietin, or vehicle control are administered subcutaneously. On days –
30 7, 6, 10, 15, and 20 the mice are bled under anesthetic by retro-orbital

puncture. Blood samples are analyzed for CBCs, and slides are made for manual differential and immature cell analysis.

When recovery of blood cell lineages is observed, the animals are sacrificed. Bone marrow is harvested for microscopic analysis and bone marrow progenitor assays. Spleen and liver are harvested for histological examination to determine extramedullary hematopoiesis. Lung, thymus, heart, testis, and kidney are analyzed for histology.

Example 17

10 Polyclonal anti-peptide antibodies were prepared by immunizing two female New Zealand white rabbits with the peptides huzveg3-1 (residues 80-104 of SEQ ID NO:2), huzveg3-2 (residues 299-314 of SEQ ID NO:2), huzveg3-3 (residues 299-326 of SEQ ID NO:2 with an N-terminal cys residue), or huzveg3-4 (residues 195-225 of SEQ ID NO:2 with a C-terminal cys residue). The peptides were synthesized using an Applied Biosystems Model 15 431A peptide synthesizer (Applied Biosystems, Inc., Foster City, CA) according to the manufacturer's instructions. The peptides huzveg3-1, huzveg3-3, and huzveg3-4 were then conjugated to the carrier protein maleimide-activated keyhole limpet hemocyanin (KLH) through cysteine residues (Pierce Chemical Co., Rockford, IL). The peptide huzveg3-2 was conjugated to the carrier protein KLH using glutaraldehyde. The rabbits were each given an initial intraperitoneal (IP) injection of 200 µg of conjugated peptide in Complete Freund's Adjuvant (Pierce Chemical Co.) followed by booster IP injections of 100 µg conjugated peptide in Incomplete Freund's Adjuvant every three weeks. 20 Seven to ten days after the administration of the third booster injection, the animals were bled and the serum was collected. The rabbits were then boosted and bled every three weeks.

The huzveg3 peptide-specific antibodies were affinity purified from the rabbit serum using an CNBr-Sepharose® 4B peptide column 30 (Pharmacia Biotech) that was prepared using 10 mg of the respective peptides per gram CNBr-Sepharose®, followed by dialysis in PBS overnight. Peptide

specific-huzveg3 antibodies were characterized by an ELISA titer check using 1 µg/ml of the appropriate peptide as an antibody target. The huzveg3-1 peptide-specific antibodies have a lower limit of detection (LLD) of 500 pg/ml by ELISA on its appropriate antibody target and recognize full-length recombinant protein (MBP-fusion; see Example 28) by ELISA. The huzveg3-2 peptide-specific antibodies had an LLD of 1 ng/ml by ELISA. The huzveg3-3 peptide-specific antibodies had an LLD of 50 pg/ml by ELISA and recognized recombinant protein by Western Blot analysis. The huzveg3-4 peptide-specific antibodies had an LLD of 50 pg/ml by ELISA and recognized recombinant protein by Western Blot analysis.

Example 18

Recombinant zveg3 was analyzed by Western blotting using antibodies to the huzveg3-1 and huzveg3-3 peptides. Protein was produced in BHK cells as disclosed above, and in 293 and MVEC (microvascular endothelial) cells transfected with adenovirus vectors according to conventional methods. Samples were electrophoresed and transferred to nitrocellulose (0.2 µm; Bio-Rad Laboratories, Hercules, CA) at room temperature using a Hoeffer Scientific Instruments (San Francisco, CA) model TE22 blotter with stirring according to directions provided in the instrument manual. The transfer was run at 500 mA for one hour or 50 mA for 12 hours in a buffer containing 25 mM Tris base, 200 mM glycine, and 20% methanol. The filters were then blocked with 10% non-fat dry milk in buffer A (50 mM Tris (pH 7.4), 5 mM EDTA (pH 8.0), 0.05% Igepal CA-630, 150 mM NaCl, 0.25% gelatin) for 10 minutes at room temperature. The nitrocellulose was quickly rinsed, then primary antibody was added in buffer A containing 2.5% non-fat dry milk. The blots were incubated for 1 hour at room temperature or overnight at 4°C with gentle shaking or rocking. Following the incubation, blots were washed three times for 10 minutes each in buffer A. Secondary antibody (goat anti-rabbit IgG conjugated to horseradish peroxidase; obtained from Rockland Inc., Gilbertsville, PA) diluted 1:4000 in buffer A containing 2.5% non-fat dry milk

was added, and the blots were incubated for one hour at room temperature with gentle shaking or rocking. The blots were then washed three times, 10 minutes each, in buffer A, then quickly rinsed in H₂O. The blots were developed using commercially available chemiluminescent substrate reagents
5 (SuperSignal® ULTRA reagents 1 and 2 mixed 1:1; reagents obtained from Pierce Chemical Co.), and exposed to film (Hyperfilm ECL™; Amersham Pharmacia Biotech, Piscataway, NJ) for times ranging from 1 second to 5 minutes or as necessary.

Using the 3-1 antibody, full-length zveg3 produced in BHK cells
10 showed two bands ($M_r \approx 46$ kDa and 30 kDa) under reducing conditions. The larger band was consistent with the size of the full-length zveg3 monomer, and the smaller band with the CUB domain + interdomain region. Under non-reducing conditions, there was a major band at $M_r \approx 78$ kDa, which appeared to be the dimerized, full-length molecule. Two smaller, minor bands were also
15 observed. Similar results were obtained with the 3-3 antibody, but under reducing conditions the smaller band ran at $M_r \approx 22$ kDa. Sequence analysis of the smaller band showed it to be the isolated growth factor domain.

The recombinant growth factor domain (produced in BHK cells), analyzed using the 3-3 antibody, ran as a broad band of $M_r \approx 18$ kDa under
20 reducing conditions. Under non-reducing conditions, the protein ran as two bands of $M_r \approx 16$ and 28 kDa, indicating the presence of both dimeric and monomeric forms of the growth factor domain.

Zveg3 protein produced in 293 cells grown in serum-free media showed sizes consistent with the predicted full-length protein. Under reducing
25 conditions, the protein ran at $M_r \approx 47$ kDa, and under non-reducing conditions at $M_r \approx 78$ kDa when blots were probed with either antibody 3-1 or 3-3. Addition of serum to the media resulted in cleavage of the protein, as seen in the BHK-produced material.

MVEC cells grown in the presence of 1% serum produced zveg3
30 protein that ran at $M_r \approx 23$ kDa under reducing conditions, and $M_r \approx 28$ kDa under non-reducing conditions using the 3-3 antibody. These results indicate

cleavage of the protein, with the antibody recognizing the monomeric and dimeric forms of the growth factor domain. When the MVEC cells were adapted to serum-free media, full-length protein was observed.

5 Example 19

Recombinant zveg3 growth factor domain was produced in BHK 570 cells grown in cell factories. Three 15-liter cultures were harvested, and the media were sterile filtered using a 0.2 μ filter. Expression levels were estimated by western blot analysis of media samples concentrated to 20X vs. 10 5K cut-off and serially diluted by two-fold to 1.25X. Signal intensity was compared to a signal on the same blot from an MBP-zveg3 fusion protein standard (see Example 28) for which the protein concentration had been determined by amino acid analysis. Expression levels were consistently between 0.25 and 0.35 mg/L of media.

15 Protein was purified from conditioned media by a combination of cation exchange chromatography and hydrophobic interaction chromatography. Culture medium was diluted with 0.1 M acetic acid, pH 3.0, containing 0.3 M NaCl at a ratio of 60%:40%, (medium:acetic acid) to deliver a process stream at 14 mS conductivity and pH 4.0. This stream was delivered 20 to a strong cation exchange resin (Poros® HS; PerSeptive Biosystems, Framingham, MA) with a bed volume of 50 ml in a 2-cm diameter column at a flow rate of 20 ml/minute. A 50-ml bed was sufficient to process 45 L of media and capture all of the target protein. Bound protein was eluted, following column washing for 10 column volumes in 10 mM acetic acid with 0.15 M NaCl 25 at pH =4.0, by forming a linear gradient to 2M NaCl in 10 mM acetic acid, pH 4.0. Ten-ml fractions were captured into tubes containing 2 ml 2.0 M Tris, pH 8.0 to neutralize the acidity. Samples from the cation exchange column were analyzed by SDS PAGE with silver staining and western blotting for the presense of zveg3. The vegf3 growth domain eluted at 0.2 - 0.5 M NaCl. 30 Protein-containing fractions were pooled. A 25-ml bed of chromatography medium (Toso Haas Ether chromatography medium) in a 2 cm diameter

column was equilibrated in 1.8 M $(\text{NH}_4)_2\text{SO}_4$ in 25 mM Na phosphate buffer at pH 7.4. The pooled protein from the cation exchange step was adjusted to 1.8 M $(\text{NH}_4)_2\text{SO}_4$ in 25 mM Na phosphate, pH 7.0. This stream was flowed over the column at 10 ml/minute. Once the loading was completed the column was washed for 10 column volumes with the equilibration buffer prior to eluting with a 10 column volume gradient formed between the equilibration buffer and 40 mM boric acid at pH 8.8. The zveg3 growth factor domain protein eluted fairly early in the gradient between 1.5 and 1.0 M $(\text{NH}_4)_2\text{SO}_4$. At this point the protein was 40 - 60% pure with a major contaminant being insulin-like growth factor binding protein 4 (IGFBP4).

Protein from the HIC (Ether) chromatography step was applied to a C4 reverse-phase HPLC column. The zveg3 growth factor domain protein eluted at 36% acetonitrile. This material still contained approximately 20% (mole/mole) IGFB4.

15

Example 20

Recombinant zveg3 growth factor domain is purified from cell-conditioned media by a combination of cation exchange chromatography, hydrophobic interaction chromatography, and nickel affinity chromatography. Protein is captured on a strong cation exchange medium and eluted essentially as disclosed in Example 19. The eluted protein is further purified by hydrophobic interaction chromatography on an ether resin (Poros® ET; PerSeptive Biosystems). The partially purified zveg3 protein is then bound to a nickel chelate resin at pH 7.0 - 8.0 in 25 mM Na phosphate buffer containing 0.25 M NaCl. The bound protein is eluted with a descending pH gradient down to pH 5.0 or an imidazole gradient. The eluate from the nickel column is adjusted to 1 M $(\text{NH}_4)_2\text{SO}_4$, 20 mM MES (morpholino ethanesulfonic acid) at pH 6.0 and passed through a phenyl ether hydrophobic interaction chromatography column (Poros® PE, PerSeptive Biosystems) that has been equilibrated in 1 M $(\text{NH}_4)_2\text{SO}_4$, 20 mM MES, pH 6.0. IGFBP4 and minor contaminants are retained on the column. The pass-through fraction, which

contains highly purified zveg3, is collected. The collected protein is desalted according to conventional methods (e.g., dialysis, ion-exchange chromatography).

5 Example 21

Recombinant zveg3 was analyzed for mitogenic activity on human aortic smooth muscle cells (HAoSMC; Clonetics Corp., Walkersville, MD) and human umbilical vein endothelial cells (HUVEC; Clonetics Corp.). HAoSMC and HUVEC were plated at a density of 5,000 cells/well in 96-well
10 culture plates and grown for approximately 24 hours in DMEM containing 10% fetal calf serum at 37°C. Cells were quiesced by incubating them for 24 hours in serum-free DMEM/Ham's F-12 medium containing insulin (5 µg/ml), transferrin (20 µg/ml), and selenium (16 pg/ml) (ITS). At the time of the assay, the medium was removed, and test samples were added to the wells in
15 triplicate. Test samples consisted of either conditioned media (CM) from adenovirally-infected HaCaT human keratinocyte cells (Boukamp et al., *J. Cell. Biol.* 106:761-771, 1988) expressing full-length zveg3, purified growth factor domain expressed in BHK cells, or control media from cells infected with parental adenovirus (Zpar). The CM was concentrated 10-fold using a 15 ml
20 centrifugal filter device with a 10K membrane filter (Ultrafree®; Millipore Corp., Bedford, MA), then diluted back to 3x with ITS medium and added to the cells. The control CM was generated from HaCaT cells infected with a parental green fluorescent protein-expressing adenovirus and treated identically to the zveg3 CM. Purified protein in a buffer containing 0.1% BSA was serially diluted into
25 ITS medium at concentrations of 1 µg/ml to 1 ng/ml and added to the test plate. A control buffer of 0.1% BSA was diluted identically to the highest concentration of zveg3 protein and added to the plate. For measurement of [³H]thymidine incorporation, 20 µl of a 50 µCi/ml stock in DMEM was added directly to the cells, for a final activity of 1 µCi/well. After another 24 hour
30 incubation, mitogenic activity was assessed by measuring the uptake of [³H]thymidine. Media were removed and cells were incubated with 0.1 ml of

trypsin until cells detached. Cells were harvested onto 96-well filter plates using a sample harvester (FilterMate™ harvester; Packard Instrument Co., Meriden, CT). The plates were then dried at 65°C for 15 minutes, sealed after adding 40 µl/well scintillation cocktail (Microscint™ O; Packard Instrument Co.) and counted on a microplate scintillation counter (Topcount®; Packard Instrument Co.).

Results presented in Table 8 demonstrate that zveg3 CM had approximately 1.5-fold higher mitogenic activity on HAoSM cells over control CM, and purified protein caused a maximal 1.8-fold increase in [³H]thymidine incorporation over the buffer control.

Table 8

<u>Sample</u>	<u>CPM Incorporated</u>	
	<u>Mean</u>	<u>St. dev.</u>
zveg3 (3x CM)	81089	8866
Zpar (3x CM)	58760	2558
zveg3 GF domain, 1 µg/ml	63884	3281
zveg3 GF domain, 500 ng/ml	57484	9744
zveg3 GF domain, 100 ng/ml	70844	10844
zveg3 GF domain, 50 ng/ml	61164	2813
zveg3 GF domain, 10 ng/ml	60676	1514
zveg3 GF domain, 5 ng/ml	60197	2481
zveg3 GF domain, 1 ng/ml	49205	5208
Buffer control	39645	9793
PDGF 10 ng/ml (maximal response)	50634	4238
Media alone (basal response)	24220	2463

Results presented in Table 9 demonstrate that zveg3 CM had no mitogenic activity on HUVEC compared to the control CM, and purified protein caused a maximal 1.3-fold increase in [³H]thymidine incorporation over the buffer control.

Table 9

<u>Sample</u>	CPM Incorporated	
	<u>Mean</u>	<u>St. dev.</u>
zveg3 (3x CM)	62723	10716
Zpar (3x CM)	61378	1553
zveg3 VEGF domain, 1 μ g/ml	44901	6592
zveg3 VEGF domain, 500 ng/ml	41921	5330
zveg3 VEGF domain, 100 ng/ml	35613	5187
zveg3 VEGF domain, 50 ng/ml	31107	525
zveg3 VEGF domain, 10 ng/ml	28505	2950
zveg3 VEGF domain, 5 ng/ml	29290	988
zveg3 VEGF domain, 1 ng/ml	28586	2718
Buffer control	33461	404
VEGF 50 ng/ml (maximal response)	53225	5229
Media alone (basal response)	22264	2814

Example 22

Recombinant zveg3 protein was assayed for stimulation of intracellular calcium release as an indicator of receptor binding and activation. Cells were cultured in chambered borosilicate coverglass slides. On the day of assay, cells were incubated for 30 minutes at room temperature in KRW buffer (KrebsRingerWollheim; 140 mM NaCl, 3.6 mM KCl, 0.5 mM NaH₂PO₄, 0.5 mM MgSO₄, 2 mM NaHCO₃, 3 mM glucose, 1.5 mM CaCl₂, 10 mM HEPES pH 7.4) containing 2 μ M fura-2 AM (obtained from Molecular Probes Inc., Eugene, OR), washed twice with KRW buffer, and allowed to sit at room temperature for at least 15 minutes before addition of growth factor or cell-conditioned culture medium (CM) to be tested. Changes in cytosolic calcium were measured by fluorescence ratio imaging (excitation at 340 nm divided by excitation at 380 nm). Digital imaging was carried out using an inverted fluorescent microscope (Nikon TE300) equipped with an oil objective (Nikon 40x Plan Fluor). Images

were acquired using a Princeton CCD digital camera and analyzed with Universal Imaging Metafluor software. Data are presented in Table 10.

Table 10

<u>Cell Line</u>	<u>Zveg3 CM</u>	<u>Control CM</u>	<u>VEGF</u>	<u>PDGF BB</u>
aortic ring cells	+	-	-	+
pericytes	+	-	-	+
aortic smooth muscle				
cells	+	-	-	+
aortic adventitial				
fibroblasts	+	-	-	+

5

Example 23

Northern blot analysis was performed using total RNA from the human neuronal and glial cell lines A172 (glioblastoma), NTera 2 (teratocarcinoma neuronal precursor; obtained from Strategene Cloning Systems, La Jolla, CA), U-87 MG (glioblastoma/astrocytoma), U-118 MG (glioblastoma), U138 MG (glioblastoma), U373 MG (glioblastoma). Except as noted, cell lines were obtained from American Type Culture Collection, Manassas, VA. Blots were prepared using 10 µg of RNA per lane. An approximately 400 bp DNA probe was generated by digestion of human zveg3 cDNA with EcoRI and BglII. The DNA probe was gel electrophoresis followed by extraction using a spin column containing a silica gel membrane (QIAquick™ Gel Extraction Kit; Qiagen, Inc., Valencia, CA). The probe was radioactively labeled with ³²P using a commercially available kit (Rediprime™ II random-prime labeling system; Amersham Corp., Arlington Heights, IL) according to the manufacturer's specifications. The probe was purified using a push column. Hybridization took place overnight at 65°C in a commercially available solution (ExpressHyb™ Hybridization Solution; Clontech Laboratories, Inc.).

The blots were then washed 4X in 2X SCC and 0.05% SDS at room temperature, followed by two washes in 0.1X SSC and 0.1% SDS at 50°C. One transcript size was detected at approximately 4 kb in A172, U-87 MG, U-118 MG, U138 MG, and U373 MG samples. Signal intensity was highest for U373 MG, U-118MG, and U-87 MG.

Example 24

10 µg of recombinant zveg3 growth factor domain protein was combined with 438 µl PBS containing 1 mCi Na-¹²⁵I (Amersham Corp.). One derivatized, nonporous polystyrene bead (ODO-Beads®; Pierce Chemical Co., Rockford, IL) was added, and the reaction mixture was incubated one minute on ice. The iodinated protein was separated from unincorporated ¹²⁵I by gel filtration, elution buffer PBS, 0.25% gelatin. The active fraction contained 4.9 µg/mL ¹²⁵I-zveg3 with a specific activity of 4.3 X 10⁴ cpm/ng.

The following cell lines were plated into the wells of a 24-well tissue culture dish and cultured in growth medium for three days:

1. Rat aortic ring pool (ARC)
2. Rat aortic ring clone 14B (ARC#14B)
3. Human umbilical vein endothelial cells, passage 5 (HUVEC)
4. Human aortic adventitial fibroblasts, passage 4 (AOAF)
5. Human aortic smooth muscle cells, passage 9 (AOSMC)
6. Human retinal pericytes, passage 4 (pericytes)

Cells were washed once with ice cold binding buffer (RPMI containing 0.1% BSA, 20 mM Tris:HCl, pH 7.2) and then 250 µl of the following solutions were added to each of three wells of the culture dishes containing the test cells.

Binding solutions were prepared in 5 mL of binding buffer with 10 ng/mL ¹²⁵I-zveg3 and:

1. No addition.
2. One µg/mL zveg3.
3. One µg/mL VEGF (R&D Systems, Minneapolis, MN).

4. One $\mu\text{g/mL}$ PDGF-BB.
5. Five $\mu\text{g/mL}$ PDGF receptor α (R&D Systems).
6. Five $\mu\text{g/mL}$ PDGF receptor β (as IgG Fc-receptor extracellular domain fusion).

5 The reaction mixtures were incubated on ice for 2 hours, then washed three times with one mL of ice-cold binding buffer. The bound ^{125}I -zveg3 was quantitated by gamma counting a NaOH extract of the cells.

Results, shown in Fig. 5, indicate binding of zveg3 to the PDGF receptor α . The data are graphed as ^{125}I -zveg3 bound/well. The error bars
10 represent standard deviations.

The experiment was repeated with the addition of rat liver stellate cells, passage 6 (Greenwel et al., *Laboratory Investigation* 69:210-216, 1993). Stellate cells bound zveg3 at a level comparable to pericytes.

15 Example 25

Binding of recombinant zveg3 to PDGF alpha and beta receptors was measured by mass spectrometry using a surface-enhanced laser desorption and ionization (SELDI) instrument (ProteinChip™, Ciphergen Biosystems, Palo Alto, CA). For this experiment an 8-spot, preactivated
20 surface array was used. To this amine-activated chip, protein-A (Zymed Laboratories, Inc., San Francisco, CA) was added at a concentration of 1 mg/ml, and the chip was incubated at 4°C for four hours. After blocking with 1M ethanolamine pH 8.0 and subsequent washes (once in 0.1% Triton X-100 in PBS; once in 100mM Na Acetate, pH4.5, 0.5 M NaCl; once in 100mM Tris-
25 HCl, pH8.5, 0.5 M NaCl; once in PBS), IgG Fc-receptor extracellular domain fusion proteins (PDGF alpha receptor, PDGF beta receptor, or unrelated control receptor) were added, and the chip was incubated at 4°C overnight. After three washes in PBS, 250 μl of zveg3 (300 ng/ml), PDGF-AA, or PDGF-BB was added, and the chip was incubated overnight at 4°C. The chip was
30 washed twice with 0.05% Triton X100, 100 mM HEPES pH 7.2, then twice with deionized water. The chip was allowed to dry at room temperature before two

additions of 0.3 microliters of sinapinic acid (Ciphergen Biosystems) in a 50:50 mixture of acetonitrile and 1% trifluoroacetic acid. Ligands that bound receptor were retained on the chip after washing and subsequently detected by mass spectrometry. Assignment of a + or - for binding was made by comparing the PDGF receptor mass spectrometry profile to that of an Fc only control for each ligand. Data are shown in Table 11.

Table 11

	<u>PDGF AA</u>	<u>PDGF AB</u>	<u>PDGF BB</u>	<u>zVEGF3</u>
PDGFR-alpha/Fc	+	+	+	+
PDGFR-beta/Fc	+/-	+/-	+	-

10

Example 26

Northern blot analysis was performed with poly(A) RNA from human vascular cell lines HUVEC (human umbilical vein endothelial cells; Cascade Biologics, Inc., Portland, OR), HPAEC (human pulmonary artery endothelial cells; Cascade Biologics), HAEC (human aortic endothelial cells; Cascade Biologics), AoSMC (aortic smooth muscle cells; Clonetics Corporation, Walkersville, MD), UASMC (umbilical artery smooth muscle cells; Clonetics Corp.), HISM (human intestinal smooth muscle; American Type Culture Collection, CRL 7130), SK5 (human dermal fibroblast cells; obtained from Dr. Russel Ross, University of Washington), NHLF (normal human lung fibroblasts; Clonetics Corp.), NHDF-neo (normal human dermal fibroblast-neonatal; Clonetics Corp.); and from leukemia cell lines Daudi, Raji, Molt-4, K562 (all obtained from Clontech Laboratories, Inc.), HL60, Jurkat, and Hut 78. RNA was loaded at 2µg per lane. An approximately 490bp DNA probe was generated by digestion of a full-length zVEGF3 clone with PvuI and StuI. The DNA probe was electrophoresed and purified using a spin column containing a silica gel membrane (QIAquick™ Gel Extraction Kit; Qiagen, Inc., Valencia, CA). The probe was radioactively labeled with ³²P using a commercially

25

available kit (Rediprime™ II random-prime labeling system; Amersham Corp., Arlington Heights, IL) according to the manufacturer's specifications. The probe was purified using a push column. Expresshyb (Clontech, Palo Alto, Ca) solution was used for the hybridizing solution for the blots. Hybridization took
5 place overnight at 65°C in a commercially available hybridization solution (ExpressHyb™ Hybridization Solution; Clontech Laboratories, Inc., Palo Alto, CA). The blots were then washed four times in 2X SCC and 0.1% SDS at room temperature, followed by two washes in 0.1X SSC and 0.1% SDS at 50°C. One transcript size was detected at approximately 4 kb in NHLF,
10 NHDF-neo, SK5, UASMC, HAEC, AoSMC, Jurkat, Hut78. Signal intensity was highest for UASMC, SK5, NHLF, and NHDF-neo.

Example 27

The effects of zveg3 on vascular endothelial regeneration and
15 intimal hyperplasia are tested in a balloon injured rat carotid artery model. Adenovirus is used as a delivery vector.

To determine the infectivity of adenovirus and the level of gene expression obtained in the vascular wall, rats are balloon injured as disclosed below and infused with an adenovirus vector comprising an expression unit for
20 green fluorescent protein (GFP). Three groups of three rats each are infused with doses of 1.5×10^{10} pfu/ml, 3×10^{10} pfu/ml, and 6×10^{10} pfu/ml. Injured and uninjured carotids are harvested 48 hours after infection and fixed in 10% buffered formalin for 24 hrs. The tissue is processed and analyzed using an anti-GFP antibody to determine % infectivity.

25 The effects of zveg3 are determined in a 14-day study using the optimal dose determined from the GFP study. Two groups of 14 animals each are balloon-injured and infected with either zveg3 adenovirus or control adenovirus. The left common carotid is isolated, and the flow of blood through the vessel is stopped by tying off the internal carotid, the external carotid, and
30 proximally, the common carotid. An arteriotomy is made between the tie on the external carotid and the bifurcation, and the vessel is rinsed out with lactated

Ringer's. A 2F-embolectomy catheter is inserted, inflated, and removed, while twisting, to remove endothelial cells; this procedure is done three times. The vessel is then rinsed again, and approximately 50 μ l of adenovirus solution is injected into it using a catheter of silastic tubing. The catheter is tied into the vessel just distal to the bifurcation and left in place for approximately 20 minutes. The catheter is then removed, and the vessel is flushed briefly with blood by loosening the proximal tie. A tie is made just distal to the bifurcation. Blood flow is restored by removing the tie on the internal carotid and the proximal tie on the common carotid. The ties on the external carotid remain.

To determine zveg3 protein production in the vessel wall, two animals from each group are sacrificed on days one and seven. Tissues are processed for immunohistochemical analysis and Western blotting analysis. For immunohistochemical analysis, tissues are kept in formalin for 24 hours, then transferred to 70% ethyl alcohol. For Western blotting analysis, tissues are flash frozen and stored at -80°C . At thirteen days animals are given BrdU tablets subcutaneously. At fourteen days (24 hours after BrdU insertion) Evan's Blue dye is given intravenously to stain for non-endothelialized segments, and the animals are bled and sacrificed. Animals are then exsanguinated and perfusion-fixed with 10% buffered formalin. Both carotids, liver, kidney and spleen are harvested. The carotids are visually inspected, and re-endothelialization is quantitated by measuring the distance from the bifurcation to the distal dye (white/blue) boundary. All tissue are kept in formalin for 24 hours then transferred to 70% ethyl alcohol. The carotids are sectioned into three pieces each and embedded in paraffin blocks. The liver, kidney, and spleen are visually inspected and processed. Slides are made of cross-sections of the carotids and stained with Hematoxylin and Eosin, then measured using the SPOT[®] diagnostic program (Diagnostic Instruments, Inc., Sterling Heights, MI). Measurements include the length of the internal elastic lamina and the areas of the media, intima, and lumen. ICC analysis includes the number of infected cells (using anti-gfp antibodies), cell proliferation (BrdU labeling), and % cell death.

A third study is conducted to determine the time course of zveg3 gene expression following balloon injury. Carotids are harvested from balloon-injured animals (5 animals/time point) at T=0 (non-injured), T=6 hrs, 1, 4, 7, and 14 days. The carotids are flash-frozen and stored at -80°C for Northern blot analysis.

Example 28

An expression plasmid containing a polynucleotide encoding human zveg3 fused N-terminally to maltose binding protein (MBP) was constructed via homologous recombination.

A fragment of human zveg3 cDNA was isolated using PCR. Two primers were used in the production of the human zveg3 fragment in a PCR reaction. Primer ZC20,572 (SEQ ID NO:44), contained 40 bp of vector flanking sequence and 25 bp corresponding to the amino terminus of human zveg3, and primer ZC20,573 (SEQ ID NO:45) contained 40 bp of the 3' end corresponding to flanking vector sequence and 25 bp corresponding to the carboxyl terminus of human zveg3. The PCR reaction conditions were 25 cycles of 94°C for 30 seconds, 50°C for 30 seconds, and 72°C for 1.5 minutes; followed by 4°C soak, run in duplicate. A 2-μl aliquot of the 100-μl reaction mixture was run on a 1.0% agarose gel with Tris/borate/EDTA buffer for analysis, and the expected band of approximately 1000 bp was seen. The remaining 90 μl of the reaction mixture was combined with the second PCR tube precipitated with 400 μl of absolute ethanol to be used for recombining into the Sma1-cut recipient vector pTAP98 to produce the construct encoding the MBP-zveg3 fusion.

Plasmid pTAP98 was derived from the plasmids pRS316 (a *Saccharomyces cerevisiae* shuttle vector; see, Hieter and Sikorski, *Genetics* 122:19-27, 1989) and pMAL™-c2X (New England Biolabs; Beverly, MA). The latter vector carries the *tac* promoter driving *MalE* (gene encoding MBP) followed by a His tag, a thrombin cleavage site, a cloning site, and the *mb* terminator. The vector pTAP98 was constructed using yeast homologous

recombination. 100ng of EcoRI-cut pMALTM-c2X was recombined with 1µg PvuI-cut pRS316, 1µg linker, and 1µg ScaI/EcoRI-cut pRS316. The linker was constructed by combining oligonucleotides ZC19,372 (SEQ ID NO:46) (100pmole), ZC19,351 (SEQ ID NO:47) (1pmole), ZC19,352 (SEQ ID NO:48) (1pmole), and ZC19,371 (SEQ ID NO:49) (100pmole) in a PCR reaction for 10 cycles of 94°C for 30 seconds, 50°C for 30 seconds, and 72°C for 30 seconds; followed by 4°C soak. PCR products were concentrated via 100% ethanol precipitation.

A vector containing the MBP-zveg3 fusion sequence was constructed by homologous recombination. One hundred microliters of competent yeast cells (*S. cerevisiae*) were combined with 10 µl of a mixture containing approximately 1 µg of the human zveg3 insert and 100 ng of SmaI digested pTAP98 vector, and transferred to a 0.2-cm electroporation cuvette. The yeast/DNA mixture was electropulsed at 0.75 kV (5 kV/cm), infinite ohms, 25 µF. To each cuvette was added 600 µl of 1.2 M sorbitol. The yeast was then plated in two 300-µl aliquots onto two -URA D plates and incubated at 30°C. After about 48 hours, the Ura⁺ yeast transformants from a single plate were resuspended in 1 ml H₂O and spun briefly to pellet the yeast cells. The cell pellet was resuspended in 1 ml of lysis buffer (2% Triton X-100, 1% SDS, 100 mM NaCl, 10 mM Tris, pH 8.0, 1 mM EDTA). Five hundred microliters of the lysis mixture was added to an Eppendorf tube containing 300 µl acid-washed glass beads and 200 µl phenol-chloroform, vortexed for 1 minute intervals two or three times, followed by a 5 minute spin in a microcentrifuge at maximum speed. Three hundred microliters of the aqueous phase was transferred to a fresh tube, and the DNA was precipitated with 600 µl ethanol (EtOH), then centrifuged for 10 minutes at 4°C. The DNA pellet was resuspended in 100 µl H₂O. Electrocompetent *E. coli* cells (MC1061; Casadaban et. al., *J. Mol. Biol.* 138: 179-207) were transformed with 1 µl of the yeast DNA prep in a volume of 40 µl. The cells were electropulsed at 2.0 kV, 25 µF, 400 ohms. Following electroporation, 0.6 ml SOC (2% BactoTM Tryptone (Difco Laboratories, Detroit, MI), 0.5% yeast extract (Difco

Laboratories), 10 mM NaCl, 2.5 mM KCl, 10 mM MgCl₂, 10 mM MgSO₄, 20 mM glucose) was plated in one aliquot on LB AMP plates (LB broth (Lennox), 1.8% agar (Bacto™; Difco Laboratories), 100 mg/L ampicillin). Individual clones harboring the correct expression construct for zveg3 were identified by expression. Cells were grown in minimal medium supplemented with casamino acids and 100 µg/ml of ampicillin overnight. 50 µl of the overnight culture was used to inoculate 2 ml of fresh medium. Cultures were grown at 37°C, shaking for 2 hours. One ml of the culture was induced with 1mM IPTG. 2-4 hours later 250 µl of each culture was mixed with 250 µl acid-washed glass beads and 250 µl Thorner buffer (8 M urea, 100 mM Tris pH 7.0, 10% glycerol, 2 mM EDTA, 5% SDS) supplemented with 5% β-ME and dye. Samples were vortexed for one minute and heated to 65°C for 5-10 minutes. 20 µl was loaded per lane on a 4%-12% PAGE gel (NOVEX, San Diego, CA). Gels were run in 1X MES buffer. The positive clones were designated pCZR236 and subjected to sequence analysis. The polynucleotide sequence of the MBP-zveg3 fusion in pCZR236 is shown in SEQ ID NO:50.

To express the fusion protein, 1 µl of sequencing DNA was used to transform *E. coli* strain W3110 (obtained from American Type Culture Collection, Manassas, VA). The cells were electropulsed at 2.0 kV, 25 µF and 400 ohms. Following electroporation, 0.6 ml SOC (2% Bacto™ Tryptone (Difco Laboratories), 0.5% yeast extract (Difco Laboratories), 10 mM NaCl, 2.5 mM KCl, 10 mM MgCl₂, 10 mM MgSO₄, 20 mM glucose) was plated in one aliquot on LB AMP plates (LB broth (Lennox), 1.8% Bacto™ Agar (Difco Laboratories), 100 mg/L ampicillin). Cells were picked from the plate and grown in minimal medium containing casamino acids overnight. A 50-µl aliquot of the overnight culture was used to inoculate 2 ml of fresh medium. Cultures were grown at 37°C with shaking for 2 hours. One ml of the culture was induced with 1mM IPTG, and the cells were lysed essentially as described above. Twenty-µl aliquots of the lysate were analyzed by gel electrophoresis as described above.

Example 29

Recombinant zveg3 was analyzed for mitogenic activity on rat stellate cells (obtained from N. Fausto, University Of Washington). Stellate cells were plated at a density of 2,000 cells/well in 96-well culture plates and
5 grown for approximately 72 hours in DMEM containing 10% fetal calf serum at 37°C. Cells were quiesced by incubating them for 20 hours in serum-free DMEM/Ham's F-12 medium containing insulin (5 µg/ml), transferrin (20 µg/ml), and selenium (16 pg/ml) (ITS). At the time of the assay, the medium was removed, and test samples were added to the wells in triplicate. Test samples
10 consisted of either conditioned media (CM) from adenovirally-infected HaCaT human keratinocyte cells (Boukamp et al., *J. Cell. Biol.* 106:761-771, 1988) expressing full-length zveg3, purified growth factor domain expressed in BHK cells, or control media from cells infected with parental adenovirus (Zpar) containing an expression unit for green fluorescent protein. The CM was
15 concentrated 10-fold using a 15-ml centrifugal filter device with a 10K membrane filter (Ultrafree®; Millipore Corp., Bedford, MA), then diluted back to 3x with ITS medium and added to the cells. Purified protein in a buffer containing 0.1% BSA was serially diluted into ITS medium at concentrations of 1 µg/ml to 1 ng/ml and added to the test plate. A control buffer of 0.1% BSA
20 was diluted identically to the highest concentration of zveg3 protein and added to the plate. For measurement of [³H]thymidine incorporation, 20 µl of a 50 µCi/ml stock in DMEM was added directly to the cells, for a final activity of 1 µCi/well. After another 24-hour incubation, mitogenic activity was assessed by measuring the uptake of [³H]thymidine. Media were removed, and cells were
25 incubated with 0.1 ml of trypsin until cells detached. Cells were harvested onto 96-well filter plates using a sample harvester (FilterMate™ harvester; Packard Instrument Co., Meriden, CT). The plates were then dried at 65°C for 15 minutes, sealed after adding 40 µl/well scintillation cocktail (Microscint™ O; Packard Instrument Co.) and counted on a microplate scintillation counter
30 (Topcount®; Packard Instrument Co.).

123

Results, presented in Table 12, demonstrated that zveg3 CM had approximately 4.4-fold higher mitogenic activity on stellate cells over control CM, and purified protein at 100 ng/ml caused a maximal 14-fold increase in [³H]thymidine incorporation over the buffer control.

5

Table 12

<u>Sample</u>	<u>CPM Incorporated</u>	
	<u>Mean</u>	<u>St. dev.</u>
zveg3 (2x CM)	42489	1306
Zpar (2x CM)	9629	540
zveg3 GF domain, 100 ng/ml	77540	4142
zveg3 GF domain, 33.3 ng/ml	74466	18142
zveg3 GF domain, 11.1 ng/ml	52462	6239
zveg3 GF domain, 3.7 ng/ml	15128	4989
Buffer control	5618	573
PDGF-BB 20 ng/ml	19741	2075
PDGF-AA 20 ng/ml	33133	3325
Media alone (basal response)	6765	226

From the foregoing, it will be appreciated that, although specific embodiments of the invention have been described herein for purposes of illustration, various modifications may be made without deviating from the spirit and scope of the invention. Accordingly, the invention is not limited except as by the appended claims.

10

CLAIMS

We claim:

1. An isolated polypeptide of at least 15 amino acid residues comprising an epitope-bearing portion of a protein of SEQ ID NO:2 or SEQ ID NO:43.
2. The polypeptide of claim 1 wherein said polypeptide comprises a segment that is at least 90% identical to a sequence selected from the group consisting of:
 - (a) residues 46-163 of SEQ ID NO:2; and
 - (b) residues 235-345 of SEQ ID NO:2.
3. The isolated polypeptide according to claim 1 wherein said polypeptide is selected from the group consisting of:
 - residues 15-163 of SEQ ID NO:2;
 - residues 46-163 of SEQ ID NO:2;
 - residues 15-170 of SEQ ID NO:2;
 - residues 46-170 of SEQ ID NO:2;
 - residues 15-234 of SEQ ID NO:2;
 - residues 46-234 of SEQ ID NO:2;
 - residues 15-229 amide of SEQ ID NO:2;
 - residues 15-230 of SEQ ID NO:2;
 - residues 15-345 of SEQ ID NO:2;
 - residues 46-345 of SEQ ID NO:2;
 - residues 164-345 of SEQ ID NO:2;
 - residues 235-345 of SEQ ID NO:2; and
 - residues 226-345 of SEQ ID NO:2.
4. An isolated polypeptide comprising a sequence of amino acids of the formula $R1_x-R2_y-R3_z$, wherein:

R1 comprises a polypeptide of from 100 to 120 residues in length that is at least 90% identical to residues 46-163 of SEQ ID NO:2, and comprises a sequence motif C[KR]Y[DNE][WYF]X_{11,15}G[KR][WYF]C (SEQ ID NO:4) corresponding to residues 104-124 of SEQ ID NO:2;

R2 is a polypeptide at least 90% identical to residues 164-234 of SEQ ID NO:2;

R3 is a polypeptide at least 90% identical in amino acid sequence to residues 235-345 of SEQ ID NO:2 and comprises cysteine residues at positions corresponding to residues 250, 280, 284, 296, 335, and 337 of SEQ ID NO:2; a glycine residue at a position corresponding to residue 282 of SEQ ID NO:2; and a sequence motif CX_{18,33}CXGX_{6,33}CX_{20,40}CXC (SEQ ID NO:3) corresponding to residues 250-337 of SEQ ID NO:2; and

each of x, y, and z is individually 0 or 1, subject to the limitations that:

at least one of x and z is 1; and

if x and z are each 1, then y is 1.

5. The isolated polypeptide of claim 4 wherein x=1.

6. The isolated polypeptide of claim 5 wherein R1 comprises residues 46-163 of SEQ ID NO:2.

7. The isolated polypeptide of claim 5 wherein R1 is at least 90% identical to residues 18-163 of SEQ ID NO:2.

8. The isolated polypeptide of claim 5 wherein y=1.

9. The isolated polypeptide of claim 8 wherein z=1.

10. The isolated polypeptide of claim 4 wherein said polypeptide comprises residues 46-229 of SEQ ID NO:2, residues 164-345 of SEQ ID NO:2, or residues 46-345 of SEQ ID NO:2.

11. The isolated polypeptide of claim 4 wherein $z=1$.
12. The isolated polypeptide of claim 11 wherein R3 comprises residues 235-345 of SEQ ID NO:2.
13. The isolated polypeptide of claim 11 wherein $y=1$.
14. The isolated polypeptide of claim 13 wherein $x=1$.
15. The isolated polypeptide of claim 11, further comprising cysteine residues at positions corresponding to residues 286, 287, 291, and 294 of SEQ ID NO:2.
16. The isolated polypeptide of any of claims 4-15, further comprising an affinity tag.
17. The isolated polypeptide of any of claims 4-15, further comprising an immunoglobulin constant domain.
18. An isolated protein comprising a first polypeptide operably linked to a second polypeptide, wherein said first polypeptide comprises a sequence of amino acids of the formula $R1_x-R2_y-R3_z$, wherein:
 - R1 comprises a polypeptide of from 100 to 120 residues in length that is at least 90% identical to residues 46-163 of SEQ ID NO:2, and comprises a sequence motif C[KR]Y[DNE][WYF]X{11,15}G[KR][WYF]C (SEQ ID NO:4) corresponding to residues 104-124 of SEQ ID NO:2;
 - R2 is a polypeptide at least 90% identical to residues 164-234 of SEQ ID NO:2;
 - R3 is a polypeptide at least 90% identical in amino acid sequence to residues 235-345 of SEQ ID NO:2 and comprises cysteine residues at positions

corresponding to residues 250, 280, 284, 296, 335, and 337 of SEQ ID NO:2; a glycine residue at a position corresponding to residue 282 of SEQ ID NO:2; and a sequence motif CX{25,33}CXGXCX{10,33}CX{20,40}CXC (SEQ ID NO:3) corresponding to residues 250-337 of SEQ ID NO:2; and

each of x, y, and z is individually 0 or 1, subject to the limitations that:

at least one of x and z is 1; and

if x and z are each 1, then y is 1,

and wherein said protein modulates cell proliferation, differentiation, metabolism, or migration.

19. The isolated protein of claim 18 wherein said protein is a heterodimer.

20. The isolated protein of claim 19 wherein z is 1 and said second polypeptide is selected from the group consisting of VEGF, VEGF-B, VEGF-C, VEGF-D, PIGF, PDGF-A, and PDGF-B.

21. The isolated protein of claim 18 wherein said protein is a homodimer.

22. The isolated protein of claim 21 wherein z=1.

23. The isolated protein according to claim 22 wherein each of said first and second polypeptides comprises residues 235-345 of SEQ ID NO:2.

24. The isolated protein of claim 21 wherein x=1.

25. The isolated protein of claim 24 wherein each of said first and second polypeptides comprises residues 46-163 of SEQ ID NO:2.

26. An isolated polynucleotide of up to approximately 4 kb in length, wherein said polynucleotide encodes a polypeptide comprising a sequence of amino acids of the formula $R1_x-R2_y-R3_z$, wherein:

R1 comprises a polypeptide of from 100 to 120 residues in length that is at least 90% identical to residues 46-163 of SEQ ID NO:2, and comprises a sequence motif C[KR]Y[DNE][WYF]X{11,15}G[KR][WYF]C (SEQ ID NO:4) corresponding to residues 104-124 of SEQ ID NO:2;

R2 is a polypeptide at least 90% identical to residues 164-234 of SEQ ID NO:2;

R3 is a polypeptide at least 90% identical in amino acid sequence to residues 235-345 of SEQ ID NO:2 and comprises cysteine residues at positions corresponding to residues 250, 280, 284, 296, 335, and 337 of SEQ ID NO:2; a glycine residue at a position corresponding to residue 282 of SEQ ID NO:2; and a sequence motif CX{25,33}CXGXCX{10,33}CX{20,40}CXC (SEQ ID NO:3) corresponding to residues 250-337 of SEQ ID NO:2; and

each of x, y, and z is individually 0 or 1, subject to the limitations that:
at least one of x and z is 1; and
if x and z are each 1, then y is 1.

27. The polynucleotide of claim 26, wherein said polynucleotide is DNA.

28. The polynucleotide of claim 27 comprising nucleotides 1 through 1035 of SEQ ID NO:6.

29. The polynucleotide of claim 27 comprising nucleotides 154 through 1188 of SEQ ID NO:1.

30. The polynucleotide of claim 27 comprising nucleotides 1049 through 2083 of SEQ ID NO:42.

31. An expression vector comprising the following operably linked elements:

a transcription promoter;
a DNA polynucleotide according to claim 27; and
a transcription terminator.

32. The expression vector of claim 31, further comprising a secretory signal sequence operably linked to the DNA polynucleotide.

33. A cultured cell into which has been introduced an expression vector according to claim 31 or claim 32, wherein said cell expresses the polypeptide encoded by the DNA segment.

34. A composition comprising a protein according to any of claims 18-25 in combination with a pharmaceutically acceptable vehicle.

35. A method of producing a protein comprising:
culturing a cell into which has been introduced an expression vector according to claim 31 or claim 32, whereby said cell expresses the polypeptide encoded by the DNA segment; and
recovering the expressed protein.

36. An antibody that specifically binds to an epitope of a polypeptide according to claim 4.

37. The antibody of claim 36 which is a monoclonal antibody.

38. The antibody of claim 36 which is a single-chain antibody.

39. The antibody of any of claims 36-38 operably linked to a reporter molecule.

40. A method for detecting a genetic abnormality in a patient, comprising:

obtaining a genetic sample from a patient;

incubating the genetic sample with a polynucleotide comprising at least 14 contiguous nucleotides of SEQ ID NO:1 or the complement of SEQ ID NO:1, under conditions wherein said polynucleotide will hybridize to complementary polynucleotide sequence, to produce a first reaction product; and

comparing said first reaction product to a control reaction product, wherein a difference between said first reaction product and said control reaction product is indicative of a genetic abnormality in the patient.

41. A method of stimulating the growth of fibroblasts or smooth muscle cells comprising applying to said cells an effective amount of a protein of any of claims 18-25.

42. A method of activating a cell-surface PDGF alpha receptor, comprising exposing a cell comprising a cell-surface PDGF alpha receptor to the polypeptide or protein of any of claims 2-25, whereby the polypeptide or protein binds to and activates the receptor.

43. A method of inhibiting a PDGF alpha receptor-mediated cellular process, comprising exposing a cell comprising a cell-surface PDGF alpha receptor to a compound that inhibits binding of a polypeptide or protein of any of claims 2-25 to the receptor.

44. A method of inhibiting zveg3 activity in a mammal comprising administering to the mammal an effective amount of a zveg3 antagonist.

45. The method of claim 44 wherein the antagonist is an antibody, a receptor, a ligand-binding receptor fragment, or a receptor IgG-Fc fusion protein.

46. An isolated, antisense polynucleotide that is the complement of the isolated polynucleotide of claim 26.

47. The antisense polynucleotide of claim 46 further comprising operably linked transcription promoter and terminator sequences.

48. A method of inhibiting zveg3 production in a cell comprising administering to the cell the antisense polynucleotide of claim 47.

1/12




Fig. 1

2/12

50	-1.10	=====	I
51	-0.80	=====	t
52	-0.18	==	V
53	-0.23	==	S
54	-0.37	=====	T
55	-0.25	===	N
56	-0.28	===	G
57	0.22		S ==
58	-0.25	===	I
59	0.05		H
60	0.05		S
61	-0.07	=	P
62	-0.05		R
63	-0.55	=====	F
64	0.37		P =====
65	0.40		H =====
66	0.42		T =====
67	0.23		Y ==
68	-0.08	=	P
69	-0.33	===	R
70	-1.40	=====	N
71	-0.93	=====	T
72	-1.17	=====	V
73	-1.17	=====	L
74	-0.95	=====	V
75	-0.95	=====	W
76	0.12		R =
77	0.12		L =
78	0.45		V =====
79	0.45		A =====
80	-0.03		V
81	-0.08	=	E
82	-0.55	=====	E
83	-1.35	=====	N
84	-1.45	=====	V
85	-1.62	=====	W
86	-0.55	=====	I
87	0.25		Q ==
88	0.72		L =====
89	0.60		T =====
90	0.67		F =====
91	0.78		D =====
92	0.78		E =====
93	0.78		R =====
94	0.28		F ==
95	1.20		G =====
96	1.70		L =====
97	2.50		E =====
98	1.70		D =====
99	1.03		P =====
100	1.53		E =====
101	0.65		D =====
102	0.65		D =====
103	-0.27	===	I
104	-0.22	==	C
105	0.45		K =====

3/12

106	-0.30	====	Y
107	0.58		D =====
108	0.58		F =====
109	1.00		V =====
110	1.30		E =====
111	1.30		V =====
112	1.55		E =====
113	0.98		E =====
114	0.18		P ==
115	-0.12		= S
116	-0.17		== D
117	-0.17		== G
118	-0.73	=====	T
119	-0.83	=====	I
120	-0.53	=====	L
121	-0.80	=====	G
122	-0.80	=====	R
123	-1.37	=====	W
124	-1.05	=====	C
125	-0.88	=====	G
126	-0.88	=====	S
127	0.18		G ==
128	0.22		T ==
129	-0.02		V
130	0.28		P ===
131	0.78		G =====
132	0.78		K =====
133	0.32		Q ===
134	0.32		I ===
135	0.32		S ===
136	0.77		K =====
137	-0.03		G
138	0.47		N =====
139	0.02		Q
140	-0.27	====	I
141	0.08		R =
142	0.08		I =
143	0.88		R =====
144	0.00		F
145	0.00		V
146	0.25		S ===
147	0.25		D ===
148	0.25		E ===
149	-0.25	====	Y
150	0.13		F =
151	0.13		P =
152	-0.03		S
153	-0.38	=====	E
154	-0.97	=====	P
155	-1.35	=====	G
156	-1.32	=====	F
157	-1.20	=====	C
158	-1.28	=====	I
159	-1.20	=====	H
160	-1.12	=====	Y
161	-0.70	=====	N

4/12

162	-1.15	=====	I
163	-0.92	=====	V
164	-0.17	==	M
165	-0.03		P
166	-0.28	===	Q
167	-0.27	===	F
168	0.15		T =
169	0.27		E ===
170	-0.48	=====	A
171	-0.70	=====	V
172	-0.45	=====	S
173	-0.50	=====	P
174	-0.45	=====	S
175	-0.58	=====	V
176	-0.63	=====	L
177	-0.33	===	P
178	-0.63	=====	P
179	-0.13	=	S
180	-0.48	=====	A
181	-0.70	=====	L
182	-0.37	=====	P
183	-0.33	=====	L
184	-0.12	=	D
185	-0.92	=====	L
186	-1.18	=====	L
187	-0.97	=====	N
188	-1.42	=====	N
189	-2.02	=====	A
190	-2.00	=====	I
191	-2.00	=====	t
192	-0.93	=====	A
193	-0.35	=====	F
194	-0.23	==	S
195	0.03		T
196	0.60		L =====
197	0.52		E =====
198	-0.28	===	D
199	-0.28	===	L
200	0.02		I
201	0.82		R =====
202	0.82		Y =====
203	1.03		L =====
204	1.37		E =====
205	0.57		P =====
206	1.07		E =====
207	0.27		R ===
208	0.27		w ===
209	0.93		Q =====
210	0.60		L =====
211	0.52		D =====
212	0.52		L =====
213	0.82		E =====
214	0.25		D ===
215	-0.42	=====	L
216	-0.08	=	Y
217	0.00		R

5/12

218	-0.80	=====	P	
219	-0.80	=====	T	
220	-0.23	==	w	
221	-0.15	=	Q	
222	-0.60	=====	L	
223	-0.55	=====	L	
224	-0.67	=====	G	
225	-0.67	=====	K	
226	-0.67	=====	A	
227	-0.08	=	F	
228	0.38		V	=====
229	1.13		F	=====
230	1.30		G	=====
231	1.05		R	=====
232	1.05		K	=====
233	0.25		S	=====
234	0.23		R	=====
235	-0.57	=====	V	
236	-0.62	=====	V	
237	-0.43	=====	D	
238	-0.43	=====	L	
239	0.37		N	=====
240	0.08		L	=====
241	0.88		L	=====
242	0.88		T	=====
243	0.57		E	=====
244	0.12		E	=====
245	-0.55	=====	V	
246	-0.37	=====	R	
247	-0.87	=====	L	
248	-0.07	=	Y	
249	0.35		S	=====
250	-0.12	=	C	
251	0.10		T	=====
252	-0.08	=	P	
253	-0.03		R	
254	-0.83	=====	N	
255	-0.37	=====	F	
256	0.55		S	=====
257	1.00		V	=====
258	0.95		S	=====
259	1.40		I	=====
260	2.20		R	=====
261	1.63		E	=====
262	1.63		E	=====
263	1.07		L	=====
264	1.07		K	=====
265	0.15		R	=====
266	-0.92	=====	T	
267	-0.85	=====	D	
268	-1.35	=====	T	
269	-1.45	=====	I	
270	-1.45	=====	F	
271	-1.33	=====	W	
272	-1.02	=====	P	
273	-0.52	=====	G	

6/12

274	-0.02		C
275	-0.02		L
276	0.28		L ===
277	0.58		V =====
278	0.87		K =====
279	0.20		R ==
280	-0.38	=====	C
281	-0.38	=====	G
282	-0.55	=====	G
283	-0.85	=====	N
284	-0.97	=====	C
285	-0.77	=====	A
286	-0.85	=====	C
287	-0.65	=====	C
288	0.02		L
289	0.15		H ==
290	0.27		N ===
291	0.07		C =
292	-0.02		N
293	-0.05		E
294	-0.50	=====	C
295	0.17		Q ==
296	-0.12	=	C
297	-0.02		V
298	0.73		P =====
299	1.23		S =====
300	1.20		K =====
301	0.62		V =====
302	1.37		T =====
303	1.18		K =====
304	0.38		K =====
305	-0.08	=	Y
306	-0.40	=====	H
307	0.18		E ==
308	-0.32	=====	V
309	0.43		L =====
310	0.67		Q =====
311	0.63		L =====
312	0.68		R =====
313	0.68		P =====
314	0.68		K =====
315	-0.12	=	T
316	-0.13	=	G
317	0.37		V =====
318	0.67		R =====
319	-0.13	=	G
320	-0.20	=====	L
321	0.60		H =====
322	0.43		K =====
323	-0.15	=	S
324	-0.50	=====	L
325	0.30		T =====
326	0.28		D =====
327	0.10		V =
328	0.85		A =====
329	1.43		L =====

7/12

330	1.57	E	=====
331	1.57	H	=====
332	1.48	h	=====
333	0.92	E	=====
334	0.25	E	=====
335	0.36	C	=====
336	0.64	D	=====
337	0.30	C	=====
338	0.51	V	=====
339	0.87	C	=====
340	1.15	R	=====
341	0.00	G	
342	0.00	S	
343	0.00	T	
344	0.00	G	
345	0.00	G	

-----|-----|-----

-2 -1 0

Hydrophobic

-----|-----|-----

0 1 2

Hydrophilic




Fig. 2

BEST AVAILABLE COPY




Fig. 3




Fig. 4

11/12




Fig. 5

12/12

MSLFGLLLLT SALAGQRQGTQAESNLSSKFQFSSNKEQNGVQDPQHERIITVSTNGSIHS
MLLLGLLLLT SALAGQRTGTRAESNLSSKLQLSSDKEQNGVQDPRHERVVTISGNGSIHS
10 20 30 40 50 60

PRFPHTYPRNTVLVWRLVAVEENVWIQLTFDERFGLEDPEDDICKYDFVEVEEPSDGTIL
PKFPHTYPRNMVLVWRLVAVDENVRIQLTFDERFGLEDPEDDICKYDFVEVEEPSDGSVL
70 80 90 100 110 120

GRWCGSGTVPKGQISKGNQIRIRFVSDEYFPSEPGFCIHYNIVMPQFTEAVSPSVLPSSA
GRWCGSGTVPKGQTSKGNHIRIRFVSDEYFPSEPGFCIHYSIIMPQVTETTSPSVLPSS
130 140 150 160 170 180

LPLDLLNNAITAFSTLEDLIRYLEPERWQLDLEDLYRPTWQLLGKAFVFGRKSRVVDLNL
LSLDLLNNAVTAFTLEELIRYLEPDRWQVDLDSLYKPTWQLLGKAFLYGKKSKVNLNL
190 200 210 220 230 240

LTEEVRLYSCTPRNFSVSIREELKRTDTIFWPGCLLVKRCGGNCACCLHNCNECQCVPSK
LKEEVKLYSCTPRNFSVSIREELKRTDTIFWPGCLLVKRCGGNCACCLHNCNECQCVPRK
250 260 270 280 290 300

VTKKYHEVLQLRPKTGVRGLHKSLTDVALEHHEECDVCVRGSTGG (SEQ ID NO:2)
VTKKYHEVLQLRPKTGVKGLHKSLTDVALEHHEECDVCVRGNAGG (SEQ ID NO:43)
310 320 330 340

Fig. 6

SEQUENCE LISTING

<110> ZymoGenetics. Inc.

<120> GROWTH FACTOR HOMOLOG ZVEGF3

<130> 98-60PC

<160> 50

<170> FastSEQ for Windows Version 3.0

<210> 1

<211> 1760

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (154)...(1191)

<400> 1

attatgtgga aactaccctg cgattctctg ctgccagagc aggcctcggcg cttccacccc	60
agtcgagcct tcccctggcg gtggtgaaag agactcggga gtcgctgctt ccaaagtgcc	120
cgccgtgagt gagctctcac ccagtcagc caa atg agc ctc ttc ggg ctt ctc	174
Met Ser Leu Phe Gly Leu Leu	
1 5	
ctg ctg aca tct gcc ctg gcc ggc cag aga cag ggg act cag gcg gaa	222
Leu Leu Thr Ser Ala Leu Ala Gly Gln Arg Gln Gly Thr Gln Ala Glu	
10 15 20	
tcc aac ctg agt agt aaa ttc cag ttt tcc agc aac aag gaa cag aac	270
Ser Asn Leu Ser Ser Lys Phe Gln Phe Ser Ser Asn Lys Glu Gln Asn	
25 30 35	
gga gta caa gat cct cag cat gag aga att att act gtg tct act aat	318
Gly Val Gln Asp Pro Gln His Glu Arg Ile Ile Thr Val Ser Thr Asn	
40 45 50 55	
gga agt att cac agc cca agg ttt cct cat act tat cca aga aat acg	366
Gly Ser Ile His Ser Pro Arg Phe Pro His Thr Tyr Pro Arg Asn Thr	
60 65 70	

2

gtc ttg gta tgg aga tta gta gca gta gag gaa aat gta tgg ata caa Val Leu Val Trp Arg Leu Val Ala Val Glu Glu Asn Val Trp Ile Gln 75 80 85	414
ctt acg ttt gat gaa aga ttt ggg ctt gaa gac cca gaa gat gac ata Leu Thr Phe Asp Glu Arg Phe Gly Leu Glu Asp Pro Glu Asp Asp Ile 90 95 100	462
tgc aag tat gat ttt gta gaa gtt gag gaa ccc agt gat gga act ata Cys Lys Tyr Asp Phe Val Glu Val Glu Glu Pro Ser Asp Gly Thr Ile 105 110 115	510
tta ggg cgc tgg tgt ggt tct ggt act gta cca gga aaa cag att tct Leu Gly Arg Trp Cys Gly Ser Gly Thr Val Pro Gly Lys Gln Ile Ser 120 125 130 135	558
aaa gga aat caa att agg ata aga ttt gta tct gat gaa tat ttt cct Lys Gly Asn Gln Ile Arg Ile Arg Phe Val Ser Asp Glu Tyr Phe Pro 140 145 150	606
tct gaa cca ggg ttc tgc atc cac tac aac att gtc atg cca caa ttc Ser Glu Pro Gly Phe Cys Ile His Tyr Asn Ile Val Met Pro Gln Phe 155 160 165	654
aca gaa gct gtg agt cct tca gtg cta ccc cct tca gct ttg cca ctg Thr Glu Ala Val Ser Pro Ser Val Leu Pro Pro Ser Ala Leu Pro Leu 170 175 180	702
gac ctg ctt aat aat gct ata act gcc ttt agt acc ttg gaa gac ctt Asp Leu Leu Asn Asn Ala Ile Thr Ala Phe Ser Thr Leu Glu Asp Leu 185 190 195	750
att cga tat ctt gaa cca gag aga tgg cag ttg gac tta gaa gat cta Ile Arg Tyr Leu Glu Pro Glu Arg Trp Gln Leu Asp Leu Glu Asp Leu 200 205 210 215	798
tat agg cca act tgg caa ctt ctt ggc aag gct ttt gtt ttt gga aga Tyr Arg Pro Thr Trp Gln Leu Leu Gly Lys Ala Phe Val Phe Gly Arg 220 225 230	846
aaa tcc aga gtg gtg gat ctg aac ctt cta aca gag gag gta aga tta Lys Ser Arg Val Val Asp Leu Asn Leu Leu Thr Glu Glu Val Arg Leu 235 240 245	894

3

tac agc tgc aca cct cgt aac ttc tca gtg tcc ata agg gaa gaa cta 942
 Tyr Ser Cys Thr Pro Arg Asn Phe Ser Val Ser Ile Arg Glu Glu Leu
 250 255 260

aag aga acc gat acc att ttc tgg cca ggt tgt ctc ctg gtt aaa cgc 990
 Lys Arg Thr Asp Thr Ile Phe Trp Pro Gly Cys Leu Leu Val Lys Arg
 265 270 275

tgt ggt ggg aac tgt gcc tgt tgt ctc cac aat tgc aat gaa tgt caa 1038
 Cys Gly Gly Asn Cys Ala Cys Cys Leu His Asn Cys Asn Glu Cys Gln
 280 285 290 295

tgt gtc cca agc aaa gtt act aaa aaa tac cac gag gtc ctt cag ttg 1086
 Cys Val Pro Ser Lys Val Thr Lys Lys Tyr His Glu Val Leu Gln Leu
 300 305 310

aga cca aag acc ggt gtc agg gga ttg cac aaa tca ctc acc gac gtg 1134
 Arg Pro Lys Thr Gly Val Arg Gly Leu His Lys Ser Leu Thr Asp Val
 315 320 325

gcc ctg gag cac cat gag gag tgt gac tgt gtg tgc aga ggg agc aca 1182
 Ala Leu Glu His His Glu Glu Cys Asp Cys Val Cys Arg Gly Ser Thr
 330 335 340

gga gga tag ccgcatcacc accagcagct cttgcccaga gctgtgcagt 1231
 Gly Gly *
 345

gcagtggctg attctattag agaacgtatg cgttatctcc atccttaatc tcagttgttt 1291
 gcttcaagga cctttcatct tcaggattta cagtgcattc tgaaagagga gacatcaaac 1351
 agaattagga gttgtgcaac agctcttttg agaggaggcc taaaggacag gagaaaaggt 1411
 cttcaatcgt ggaaagaaaa ttaaatgttg tattaatatg atcaccagct agtttcagag 1471
 ttaccatgta cgtattccac tagctgggtt ctgtatttca gttctttcga tacggcttag 1531
 ggtaatgtca gtacaggaaa aaaactgtgc aagtgagcac ctgattccgt tgccttgctt 1591
 aactctaaag ctccatgtcc tgggcctaaa atcgtataaa atctggattt tttttttttt 1651
 tttttgctca tattcacata tgtaaaccag aacattctat gtactacaaa cctggttttt 1711
 aaaaaggaac tatgttgcta tgaattaaac ttgtgtcgtg ctgatagga 1760

<210> 2

<211> 345

<212> PRT

<213> Homo sapiens

<400> 2

Met Ser Leu Phe Gly Leu Leu Leu Leu Thr Ser Ala Leu Ala Gly Gln
 1 5 10 15
 Arg Gln Gly Thr Gln Ala Glu Ser Asn Leu Ser Ser Lys Phe Gln Phe
 20 25 30
 Ser Ser Asn Lys Glu Gln Asn Gly Val Gln Asp Pro Gln His Glu Arg
 35 40 45
 Ile Ile Thr Val Ser Thr Asn Gly Ser Ile His Ser Pro Arg Phe Pro
 50 55 60
 His Thr Tyr Pro Arg Asn Thr Val Leu Val Trp Arg Leu Val Ala Val
 65 70 75 80
 Glu Glu Asn Val Trp Ile Gln Leu Thr Phe Asp Glu Arg Phe Gly Leu
 85 90 95
 Glu Asp Pro Glu Asp Asp Ile Cys Lys Tyr Asp Phe Val Glu Val Glu
 100 105 110
 Glu Pro Ser Asp Gly Thr Ile Leu Gly Arg Trp Cys Gly Ser Gly Thr
 115 120 125
 Val Pro Gly Lys Gln Ile Ser Lys Gly Asn Gln Ile Arg Ile Arg Phe
 130 135 140
 Val Ser Asp Glu Tyr Phe Pro Ser Glu Pro Gly Phe Cys Ile His Tyr
 145 150 155 160
 Asn Ile Val Met Pro Gln Phe Thr Glu Ala Val Ser Pro Ser Val Leu
 165 170 175
 Pro Pro Ser Ala Leu Pro Leu Asp Leu Leu Asn Asn Ala Ile Thr Ala
 180 185 190
 Phe Ser Thr Leu Glu Asp Leu Ile Arg Tyr Leu Glu Pro Glu Arg Trp
 195 200 205
 Gln Leu Asp Leu Glu Asp Leu Tyr Arg Pro Thr Trp Gln Leu Leu Gly
 210 215 220
 Lys Ala Phe Val Phe Gly Arg Lys Ser Arg Val Val Asp Leu Asn Leu
 225 230 235 240
 Leu Thr Glu Glu Val Arg Leu Tyr Ser Cys Thr Pro Arg Asn Phe Ser
 245 250 255
 Val Ser Ile Arg Glu Glu Leu Lys Arg Thr Asp Thr Ile Phe Trp Pro
 260 265 270
 Gly Cys Leu Leu Val Lys Arg Cys Gly Gly Asn Cys Ala Cys Cys Leu
 275 280 285
 His Asn Cys Asn Glu Cys Gln Cys Val Pro Ser Lys Val Thr Lys Lys
 290 295 300
 Tyr His Glu Val Leu Gln Leu Arg Pro Lys Thr Gly Val Arg Gly Leu
 305 310 315 320
 His Lys Ser Leu Thr Asp Val Ala Leu Glu His His Glu Glu Cys Asp
 325 330 335
 Cys Val Cys Arg Gly Ser Thr Gly Gly
 340 345

<210> 3
<211> 116
<212> PRT
<213> Artificial Sequence

<220>
<223> peptide motif

<221> VARIANT
<222> (2)...(19)
<223> Xaa is any amino acid

<221> VARIANT
<222> (20)...(34)
<223> Xaa is any amino acid or not present

<221> VARIANT
<222> (36)...(36)
<223> Xaa is any amino acid

<221> VARIANT
<222> (38)...(38)
<223> Xaa is any amino acid

<221> VARIANT
<222> (40)...(45)
<223> Xaa is any amino acid

<221> VARIANT
<222> (46)...(72)
<223> Xaa is any amino acid or not present

<221> VARIANT
<222> (74)...(93)
<223> Xaa is any amino acid

<221> VARIANT
<222> (94)...(113)
<223> Xaa is any amino acid not present

<221> VARIANT
<222> (115)...(115)
<223> Xaa is any amino acid

<400> 3

6

Cys Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa
 1 5 10 15
 Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa
 20 25 30
 Xaa Xaa Cys Xaa Gly Xaa Cys Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa
 35 40 45
 Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa
 50 55 60
 Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Cys Xaa Xaa Xaa Xaa Xaa Xaa Xaa
 65 70 75 80
 Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa
 85 90 95
 Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa
 100 105 110
 Xaa Cys Xaa Cys
 115

<210> 4
 <211> 24
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> peptide motif

<221> VARIANT
 <222> (2)...(2)
 <223> Xaa is Lys or Arg

<221> VARIANT
 <222> (4)...(4)
 <223> Xaa is Asp, Asn or Glu

<221> VARIANT
 <222> (5)...(5)
 <223> Xaa is Trp, Tyr or Phe

<221> VARIANT
 <222> (6)...(16)
 <223> Xaa is any amino acid

<221> VARIANT
 <222> (17)...(20)
 <223> Xaa is any amino acid or not present

<221> VARIANT
 <222> (22)...(22)
 <223> Xaa is Lys or Arg

<221> VARIANT
 <222> (23)...(23)
 <223> Xaa is Trp. Tyr or Phe

<400> 4
 Cys Xaa Tyr Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa
 1 5 10 15
 Xaa Xaa Xaa Xaa Gly Xaa Xaa Cys
 20

<210> 5
 <211> 6
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> peptide tag

<400> 5
 Glu Tyr Met Pro Met Glu
 1 5

<210> 6
 <211> 1035
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> degenerate sequence derived from SEQ ID NOS: 1 and
 2

<221> misc_feature
 <222> (1)...(1035)
 <223> n = A,T,C or G

<400> 6
 atgwsnytny tyggnytnyt nytnytnacn wsngcnytny cnggncarmg ncarggnacn 60
 cargcngarw snaayytnws nwsnaartty carttywsnw snaayaarga rcaraayggn 120
 gtncargayc cncarcayga rmgnathath acngtnwsna cnaayggnws nathcaywsn 180
 ccnmgnttyc cncayacnta yccnmgnaay acngtnytny tntggmgnyt ngtnngcngtn 240
 gargaraayg tntggathca rytnacntty gaygarmgnt tyggnytnga rgayccngar 300

gaygayatht	gyaartayga	yttygtngar	gtngargarc	cnwsngaygg	nacnathytn	360
ggnmntggt	gyggnwsngg	nacngtnccn	ggnaarcara	thwsnaargg	naaycarath	420
mgnathmgt	tygtwnsnga	ygartaytty	ccnwsngarc	cnggnttytg	yathcaytay	480
aayathgtna	tgccncartt	yacngargcn	gtnwsnccnw	sngtnytnc	nccnwsngcn	540
ytnccnytn	ayytnytnaa	yaaygcnath	acngcnttyw	snacnytna	rgayytnath	600
mgntayytn	arccngarmg	ntggcarytn	gayytnarg	ayytnaymg	nccnacntgg	660
carytnytn	gnaargcntt	ygtnttygg	mgnaarwsnm	gngtngtna	yytnaayytn	720
ytnacngarg	argtnmgnyt	ntaywsntgy	acnccnmgna	aytywsngt	nwsnathmgn	780
gargarytna	armgnacnga	yacnathtty	tggccnggnt	gyytnytn	naarmgntgy	840
ggnggnaayt	gygcntgytg	yytncayaay	tgyaaygart	gycartgygt	nccnwsnaar	900
gtnacnaara	artaycayga	rgtnytnar	ytnmgncna	aracnggnt	nmgnggnytn	960
cayaarwsny	tnacngaygt	ngcnytnar	caycaygarg	artgygaytg	ygtntgymgn	1020
ggnwsnacng	gnggn					1035

<210> 7

<211> 17

<212> DNA

<213> Artificial Sequence

<220>

<223> Oligonucleotide primer

<221> misc_feature

<222> (1)...(17)

<223> n = A.T.C or G

<400> 7

mgntgyggng gnaaytg

17

<210> 8

<211> 17

<212> DNA

<213> Artificial Sequence

<220>

<223> Oligonucleotide primer

<221> misc_feature

<222> (1)...(17)

<223> n = A.T.C or G

<400> 8

mgntgydsng gnwrytg

17

<210> 9

<211> 17
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Oligonucleotide primer

<221> misc_feature
 <222> (1)...(17)
 <223> n = A.T.C or G

<400> 9
 carywnccns hrcanck 17

<210> 10
 <211> 17
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Oligonucleotide primer

<221> misc_feature
 <222> (1)...(17)
 <223> n = A.T.C or G

<400> 10
 ttytgccng gntgyyt 17

<210> 11
 <211> 17
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Oligonucleotide primer

<221> misc_feature
 <222> (1)...(17)
 <223> n = A.T.C or G

<400> 11
 ntnddnccnn sntgybt 17

<210> 12

10

<211> 17
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Oligonucleotide primer

<221> misc_feature
 <222> (1)...(17)
 <223> n = A.T.C or G

<400> 12
 avrcansnng gnhhnan 17

<210> 13
 <211> 17
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Oligonucleotide primer

<400> 13
 caygargart gygaytg 17

<210> 14
 <211> 17
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Oligonucleotide primer

<221> misc_feature
 <222> (1)...(17)
 <223> n = A.T.C or G

<400> 14
 caynnnnvnt gyvvntg 17

<210> 15
 <211> 17
 <212> DNA
 <213> Artificial Sequence

11

<220>
<223> Oligonucleotide primer

<221> misc_feature
<222> (1)...(17)
<223> n = A.T.C or G

<400> 15
canbbrcanb nnnnrtg 17

<210> 16
<211> 17
<212> DNA
<213> Artificial Sequence

<220>
<223> Oligonucleotide primer

<221> misc_feature
<222> (1)...(17)
<223> n = A.T.C or G

<400> 16
tgyacnccnm gnaaytt 17

<210> 17
<211> 17
<212> DNA
<213> Artificial Sequence

<220>
<223> Oligonucleotide primer

<221> misc_feature
<222> (1)...(17)
<223> n = A.T.C or G

<400> 17
tgyhnnmcnm knrmdh 17

<210> 18
<211> 17
<212> DNA
<213> Artificial Sequence

12

<220>
<223> Oligonucleotide primer

<221> misc_feature
<222> (1)...(17)
<223> n = A.T.C or G

<400> 18
dhnkynmkng knndrca

17

<210> 19
<211> 17
<212> DNA
<213> Artificial Sequence

<220>
<223> Oligonucleotide primer

<400> 19
tgyaartayg aytwygt

17

<210> 20
<211> 17
<212> DNA
<213> Artificial Sequence

<220>
<223> Oligonucleotide primer

<400> 20
acrwartcrt aytrca

17

<210> 21
<211> 17
<212> DNA
<213> Artificial Sequence

<220>
<223> Oligonucleotide primer

<221> misc_feature
<222> (1)...(17)
<223> n = A.T.C or G

<400> 21

13

ywnngnmrnt dbtgygg

17

<210> 22

<211> 17

<212> DNA

<213> Artificial Sequence

<220>

<223> Oligonucleotide primer

<221> misc_feature

<222> (1)...(17)

<223> n = A.T.C or G

<400> 22

ccrcavhany knccnwr

17

<210> 23

<211> 17

<212> DNA

<213> Artificial Sequence

<220>

<223> Oligonucleotide primer

<221> misc_feature

<222> (1)...(17)

<223> n = A.T.C or G

<400> 23

tdbccnmand vntaycc

17

<210> 24

<211> 17

<212> DNA

<213> Artificial Sequence

<220>

<223> Oligonucleotide primer

<221> misc_feature

<222> (1)...(17)

<223> n = A.T.C or G

<400> 24

14

ggrtanbhnt knggvha

17

<210> 25

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Oligonucleotide primer

<400> 25

agcaggtcca gtggcaaagc

20

<210> 26

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> Oligonucleotide primer

<400> 26

cgtttgatga aagatttggg c

21

<210> 27

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> Oligonucleotide primer

<400> 27

ggaggtctat ataagcagag c

21

<210> 28

<211> 18

<212> DNA

<213> Artificial Sequence

<220>

<223> Oligonucleotide primer

<400> 28

taacagagga ggtaagat

18

15

<210> 29
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> Oligonucleotide primer

<400> 29
tcggttctct ttagttct 18

<210> 30
<211> 25
<212> DNA
<213> Artificial Sequence

<220>
<223> Oligonucleotide primer

<400> 30
tctggacgtc ctctgctgg tatag 25

<210> 31
<211> 25
<212> DNA
<213> Artificial Sequence

<220>
<223> Oligonucleotide primer

<400> 31
ggtatggagc caggggcaag ttggg 25

<210> 32
<211> 27
<212> DNA
<213> Artificial Sequence

<220>
<223> Oligonucleotide primer

<400> 32
gagtggcaac ttccagggcc aggagag 27

16

<210> 33
 <211> 27
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Oligonucleotide primer

<400> 33
 cttttgctag cctcaaccct gactatc 27

<210> 34
 <211> 35
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Oligonucleotide primer ZC20.180

<400> 34
 cgcgcggttt aaacgccacc atgagcctct tcggg 35

<210> 35
 <211> 32
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Oligonucleotide primer ZC20.181

<400> 35
 cgtatcggcg cgccctatcc tcctgtgctc cc 32

<210> 36
 <211> 1882
 <212> DNA
 <213> Homo sapiens

<220>
 <221> CDS
 <222> (226)...(1338)

<400> 36
 ccgtcaccat ttatcagctc agcaccacaa ggaagtgcgg caccacacg cgctcggaag 60
 gttcagcatg caggaagttt ggggagagct cggcgattag cacagcgacc cgggcccagcg 120

cagggcgagc	gcaggcggcg	agagcgcagg	gcggcgcggc	gtcgggtcccg	ggagcagaac	180
ccggcttttt	cttggagcga	cgctgtctct	agtcgctgat	cccaa	atg cac cgg ctc	237
					Met His Arg Leu	
					1	
atc ttt gtc tac act cta atc tgc gca aac ttt tgc agc tgt cgg gac	285					
Ile Phe Val Tyr Thr Leu Ile Cys Ala Asn Phe Cys Ser Cys Arg Asp						
5 10 15 20						
act tct gca acc ccg cag agc gca tcc atc aaa gct ttg cgc aac gcc	333					
Thr Ser Ala Thr Pro Gln Ser Ala Ser Ile Lys Ala Leu Arg Asn Ala						
25 30 35						
aac ctc agg cga gat gag agc aat cac ctc aca gac ttg tac cga aga	381					
Asn Leu Arg Arg Asp Glu Ser Asn His Leu Thr Asp Leu Tyr Arg Arg						
40 45 50						
gat gag acc atc cag gtg aaa gga aac ggc tac gtg cag agt cct aga	429					
Asp Glu Thr Ile Gln Val Lys Gly Asn Gly Tyr Val Gln Ser Pro Arg						
55 60 65						
ttc ccg aac agc tac ccc agg aac ctg ctc ctg aca tgg cgg ctt cac	477					
Phe Pro Asn Ser Tyr Pro Arg Asn Leu Leu Leu Thr Trp Arg Leu His						
70 75 80						
tct cag gag aat aca cgg ata cag cta gtg ttt gac aat cag ttt gga	525					
Ser Gln Glu Asn Thr Arg Ile Gln Leu Val Phe Asp Asn Gln Phe Gly						
85 90 95 100						
tta gag gaa gca gaa aat gat atc tgt agg tat gat ttt gtg gaa gtt	573					
Leu Glu Glu Ala Glu Asn Asp Ile Cys Arg Tyr Asp Phe Val Glu Val						
105 110 115						
gaa gat ata tcc gaa acc agt acc att att aga gga cga tgg tgt gga	621					
Glu Asp Ile Ser Glu Thr Ser Thr Ile Ile Arg Gly Arg Trp Cys Gly						
120 125 130						
cac aag gaa gtt cct cca agg ata aaa tca aga acg aac caa att aaa	669					
His Lys Glu Val Pro Pro Arg Ile Lys Ser Arg Thr Asn Gln Ile Lys						
135 140 145						
atc aca ttc aag tcc gat gac tac ttt gtg gct aaa cct gga ttc aag	717					
Ile Thr Phe Lys Ser Asp Asp Tyr Phe Val Ala Lys Pro Gly Phe Lys						
150 155 160						

18

att tat tat tct ttg ctg gaa gat ttc caa ccc gca gca gct tca gag Ile Tyr Tyr Ser Leu Leu Glu Asp Phe Gln Pro Ala Ala Ala Ser Glu 165 170 175 180	765
acc aac tgg gaa tct gtc aca agc tct att tca ggg gta tcc tat aac Thr Asn Trp Glu Ser Val Thr Ser Ser Ile Ser Gly Val Ser Tyr Asn 185 190 195	813
tct cca tca gta acg gat ccc act ctg att gcg gat gct ctg gac aaa Ser Pro Ser Val Thr Asp Pro Thr Leu Ile Ala Asp Ala Leu Asp Lys 200 205 210	861
aaa att gca gaa ttt gat aca gtg gaa gat ctg ctc aag tac ttc aat Lys Ile Ala Glu Phe Asp Thr Val Glu Asp Leu Leu Lys Tyr Phe Asn 215 220 225	909
cca gag tca tgg caa gaa gat ctt gag aat atg tat ctg gac acc cct Pro Glu Ser Trp Gln Glu Asp Leu Glu Asn Met Tyr Leu Asp Thr Pro 230 235 240	957
cgg tat cga ggc agg tca tac cat gac cgg aag tca aaa gtt gac ctg Arg Tyr Arg Gly Arg Ser Tyr His Asp Arg Lys Ser Lys Val Asp Leu 245 250 255 260	1005
gat agg ctc aat gat gat gcc aag cgt tac agt tgc act ccc agg aat Asp Arg Leu Asn Asp Asp Ala Lys Arg Tyr Ser Cys Thr Pro Arg Asn 265 270 275	1053
tac tcg gtc aat ata aga gaa gag ctg aag ttg gcc aat gtg gtc ttc Tyr Ser Val Asn Ile Arg Glu Glu Leu Lys Leu Ala Asn Val Val Phe 280 285 290	1101
ttt cca cgt tgc ctc ctc gtg cag cgc tgt gga gga aat tgt ggc tgt Phe Pro Arg Cys Leu Leu Val Gln Arg Cys Gly Gly Asn Cys Gly Cys 295 300 305	1149
gga act gtc aac tgg agg tcc tgc aca tgc aat tca ggg aaa acc gtg Gly Thr Val Asn Trp Arg Ser Cys Thr Cys Asn Ser Gly Lys Thr Val 310 315 320	1197
aaa aag tat cat gag gta tta cag ttt gag cct ggc cac atc aag agg Lys Lys Tyr His Glu Val Leu Gln Phe Glu Pro Gly His Ile Lys Arg 325 330 335 340	1245

19

agg ggt aga gct aag acc atg gct cta gtt gac atc cag ttg gat cac 1293
 Arg Gly Arg Ala Lys Thr Met Ala Leu Val Asp Ile Gln Leu Asp His
 345 350 355

cat gaa cga tgc gat tgt atc tgc agc tca aga cca cct cga taa 1338
 His Glu Arg Cys Asp Cys Ile Cys Ser Ser Arg Pro Pro Arg *
 360 365 370

gagaatgtgc acatccttac attaagcctg aaagaacctt tagtttaagg aggggtgagat 1398
 aagagaccct tttcctacca gcaaccaaac ttactactag cctgcaatgc aatgaacaca 1458
 agtggttgct gagtctcagc cttgctttgt taatgccatg gcaagtagaa aggtatatca 1518
 tcaacttcta tacctaagaa tataggattg catttaataa tagtgtttga ggttatatat 1578
 gcacaaacac acacagaaat atattcatgt ctatgtgtat atagatcaaa tgtttttttt 1638
 ttttggtata tataaccagg tacaccagag gttacatatg tttgagttag actcttaaaa 1698
 tcctttgccca aaataaggga tgggtcaaata tatgaaacat gtcttttagaa aatttaggag 1758
 ataaatttat ttttaattt tgaaacacga aacaattttg aatcttgctc tcttaaagaa 1818
 agcatcttgt atattaaaa tcaaaagatg aggctttctt acatatacat cttagttgat 1878
 tatt 1882

<210> 37
 <211> 370
 <212> PRT
 <213> Homo sapiens

<400> 37

Met His Arg Leu Ile Phe Val Tyr Thr Leu Ile Cys Ala Asn Phe Cys
 1 5 10 15
 Ser Cys Arg Asp Thr Ser Ala Thr Pro Gln Ser Ala Ser Ile Lys Ala
 20 25 30
 Leu Arg Asn Ala Asn Leu Arg Arg Asp Glu Ser Asn His Leu Thr Asp
 35 40 45
 Leu Tyr Arg Arg Asp Glu Thr Ile Gln Val Lys Gly Asn Gly Tyr Val
 50 55 60
 Gln Ser Pro Arg Phe Pro Asn Ser Tyr Pro Arg Asn Leu Leu Leu Thr
 65 70 75 80
 Trp Arg Leu His Ser Gln Glu Asn Thr Arg Ile Gln Leu Val Phe Asp
 85 90 95
 Asn Gln Phe Gly Leu Glu Glu Ala Glu Asn Asp Ile Cys Arg Tyr Asp
 100 105 110
 Phe Val Glu Val Glu Asp Ile Ser Glu Thr Ser Thr Ile Ile Arg Gly
 115 120 125
 Arg Trp Cys Gly His Lys Glu Val Pro Pro Arg Ile Lys Ser Arg Thr
 130 135 140
 Asn Gln Ile Lys Ile Thr Phe Lys Ser Asp Asp Tyr Phe Val Ala Lys
 145 150 155 160

20

Pro Gly Phe Lys Ile Tyr Tyr Ser Leu Leu Glu Asp Phe Gln Pro Ala
 165 170 175
 Ala Ala Ser Glu Thr Asn Trp Glu Ser Val Thr Ser Ser Ile Ser Gly
 180 185 190
 Val Ser Tyr Asn Ser Pro Ser Val Thr Asp Pro Thr Leu Ile Ala Asp
 195 200 205
 Ala Leu Asp Lys Lys Ile Ala Glu Phe Asp Thr Val Glu Asp Leu Leu
 210 215 220
 Lys Tyr Phe Asn Pro Glu Ser Trp Gln Glu Asp Leu Glu Asn Met Tyr
 225 230 235 240
 Leu Asp Thr Pro Arg Tyr Arg Gly Arg Ser Tyr His Asp Arg Lys Ser
 245 250 255
 Lys Val Asp Leu Asp Arg Leu Asn Asp Asp Ala Lys Arg Tyr Ser Cys
 260 265 270
 Thr Pro Arg Asn Tyr Ser Val Asn Ile Arg Glu Glu Leu Lys Leu Ala
 275 280 285
 Asn Val Val Phe Phe Pro Arg Cys Leu Leu Val Gln Arg Cys Gly Gly
 290 295 300
 Asn Cys Gly Cys Gly Thr Val Asn Trp Arg Ser Cys Thr Cys Asn Ser
 305 310 315 320
 Gly Lys Thr Val Lys Lys Tyr His Glu Val Leu Gln Phe Glu Pro Gly
 325 330 335
 His Ile Lys Arg Arg Gly Arg Ala Lys Thr Met Ala Leu Val Asp Ile
 340 345 350
 Gln Leu Asp His His Glu Arg Cys Asp Cys Ile Cys Ser Ser Arg Pro
 355 360 365
 Pro Arg
 370

<210> 38
 <211> 20
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> oligonucleotide primer ZC21.222

<400> 38
 tgagccctcg cccagtcag

20

<210> 39
 <211> 25
 <212> DNA
 <213> Artificial Sequence

21

<220>

<223> oligonucleotide primer ZC21.224

<400> 39

acatacagga aagccttgcc caaaa

25

<210> 40

<211> 25

<212> DNA

<213> Artificial Sequence

<220>

<223> oligonucleotide primer ZC21.223

<400> 40

aaactaccct gcgattctct gctgc

25

<210> 41

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> oligonucleotide primer ZC21.334

<400> 41

ggtaaagtga gcttggctga g

21

<210> 42

<211> 3571

<212> DNA

<213> Mus musculus

<220>

<221> CDS

<222> (1049)...(2086)

<400> 42

gaattcccg	gtcgaccac	gcgtccggc	gccagggga	aaggaagctg	ggggccgcct	60
ggcggcattc	ctcgccgcag	tgtgggtcc	gtctgccgcg	ggggccgcag	tgccccctgt	120
ctgcgccagc	acctgttggc	ccgccagctg	gccgcccgcg	ccccccgcgc	ccccccgcgc	180
cgcccggccg	ccagccccgc	gccccgcgcg	ccgcccgcctg	ggggaaaagt	gagacgggga	240
ggggacaaga	gcgatcctcc	aggccagcca	ggccttcctt	tagccgcccc	tgcttagccg	300
ccacctctcc	tcagccctgc	gtcctgccct	gccttagggc	aggcatccga	gcgctcgcga	360
ctccgagccg	cccaagctct	cccggcttcc	cgcagcactt	cgccggtacc	cgagggaact	420

22

tcggtggcca ccgactgcag caaggaggag gctccgcggt ggatccgggc cagtcccagag 480
 tcgtccccgc ggccctctctg cccgcccggg acccgcgcg cactcgcagg gcacgggtccc 540
 ctccccccag gtgggggttg ggcgccgcct gccgccccga tcagcagctt tgtcattgat 600
 cccaaggtgc tcgcctcgct gccgacctgg cttccagtct ggcttggcgg gacccccagt 660
 cctcgctgt gtccgtgccc ccaaactgac aggtgctccc tgcgagtcgc cacgactcat 720
 cgccgctccc ccgcgtcccc accccttctt tctccctcg cctacccccca cccccgcac 780
 ttcggcacag ctccaggattt gtttaaactt tgggaaactg gttcagggtcc aggttttgc 840
 ttgatccctt tcaaaaactg gagacacaga agagggtctt aggaaaaact tttggatggg 900
 attatgtgga aactaccctg cgattctctg ctgccagagc cggccaggcg cttccaccgc 960
 agcgagcct tccccggct gggctgagcc ttggagtcgt cgcttcccca gtgcccgcg 1020
 cgagtgcgccc ctgcgccag tcagccaa atg ctc ctc ctc ggc ctc ctc ctg 1072

Met Leu Leu Leu Gly Leu Leu Leu

1

5

ctg aca tct gcc ctg gcc ggc caa aga acg ggg act cgg gct gag tcc 1120
 Leu Thr Ser Ala Leu Ala Gly Gln Arg Thr Gly Thr Arg Ala Glu Ser
 10 15 20

aac ctg agc agc aag ttg cag ctc tcc agc gac aag gaa cag aac gga 1168
 Asn Leu Ser Ser Lys Leu Gln Leu Ser Ser Asp Lys Glu Gln Asn Gly
 25 30 35 40

gtg caa gat ccc cgg cat gag aga gtt gtc act ata tct ggt aat ggg 1216
 Val Gln Asp Pro Arg His Glu Arg Val Val Thr Ile Ser Gly Asn Gly
 45 50 55

agc atc cac agc ccg aag ttt cct cat aca tac cca aga aat atg gtg 1264
 Ser Ile His Ser Pro Lys Phe Pro His Thr Tyr Pro Arg Asn Met Val
 60 65 70

ctg gtg tgg aga tta gtt gca gta gat gaa aat gtg cgg atc cag ctg 1312
 Leu Val Trp Arg Leu Val Ala Val Asp Glu Asn Val Arg Ile Gln Leu
 75 80 85

aca ttt gat gag aga ttt ggg ctg gaa gat cca gaa gac gat ata tgc 1360
 Thr Phe Asp Glu Arg Phe Gly Leu Glu Asp Pro Glu Asp Asp Ile Cys
 90 95 100

aag tat gat ttt gta gaa gtt gag gag ccc agt gat gga agt gtt tta 1408
 Lys Tyr Asp Phe Val Glu Val Glu Glu Pro Ser Asp Gly Ser Val Leu
 105 110 115 120

gga cgc tgg tgt ggt tct ggg act gtg cca gga aag cag act tct aaa 1456
 Gly Arg Trp Cys Gly Ser Gly Thr Val Pro Gly Lys Gln Thr Ser Lys
 125 130 135

23

gga aat cat atc agg ata aga ttt gta tct gat gag tat ttt cca tct Gly Asn His Ile Arg Ile Arg Phe Val Ser Asp Glu Tyr Phe Pro Ser 140 145 150	1504
gaa ccc gga ttc tgc atc cac tac agt att atc atg cca caa gtc aca Glu Pro Gly Phe Cys Ile His Tyr Ser Ile Ile Met Pro Gln Val Thr 155 160 165	1552
gaa acc acg agt cct tgc gtg ttg ccc cct tca tct ttg tca ttg gac Glu Thr Thr Ser Pro Ser Val Leu Pro Pro Ser Ser Leu Ser Leu Asp 170 175 180	1600
ctg ctc aac aat gct gtg act gcc ttc agt acc ttg gaa gag ctg att Leu Leu Asn Asn Ala Val Thr Ala Phe Ser Thr Leu Glu Glu Leu Ile 185 190 195 200	1648
cgg tac cta gag cca gat cga tgg cag gtg gac ttg gac agc ctc tac Arg Tyr Leu Glu Pro Asp Arg Trp Gln Val Asp Leu Asp Ser Leu Tyr 205 210 215	1696
aag cca aca tgg cag ctt ttg ggc aag gct ttc ctg tat ggg aaa aaa Lys Pro Thr Trp Gln Leu Leu Gly Lys Ala Phe Leu Tyr Gly Lys Lys 220 225 230	1744
agc aaa gtg gtg aat ctg aat ctc ctc aag gaa gag gta aaa ctc tac Ser Lys Val Val Asn Leu Asn Leu Leu Lys Glu Glu Val Lys Leu Tyr 235 240 245	1792
agc tgc aca ccc cgg aac ttc tca gtg tcc ata cgg gaa gag cta aag Ser Cys Thr Pro Arg Asn Phe Ser Val Ser Ile Arg Glu Glu Leu Lys 250 255 260	1840
agg aca gat acc ata ttc tgg cca ggt tgt ctc ctg gtc aag cgc tgt Arg Thr Asp Thr Ile Phe Trp Pro Gly Cys Leu Leu Val Lys Arg Cys 265 270 275 280	1888
gga gga aat tgt gcc tgt tgt ctc cat aat tgc aat gaa tgt cag tgt Gly Gly Asn Cys Ala Cys Cys Leu His Asn Cys Asn Glu Cys Gln Cys 285 290 295	1936
gtc cca cgt aaa gtt aca aaa aag tac cat gag gtc ctt cag ttg aga Val Pro Arg Lys Val Thr Lys Lys Tyr His Glu Val Leu Gln Leu Arg 300 305 310	1984

24

cca aaa act gga gtc aag gga ttg cat aag tca ctc act gat gtg gct 2032
 Pro Lys Thr Gly Val Lys Gly Leu His Lys Ser Leu Thr Asp Val Ala
 315 320 325

ctg gaa cac cac gag gaa tgt gac tgt gtg tgt aga gga aac gca gga 2080
 Leu Glu His His Glu Glu Cys Asp Cys Val Cys Arg Gly Asn Ala Gly
 330 335 340

ggg taa ctgcagcctt cgtagcagca cacgtgagca ctggcattct gtgtaccccc 2136
 Gly *
 345

acaagcaacc ttcatcccca ccagcgttgg ccgcagggct ctcagctgct gatgctggct 2196
 atggtaaaga tcttactcgt ctccaaccaa attctcagtt gtttgcttca atagccttcc 2256
 cctgcaggac ttcaagtgtc ttctaaaaga ccagaggcac caagaggagt caatcacaaa 2316
 gcactgcctt ctagaggaag cccagacaat ggtccttctga ccacagaaac aaatgaaatg 2376
 aatgtagatc gctagcaaac tctggagtga cagcatttct tttccactga cagaatgggtg 2436
 tagcttagtt gtcttgatat gggcaagtga tgtcagcaca agaaaatggg gaaaaacaca 2496
 cacttgattg tgaacaatgc agaaatactt ggatttctcc aacctgtttg catagataga 2556
 cagatgctct gttttctaca aactcaaagc ttttagagag cagctatgtt aataggaatt 2616
 aaatgtgcca tgctgaaagg aaagactgaa gttttcaatg cttggcaact tctccgcaat 2676
 ttggaggaaa ggtgcggtca tggtttgagg aaagcacacc tgcacagagg agtggccttc 2736
 ccttcccttc cctctgaggt ggcttctgtg tttcattgtg tatattttta tattctcctt 2796
 ttgacattat aactgttggc ttttctaata ttgttaaata tttctatttt taccaaaggt 2856
 atttaatat tttttttatg acaacctaga gcaattattt ttagcttgat aatttttttt 2916
 tctaaacaaa attgttatag ccagaagaac aaagatgatt gatataaaaa tcttgttgct 2976
 ctgacaaaaa catatgtatt tcttccttgt atggtgctag agcttagcgt catctgcatt 3036
 tgaaaagatg gaatggggaa gtttttagaa ttggtaggct gcagggacag tttgataaca 3096
 actgtactat catcaattcc caattctgtt cttagagcta cgaacagaac agagcttgag 3156
 taaatatgga gccattgcta acctaccctt ttctatggga aataggagta tagctcagag 3216
 aagcacgtcc ccagaaacct cgaccatttc taggcacagt gttctgggct atgctgcgct 3276
 gtatggacat atcctattta tttcaatact agggttttat tacctttaaa ctctgctcca 3336
 tacacttgta ttaatacatg gatattttta tgtacagaag tataatcattt aaggagtcca 3396
 cttattatac tctttggcaa ttgcaaagaa aatcaacata atacattgct tgtaaattgct 3456
 taatctgtgc ccaagttttg tgggtgactat ttgaattaaa atgtattgaa tcatcaaata 3516
 aaataatctg gctatttttg ggaaaaaaaa aaaaaaaaaa aaaaagggcg gccgc 3571

<210> 43

<211> 345

<212> PRT

<213> Mus musculus

<400> 43

Met Leu Leu Leu Gly Leu Leu Leu Leu Thr Ser Ala Leu Ala Gly Gln
 1 5 10 15

25

Arg Thr Gly Thr Arg Ala Glu Ser Asn Leu Ser Ser Lys Leu Gln Leu
 20 25 30
 Ser Ser Asp Lys Glu Gln Asn Gly Val Gln Asp Pro Arg His Glu Arg
 35 40 45
 Val Val Thr Ile Ser Gly Asn Gly Ser Ile His Ser Pro Lys Phe Pro
 50 55 60
 His Thr Tyr Pro Arg Asn Met Val Leu Val Trp Arg Leu Val Ala Val
 65 70 75 80
 Asp Glu Asn Val Arg Ile Gln Leu Thr Phe Asp Glu Arg Phe Gly Leu
 85 90 95
 Glu Asp Pro Glu Asp Asp Ile Cys Lys Tyr Asp Phe Val Glu Val Glu
 100 105 110
 Glu Pro Ser Asp Gly Ser Val Leu Gly Arg Trp Cys Gly Ser Gly Thr
 115 120 125
 Val Pro Gly Lys Gln Thr Ser Lys Gly Asn His Ile Arg Ile Arg Phe
 130 135 140
 Val Ser Asp Glu Tyr Phe Pro Ser Glu Pro Gly Phe Cys Ile His Tyr
 145 150 155 160
 Ser Ile Ile Met Pro Gln Val Thr Glu Thr Thr Ser Pro Ser Val Leu
 165 170 175
 Pro Pro Ser Ser Leu Ser Leu Asp Leu Leu Asn Asn Ala Val Thr Ala
 180 185 190
 Phe Ser Thr Leu Glu Glu Leu Ile Arg Tyr Leu Glu Pro Asp Arg Trp
 195 200 205
 Gln Val Asp Leu Asp Ser Leu Tyr Lys Pro Thr Trp Gln Leu Leu Gly
 210 215 220
 Lys Ala Phe Leu Tyr Gly Lys Lys Ser Lys Val Val Asn Leu Asn Leu
 225 230 235 240
 Leu Lys Glu Glu Val Lys Leu Tyr Ser Cys Thr Pro Arg Asn Phe Ser
 245 250 255
 Val Ser Ile Arg Glu Glu Leu Lys Arg Thr Asp Thr Ile Phe Trp Pro
 260 265 270
 Gly Cys Leu Leu Val Lys Arg Cys Gly Gly Asn Cys Ala Cys Cys Leu
 275 280 285
 His Asn Cys Asn Glu Cys Gln Cys Val Pro Arg Lys Val Thr Lys Lys
 290 295 300
 Tyr His Glu Val Leu Gln Leu Arg Pro Lys Thr Gly Val Lys Gly Leu
 305 310 315 320
 His Lys Ser Leu Thr Asp Val Ala Leu Glu His His Glu Glu Cys Asp
 325 330 335
 Cys Val Cys Arg Gly Asn Ala Gly Gly
 340 345

<210> 44

<211> 65

26

<212> DNA

<213> Artificial Sequence

<220>

<223> oligonucleotide primer ZC20.572

<400> 44

tcaccacgcg aattcggtag cgctggttcc gcgtggatcc ggccagagac aggggactca 60
ggcgg 65

<210> 45

<211> 65

<212> DNA

<213> Artificial Sequence

<220>

<223> oligonucleotide primer ZC20.573

<400> 45

tctgtatcag gctgaaaatc ttatctcatc cgccaaaaca ctatcctcct gtgctccctc 60
tgcac 65

<210> 46

<211> 40

<212> DNA

<213> Artificial Sequence

<220>

<223> oligonucleotide primer ZC19.372

<400> 46

tgatgatgaa gccctgaaag acgcgcagac taattcgagc 40

<210> 47

<211> 60

<212> DNA

<213> Artificial Sequence

<220>

<223> oligonucleotide primer ZC19.351

<400> 47

acgcgcagac taattcgagc tcccaccatc accatcacca cgcgaattcg gtaccgctgg 60

<210> 48

27

<211> 60
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> oligonucleotide primer ZC19.352

<400> 48
 actcactata gggcgaattg cccgggggat ccacgcggaa ccagcggtac cgaattcgcg 60

<210> 49
 <211> 42
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> oligonucleotide primer ZC19.371

<400> 49
 acggccagtg aattgtaata cgactcacta tagggcgaat tg 42

<210> 50
 <211> 1095
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Fused DNA

<400> 50
 ctgaaagacg cgcagactaa ttcgagctcc caccatcacc atcaccacgc gaattcggta 60
 ccgctgggtc cgcgtggatc cggccagaga caggggactc aggcggaatc caacctgagt 120
 agtaaattcc agttttccag caacaaggaa cagaacggag tacaagatcc tcagcatgag 180
 agaattatta ctgtgtctac taatggaagt attcacagcc caaggtttcc tcatacttat 240
 ccaagaaata cggctcttgg atggagatta gtagcagtag aggaaaatgt atggatacaa 300
 cttacgtttg atgaaagatt tgggcttgaa gaccagaag atgacatatg caagtatgat 360
 tttgtagaag ttgaggaacc cagtgatgga actatattag ggcgctggtg tggttctggt 420
 actgtaccag gaaaacagat ttctaaagga aatcaaatta ggataagatt tgtatctgat 480
 gaatatattc cttctgaacc agggttctgc atccactaca acattgtcat gccacaattc 540
 acagaagctg tgagtccttc agtgctaccc cttcagctt tgccactgga cctgcttaat 600
 aatgctataa ctgcctttag taccttgga gaccttattc gatatcttga accagagaga 660
 tggcagttgg acttagaaga tctatatagg ccaacttggc aacttcttgg caaggctttt 720
 gtttttgga gaaaatccag agtggtggat ctgaaccttc taacagagga ggtaagatta 780
 tacagctgca cacctcgtaa cttctcagtg tccataaggg aagaactaaa gagaaccgat 840
 accattttct ggccaggttg tctcctggtt aaacgctgtg gtgggaactg tgcctgttgt 900

ctccacaatt	gcaatgaatg	tcaatgtgtc	ccaagcaaag	ttactaaaaa	ataccacgag	960
gtccttcagt	tgagaccaa	gaccggtgtc	aggggattgc	acaaatcact	caccgacgtg	1020
gccctggagc	accatgagga	gtgtgactgt	gtgtgcagag	ggagcacagg	aggatagtgt	1080
tttggcggat	gagat					1095

THIS PAGE BLANK (USPTO)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
15 June 2000 (15.06.2000)

PCT

(10) International Publication Number
WO 00/34474 A3

(51) International Patent Classification⁷: **C12N 15/12**,
15/11, 5/10, 15/62, C07K 14/52, 16/22, G01N 33/68,
C12Q 1/68, A61K 38/18

Seattle, WA 98155 (US). **WEST, James, W.**; 7539 19th
Northeast, Seattle, WA 98115 (US).

(21) International Application Number: **PCT/US99/28968**

(74) Agent: **PARKER, Gary, E.**; ZymoGenetics, Inc., 1201
Eastlake Avenue East, Seattle, WA 98102 (US).

(22) International Filing Date: 7 December 1999 (07.12.1999)

(81) Designated States (*national*): AE, AL, AM, AT, AU, AZ,
BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK,
DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL,
IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU,
LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO,
RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG,
UZ, VN, YU, ZA, ZW.

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

09/207,120	7 December 1998 (07.12.1998)	US
60/142,576	6 July 1999 (06.07.1999)	-US
60/161,653	21 October 1999 (21.10.1999)	US
60/165,255	12 November 1999 (12.11.1999)	US

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent
(AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent
(AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU,
MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM,
GA, GN, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant: **ZYMOGENETICS, INC.** [US/US]; 1201
Eastlake Avenue East, Seattle, WA 98102 (US).

(72) Inventors: **GAO, Zeren**; 9502 179th Place Northeast
#3; Redmond, WA 98052 (US). **HART, Charles, E.**;
20114 183rd Place Northeast, Woodinville, WA 98072
(US). **PIDDINGTON, Christopher, S.**; 2838 Irongate
Place, Thousand Oaks, CA 91362 (US). **SHEPPARD,**
Paul, O.; 13532 278th Drive Northeast, Granite Falls,
WA 98252 (US). **SHOEMAKER, Kimberly, E.**; 4634
Somerset Avenue Southeast, Bellevue, WA 98006 (US).
GILBERTSON, Debra, G.; 4025 Northeast 195th Place,

Published:

— With international search report.

(88) Date of publication of the international search report: 28 December 2000

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: **GROWTH FACTOR HOMOLOG ZVEGF3**

(57) Abstract: Polypeptide growth factors, methods of making them, polynucleotides encoding them, antibodies to them, and methods of using them are disclosed. The polypeptides comprises an amino acid segment that is at least 90% identical to residues 46-163 of SEQ ID NO:2 or residues 235-345 of SEQ ID NO:2. Multimers of the polypeptides are also disclosed. The polypeptides, multimeric proteins, and polynucleotides can be used in the study and regulation of cell and tissue development, as components of cell culture media, and as diagnostic agents.

WO 00/34474 A3

INTERNATIONAL SEARCH REPORT

Intern. Appl. No.

PCT/US 99/28968

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N15/12 C12N15/11 C12N5/10 C12N15/62 C07K14/52
C07K16/22 G01N33/68 C12Q1/68 A61K38/18

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N C07K G01N C12Q A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, STRAND, BIOSIS

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P, X	WO 99 47677 A (GENENTECH INC) 23 September 1999 (1999-09-23) figure 2	1-48
P, X	WO 99 37671 A (NA SONGQING ; SONG HO YEONG (US); DOU SHENSHEN (US); LILLY CO ELI () 29 July 1999 (1999-07-29) page 56 -page 58	1-48
E	EP 0 984 063 A (LILLY CO ELI) 8 March 2000 (2000-03-08) the whole document	1-48
A	WO 98 07832 A (UNIV HELSINKI LICENSING ; LUDWIG INST CANCER RES (US)) 26 February 1998 (1998-02-26) abstract seq. ID 2	1-48



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

26 June 2000

Date of mailing of the international search report

03/07/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Galli, I

INTERNATIONAL SEARCH REPORT

Intern. Application No

PCT/US 99/28968

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>WO 98 24811 A (ZYMOGENETICS INC) 11 June 1998 (1998-06-11) abstract seq. ID 2</p> <p>-----</p>	1-48

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 99/28968

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9947677 A	23-09-1999	AU 3072199 A AU 3075099 A WO 9946281 A	27-09-1999 11-10-1999 16-09-1999
WO 9937671 A	29-07-1999	AU 2562199 A EP 0984063 A	09-08-1999 08-03-2000
EP 0984063 A	08-03-2000	AU 2562199 A WO 9937671 A	09-08-1999 29-07-1999
WO 9807832 A	26-02-1998	AU 4079697 A EP 0956339 A	06-03-1998 17-11-1999
WO 9824811 A	11-06-1998	AU 5439998 A	29-06-1998

CORRECTED VERSION

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
15 June 2000 (15.06.2000)

(10) International Publication Number
WO 00/034474 A3

PCT

(51) International Patent Classification⁷: **C12N 15/12**,
15/11, 5/10, 15/62, C07K 14/52, 16/22, G01N 33/68,
C12Q 1/68, A61K 38/18

(74) Agent: **PARKER, Gary, E.**; ZymoGenetics, Inc., 1201
Eastlake Avenue East, Seattle, WA 98102 (US).

(21) International Application Number: **PCT/US99/28968**

(22) International Filing Date: 7 December 1999 (07.12.1999)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
09/207,120 7 December 1998 (07.12.1998) US
60/142,576 6 July 1999 (06.07.1999) US
60/161,653 21 October 1999 (21.10.1999) US
60/165,255 12 November 1999 (12.11.1999) US

(81) Designated States (*national*): AE, AL, AM, AT, AU, AZ,
BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK,
DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL,
IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU,
LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO,
RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG,
UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent
(AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent
(AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU,
MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM,
GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

(71) Applicant: **ZYMOGENETICS, INC.** [US/US]; 1201
Eastlake Avenue East, Seattle, WA 98102 (US).

(88) Date of publication of the international search report:
28 December 2000

(72) Inventors: **GAO, Zeren**; 9502 179th Place Northeast
#3, Redmond, WA 98052 (US). **HART, Charles, E.**;
20114 183rd Place Northeast, Woodinville, WA 98072
(US). **PIDDINGTON, Christopher, S.**; 2838 Irongate
Place, Thousand Oaks, CA 91362 (US). **SHEPPARD,
Paul, O.**; 13532 278th Drive Northeast, Granite Falls,
WA 98252 (US). **SHOEMAKER, Kimberly, E.**; 4634
Somerset Avenue Southeast, Bellevue, WA 98006 (US).
GILBERTSON, Debra, G.; 4025 Northeast 195th Place,
Seattle, WA 98155 (US). **WEST, James, W.**; 7539 19th
Northeast, Seattle, WA 98115 (US).

(48) Date of publication of this corrected version:
29 August 2002

(15) Information about Correction:
see PCT Gazette No. 35/2002 of 29 August 2002, Section
II

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: **GROWTH FACTOR HOMOLOG ZVEGF3**

(57) Abstract: Polypeptide growth factors, methods of making them, polynucleotides encoding them, antibodies to them, and meth-
ods of using them are disclosed. The polypeptides comprises an amino acid segment that is at least 90% identical to residues 46-163
of SEQ ID NO:2 or residues 235-345 of SEQ ID NO:2. Multimers of the polypeptides are also disclosed. The polypeptides, mul-
timeric proteins, and polynucleotides can be used in the study and regulation of cell and tissue development, as components of cell
culture media, and as diagnostic agents.

WO 00/034474 A3

Description

GROWTH FACTOR HOMOLOG ZVEGF3

5

BACKGROUND OF THE INVENTION

In multicellular animals, cell growth, differentiation, and migration are controlled by polypeptide growth factors. These growth factors play a role in both normal development and pathogenesis, including the development of solid tumors.

Polypeptide growth factors influence cellular events by binding to cell-surface receptors, many of which are tyrosine kinases. Binding initiates a chain of signalling events within the cell, which ultimately results in phenotypic changes, such as cell division, protease production, and cell migration.

Growth factors can be classified into families on the basis of structural similarities. One such family, the PDGF (platelet derived growth factor) family, is characterized by a dimeric structure stabilized by disulfide bonds. This family includes PDGF, the placental growth factors (PIGFs), and the vascular endothelial growth factors (VEGFs). The individual polypeptide chains of these proteins form characteristic higher-order structures having a bow tie-like configuration about a cystine knot, formed by disulfide bonding between pairs of cysteine residues. Hydrophobic interactions between loops contribute to the dimerization of the two monomers. See, Daopin et al., *Science* 257:369, 1992; Lapthorn et al., *Nature* 369:455, 1994. Members of this family are active as both homodimers and heterodimers. See, for example, Heldin et al., *EMBO J.* 7:1387-1393, 1988; Cao et al., *J. Biol. Chem.* 271:3154-3162, 1996. The cystine knot motif and bow tie fold are also characteristic of the growth factors transforming growth factor-beta (TGF- β) and nerve growth factor (NGF), and the glycoprotein hormones. Although their amino acid sequences are quite divergent, these proteins all contain the six conserved cysteine residues of the cystine knot.